

Juan Marzoa Fandiño

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/893219/publications.pdf>

Version: 2024-02-01

16

papers

532

citations

759233

12

h-index

940533

16

g-index

16

all docs

16

docs citations

16

times ranked

802

citing authors

#	ARTICLE	IF	CITATIONS
1	Impact of human-associated Escherichia coli clonal groups in Antarctic pinnipeds: presence of ST73, ST95, ST141 and ST131. <i>Scientific Reports</i> , 2018, 8, 4678.	3.3	37
2	Liposomes or traditional adjuvants: induction of bactericidal activity by the macrophage infectivity potentiator protein (Mip) of <i>Neisseria meningitidis</i> . <i>Apmis</i> , 2017, 125, 725-731.	2.0	5
3	Lambs are an important source of atypical enteropathogenic Escherichia coli in southern Brazil. <i>Veterinary Microbiology</i> , 2016, 196, 72-77.	1.9	15
4	Diversity of Shiga toxin-producing Escherichia coli in sheep flocks of Paraná State, southern Brazil. <i>Veterinary Microbiology</i> , 2015, 175, 150-156.	1.9	34
5	Virulence Patterns in a Murine Sepsis Model of ST131 Escherichia coli Clinical Isolates Belonging to Serotypes O25b:H4 and O16:H5 Are Associated to Specific Viotypes. <i>PLoS ONE</i> , 2014, 9, e87025.	2.5	44
6	Molecular epidemiology and virulence of Escherichia coli O16:H5-ST131: Comparison with H30 and H30-Rx subclones of O25b:H4-ST131. <i>International Journal of Medical Microbiology</i> , 2014, 304, 1247-1257.	3.6	64
7	Poultry as reservoir for extraintestinal pathogenic Escherichia coli O45:K1:H7-B2-ST95 in humans. <i>Veterinary Microbiology</i> , 2013, 167, 506-512.	1.9	87
8	High Resolution Clear Native Electrophoresis (hrCNE) Allows a Detailed Analysis of the Heterotrimeric Structure of Recombinant <i>Neisseria meningitidis</i> Porins Inserted into Liposomes. <i>Journal of Proteome Research</i> , 2013, 12, 777-784.	3.7	6
9	Emergence of new variants of ST131 clonal group among extraintestinal pathogenic Escherichia coli producing extended-spectrum β -lactamases. <i>International Journal of Antimicrobial Agents</i> , 2013, 42, 347-351.	2.5	55
10	Detection of quinolone-resistant Escherichia coli isolates belonging to clonal groups O25b:H4-B2-ST131 and O25b:H4-D-ST69 in raw sewage and river water in Barcelona, Spain. <i>Journal of Antimicrobial Chemotherapy</i> , 2013, 68, 758-765.	3.0	44
11	Four Main Viotypes among Extended-Spectrum- β -Lactamase-Producing Isolates of Escherichia coli O25b:H4-B2-ST131: Bacterial, Epidemiological, and Clinical Characteristics. <i>Journal of Clinical Microbiology</i> , 2013, 51, 3358-3367.	3.9	76
12	Induction of immune responses by purified outer membrane protein complexes from <i>Neisseria meningitidis</i> . <i>Vaccine</i> , 2012, 30, 2387-2395.	3.8	13
13	Identification of <i>Neisseria meningitidis</i> Outer Membrane Vesicle Complexes Using 2-D High Resolution Clear Native/SDS-PAGE. <i>Journal of Proteome Research</i> , 2010, 9, 611-619.	3.7	17
14	Analysis of outer membrane porin complexes of <i>Neisseria meningitidis</i> in wild-type and specific knock-out mutant strains. <i>Proteomics</i> , 2009, 9, 648-656.	2.2	17
15	Characterisation and immune responses to meningococcal recombinant porin complexes incorporated into liposomes. <i>Vaccine</i> , 2009, 27, 5338-5343.	3.8	7
16	A cross-reactive neisserial antigen encoded by the NMB0035 locus shows high sequence conservation but variable surface accessibility. <i>Journal of Medical Microbiology</i> , 2008, 57, 80-87.	1.8	11