Maria Papp

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/8934492/maria-papp-publications-by-year.pdf

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

102
papers2,416
citations29
h-index45
g-index111
ext. papers3,034
ext. citations3.7
avg, IF4.37
L-index

#	Paper	IF	Citations
102	current acute pancreatitis prevention by the elimination of alcohol and cigaette smoking (REAPPEAR): protocol of a randomised controlled trial and a cohort study <i>BMJ Open</i> , 2022 , 12, e05082	1 3	3
101	In-Hospital Patient Education Markedly Reduces Alcohol Consumption after Alcohol-Induced Acute Pancreatitis. <i>Nutrients</i> , 2022 , 14, 2131	6.7	0
100	Serological biomarkers for management of primary sclerosing cholangitis. <i>World Journal of Gastroenterology</i> , 2022 , 28, 2291-2301	5.6	O
99	EASY-APP: An artificial intelligence model and application for early and easy prediction of severity in acute pancreatitis. <i>Clinical and Translational Medicine</i> , 2022 , 12,	5.7	1
98	Hypoalbuminemia affects one third of acute pancreatitis patients and is independently associated with severity and mortality <i>Scientific Reports</i> , 2021 , 11, 24158	4.9	2
97	Early infection is an independent risk factor for increased mortality in patients with culture-confirmed infected pancreatic necrosis. <i>Pancreatology</i> , 2021 ,	3.8	1
96	Abnormal ferritin levels predict development of poor outcomes in cirrhotic outpatients: a cohort study. <i>BMC Gastroenterology</i> , 2021 , 21, 94	3	O
95	Glucose levels show independent and dose-dependent association with worsening acute pancreatitis outcomes: Post-hoc analysis of a prospective, international cohort of 2250 acute pancreatitis cases. <i>Pancreatology</i> , 2021 , 21, 1237-1246	3.8	1
94	Acute-on-chronic liver failure: A distinct clinical syndrome. <i>Journal of Hepatology</i> , 2021 , 75 Suppl 1, S27-	·SB;54	7
93	PREDICT identifies precipitating events associated with the clinical course of acutely decompensated cirrhosis. <i>Journal of Hepatology</i> , 2021 , 74, 1097-1108	13.4	41
92	Evidence for diagnosis of early chronic pancreatitis after three episodes of acute pancreatitis: a cross-sectional multicentre international study with experimental animal model. <i>Scientific Reports</i> , 2021 , 11, 1367	4.9	8
91	Initial Renal Function (eGFR) Is a Prognostic Marker of Severe Acute Pancreatitis: A Cohort-Analysis of 1,224 Prospectively Collected Cases. <i>Frontiers in Medicine</i> , 2021 , 8, 671917	4.9	О
90	Increased sTREM-1 levels identify cirrhotic patients with bacterial infection and predict their 90-day mortality. <i>Clinics and Research in Hepatology and Gastroenterology</i> , 2021 , 45, 101579	2.4	3
89	Early occurrence of pseudocysts in acute pancreatitis - A multicenter international cohort analysis of 2275 cases. <i>Pancreatology</i> , 2021 ,	3.8	3
88	Anti-neutrophil cytoplasmic antibody testing by indirect immunofluorescence: Computer-aided versus conventional microscopic evaluation of routine diagnostic samples from patients with vasculitis or other inflammatory diseases. <i>Clinica Chimica Acta</i> , 2020 , 511, 117-124	6.2	1
87	Hypertriglyceridemia-induced acute pancreatitis: A prospective, multicenter, international cohort analysis of 716 acute pancreatitis cases. <i>Pancreatology</i> , 2020 , 20, 608-616	3.8	29
86	High discontinuation rate of azathioprine in autoimmune hepatitis, independent of time of treatment initiation. <i>Liver International</i> , 2020 , 40, 2164-2171	7.9	8

(2019-2020)

85	LIFEStyle, Prevention and Risk of Acute PaNcreatitis (LIFESPAN): protocol of a multicentre and multinational observational case-control study. <i>BMJ Open</i> , 2020 , 10, e029660	3	3
84	Development of disturbance of consciousness is associated with increased severity in acute pancreatitis. <i>Pancreatology</i> , 2020 , 20, 806-812	3.8	2
83	The search for the Holy Grail: autoantigenic targets in primary sclerosing cholangitis associated with disease phenotype and neoplasia. <i>Autoimmunity Highlights</i> , 2020 , 11, 6	3.7	2
82	Golgi protein-73: A biomarker for assessing cirrhosis and prognosis of liver disease patients. <i>World Journal of Gastroenterology</i> , 2020 , 26, 5130-5145	5.6	11
81	Rapid Response to Treatment of Autoimmune Hepatitis Associated With Remission at 6 and 12 Months. <i>Clinical Gastroenterology and Hepatology</i> , 2020 , 18, 1609-1617.e4	6.9	8
80	EarLy Elimination of Fatty Acids iN hypertriglyceridemia-induced acuTe pancreatitis (ELEFANT trial): Protocol of an open-label, multicenter, adaptive randomized clinical trial. <i>Pancreatology</i> , 2020 , 20, 369-	3 7 8	15
79	Deleterious effect of proton pump inhibitors on the disease course of cirrhosis. <i>European Journal of Gastroenterology and Hepatology</i> , 2020 , 32, 257-264	2.2	15
78	Real-life efficacy of vedolizumab on endoscopic healing in inflammatory bowel disease - A nationwide Hungarian cohort study. <i>Expert Opinion on Biological Therapy</i> , 2020 , 20, 205-213	5.4	4
77	Acid suppression therapy, gastrointestinal bleeding and infection in acute pancreatitis - An international cohort study. <i>Pancreatology</i> , 2020 , 20, 1323-1331	3.8	6
76	The PREDICT study uncovers three clinical courses of acutely decompensated cirrhosis that have distinct pathophysiology. <i>Journal of Hepatology</i> , 2020 , 73, 842-854	13.4	108
76 75		13.4 4.6	108
	A Multicenter, International Cohort Analysis of 1435 Cases to Support Clinical Trial Design in Acute		
75	A Multicenter, International Cohort Analysis of 1435 Cases to Support Clinical Trial Design in Acute Pancreatitis. <i>Frontiers in Physiology</i> , 2019 , 10, 1092 Antibiotic therapy in acute pancreatitis: From global overuse to evidence based recommendations.	4.6	15
75 74	A Multicenter, International Cohort Analysis of 1435 Cases to Support Clinical Trial Design in Acute Pancreatitis. <i>Frontiers in Physiology</i> , 2019 , 10, 1092 Antibiotic therapy in acute pancreatitis: From global overuse to evidence based recommendations. <i>Pancreatology</i> , 2019 , 19, 488-499 Predniso(lo)ne Dosage and Chance of Remission in Patients With Autoimmune Hepatitis. <i>Clinical</i>	4.6	15 31
75 74 73	A Multicenter, International Cohort Analysis of 1435 Cases to Support Clinical Trial Design in Acute Pancreatitis. Frontiers in Physiology, 2019, 10, 1092 Antibiotic therapy in acute pancreatitis: From global overuse to evidence based recommendations. Pancreatology, 2019, 19, 488-499 Predniso(lo)ne Dosage and Chance of Remission in Patients With Autoimmune Hepatitis. Clinical Gastroenterology and Hepatology, 2019, 17, 2068-2075.e2 HLA-DQ2 homozygosis increases tTGA levels at diagnosis but does not influence the clinical phenotype of coeliac disease: A multicentre study. International Journal of Immunogenetics, 2019,	4.6 3.8 6.9	15 31 31
75 74 73 72	A Multicenter, International Cohort Analysis of 1435 Cases to Support Clinical Trial Design in Acute Pancreatitis. Frontiers in Physiology, 2019, 10, 1092 Antibiotic therapy in acute pancreatitis: From global overuse to evidence based recommendations. Pancreatology, 2019, 19, 488-499 Predniso(lo)ne Dosage and Chance of Remission in Patients With Autoimmune Hepatitis. Clinical Gastroenterology and Hepatology, 2019, 17, 2068-2075.e2 HLA-DQ2 homozygosis increases tTGA levels at diagnosis but does not influence the clinical phenotype of coeliac disease: A multicentre study. International Journal of Immunogenetics, 2019, 46, 74-81 A DNA pool of FLT3-ITD positive DNA samples can be used efficiently for analytical evaluation of NGS-based FLT3-ITD quantitation - Testing several different ITD sequences and rates,	4.6 3.8 6.9	15 31 31 8
75 74 73 72 71	A Multicenter, International Cohort Analysis of 1435 Cases to Support Clinical Trial Design in Acute Pancreatitis. Frontiers in Physiology, 2019, 10, 1092 Antibiotic therapy in acute pancreatitis: From global overuse to evidence based recommendations. Pancreatology, 2019, 19, 488-499 Predniso(lo)ne Dosage and Chance of Remission in Patients With Autoimmune Hepatitis. Clinical Gastroenterology and Hepatology, 2019, 17, 2068-2075.e2 HLA-DQ2 homozygosis increases tTGA levels at diagnosis but does not influence the clinical phenotype of coeliac disease: A multicentre study. International Journal of Immunogenetics, 2019, 46, 74-81 A DNA pool of FLT3-ITD positive DNA samples can be used efficiently for analytical evaluation of NGS-based FLT3-ITD quantitation - Testing several different ITD sequences and rates, simultaneously. Journal of Biotechnology, 2019, 303, 25-29 Multiple Hits in Acute Pancreatitis: Components of Metabolic Syndrome Synergize Each Other's	4.6 3.8 6.9 2.3	15 31 31 8

67	Observational longitudinal multicentre investigation of acute pancreatitis (GOULASH PLUS): follow-up of the GOULASH study, protocol. <i>BMJ Open</i> , 2019 , 9, e025500	3	2
66	Multidrug-resistant bacterial infections in patients with decompensated cirrhosis and with acute-on-chronic liver failure in Europe. <i>Journal of Hepatology</i> , 2019 , 70, 398-411	13.4	120
65	Aging and Comorbidities in Acute Pancreatitis II.: A Cohort-Analysis of 1203 Prospectively Collected Cases. <i>Frontiers in Physiology</i> , 2018 , 9, 1776	4.6	20
64	Loss of tolerance to gut immunity protein, glycoprotein 2 (GP2) is associated with progressive disease course in primary sclerosing cholangitis. <i>Scientific Reports</i> , 2018 , 8, 399	4.9	11
63	Significance of serological markers in the disease course of ulcerative colitis in a prospective clinical cohort of patients. <i>PLoS ONE</i> , 2018 , 13, e0194166	3.7	14
62	Functional polymorphisms of innate immunity receptors are not risk factors for the non-SBP type bacterial infections in cirrhosis. <i>Liver International</i> , 2018 , 38, 1242-1252	7.9	6
61	Mucosal Autoimmunity to Cell-Bound GP2 Isoforms Is a Sensitive Marker in PSC and Associated With the Clinical Phenotype. <i>Frontiers in Immunology</i> , 2018 , 9, 1959	8.4	8
60	Major Changes of von Willebrand Factor Multimer Distribution in Cirrhotic Patients with Stable Disease or Acute Decompensation. <i>Thrombosis and Haemostasis</i> , 2018 , 118, 1397-1408	7	8
59	Lectin-complement pathway molecules are decreased in patients with cirrhosis and constitute the risk of bacterial infections. <i>Liver International</i> , 2017 , 37, 1023-1031	7.9	17
58	Prediction of Short- and Medium-term Efficacy of Biosimilar Infliximab Therapy. Do Trough Levels and Antidrug Antibody Levels or Clinical And Biochemical Markers Play the More Important Role?. <i>Journal of Crohnps and Colitis</i> , 2017 , 11, 697-705	1.5	25
57	High versus low energy administration in the early phase of acute pancreatitis (GOULASH trial): protocol of a multicentre randomised double-blind clinical trial. <i>BMJ Open</i> , 2017 , 7, e015874	3	17
56	Long-term Efficacy, Safety, and Immunogenicity of Biosimilar Infliximab After One Year in a Prospective Nationwide Cohort. <i>Inflammatory Bowel Diseases</i> , 2017 , 23, 1908-1915	4.5	36
55	Response to Low L-Ficolin associated with disease severity during sepsis in adult ICU patients. <i>Liver International</i> , 2017 , 37, 1410	7.9	1
54	Gut barrier failure biomarkers are associated with poor disease outcome in patients with primary sclerosing cholangitis. <i>World Journal of Gastroenterology</i> , 2017 , 23, 5412-5421	5.6	23
53	Efficacy and Safety of the Biosimilar Infliximab CT-P13 Treatment in Inflammatory Bowel Diseases: A Prospective, Multicentre, Nationwide Cohort. <i>Journal of Crohnps and Colitis</i> , 2016 , 10, 133-40	1.5	133
52	Presepsin teardown - pitfalls of biomarkers in the diagnosis and prognosis of bacterial infection in cirrhosis. <i>World Journal of Gastroenterology</i> , 2016 , 22, 9172-9185	5.6	12
51	Mannose-Binding Lectin Levels and Carotid Intima-Media Thickness in Type 2 Diabetic Patients. Journal of Diabetes Research, 2016 , 2016, 8132925	3.9	8
50	Baseline Characteristics and Disease Phenotype in Inflammatory Bowel Disease. <i>Journal of Pediatric Gastroenterology and Nutrition</i> , 2016 , 62, 50-5	2.8	10

(2012-2016)

49	Macrophage activation marker, soluble CD163, is an independent predictor of short-term mortality in patients with cirrhosis and bacterial infection. <i>Liver International</i> , 2016 , 36, 1628-1638	7.9	12
48	Seasonal variability of vitamin D and bone metabolism in infliximab-treated paediatric Crohn's disease. <i>Digestive and Liver Disease</i> , 2015 , 47, 652-7	3.3	6
47	Diagnostic and clinical significance of Crohn's disease-specific anti-MZGP2 pancreatic antibodies by a novel ELISA. <i>Clinica Chimica Acta</i> , 2015 , 441, 176-81	6.2	19
46	Risk Matrix for Prediction of Disease Progression in a Referral Cohort of Patients with Crohn's Disease. <i>Journal of Crohnys and Colitis</i> , 2015 , 9, 891-8	1.5	8
45	Rediscovery of the Anti-Pancreatic Antibodies and Evaluation of their Prognostic Value in a Prospective Clinical Cohort of Crohn's Patients: The Importance of Specific Target Antigens [GP2 and CUZD1]. <i>Journal of Crohnps and Colitis</i> , 2015 , 9, 659-68	1.5	30
44	Prevalence, significance and predictive value of antiphospholipid antibodies in Crohn's disease. <i>World Journal of Gastroenterology</i> , 2015 , 21, 6952-64	5.6	14
43	Autoregressive cross-lagged models of IMPACT-III and Pediatric Crohn's Disease Activity Indexes during one year infliximab therapy in pediatric patients with Crohn's disease. <i>Journal of Crohn's and Colitis</i> , 2014 , 8, 747-55	1.5	9
42	New serological markers in pediatric patients with inflammatory bowel disease. <i>World Journal of Gastroenterology</i> , 2014 , 20, 4873-82	5.6	16
41	Immune dysfunction in cirrhosis. World Journal of Gastroenterology, 2014, 20, 2564-77	5.6	105
40	Serological studies in inflammatory bowel disease: how important are they?. <i>Current Opinion in Gastroenterology</i> , 2014 , 30, 359-64	3	18
39	Incidence and paris classification of pediatric inflammatory bowel disease. <i>Gastroenterology Research and Practice</i> , 2014 , 2014, 904307	2	32
38	Pathophysiological and clinical approach to cirrhotic cardiomyopathy. <i>Journal of Gastrointestinal and Liver Diseases</i> , 2014 , 23, 301-10	1.4	16
37	Predictors of relapse in patients with ulcerative colitis in remission after one-year of infliximab therapy. <i>Scandinavian Journal of Gastroenterology</i> , 2013 , 48, 1394-8	2.4	41
36	Low mannose-binding lectin (MBL) is associated with paediatric inflammatory bowel diseases and ileal involvement in patients with Crohn disease. <i>Journal of Crohnps and Colitis</i> , 2013 , 7, 134-41	1.5	8
35	High prevalence of IgA class anti-neutrophil cytoplasmic antibodies (ANCA) is associated with increased risk of bacterial infection in patients with cirrhosis. <i>Journal of Hepatology</i> , 2013 , 59, 457-66	13.4	27
34	Incidence, Paris classification, and follow-up in a nationwide incident cohort of pediatric patients with inflammatory bowel disease. <i>Journal of Pediatric Gastroenterology and Nutrition</i> , 2013 , 57, 576-82	2.8	75
33	Foreign body impaction in the sigmoid colon: a twenty euro bet. <i>World Journal of Gastroenterology</i> , 2013 , 19, 3892-4	5.6	9
32	Association of peroxisome proliferator-activated receptor gamma polymorphisms with inflammatory bowel disease in a Hungarian cohort. <i>Inflammatory Bowel Diseases</i> , 2012 , 18, 472-9	4.5	12

31	Acute phase proteins in the diagnosis and prediction of cirrhosis associated bacterial infections. Liver International, 2012 , 32, 603-11	7.9	67
30	Diagnostic yield of upper endoscopy in paediatric patients with Crohn's disease and ulcerative colitis. Subanalysis of the HUPIR registry. <i>Journal of Crohnps and Colitis</i> , 2012 , 6, 86-94	1.5	29
29	High-sensitivity C-reactive protein for identification of disease phenotype, active disease, and clinical relapses in Crohn's disease: a marker for patient classification?. <i>Inflammatory Bowel Diseases</i> , 2012 , 18, 1647-54	4.5	58
28	Is there any association between impaired health-related quality of life and non-adherence to medical therapy in inflammatory bowel disease?. <i>Scandinavian Journal of Gastroenterology</i> , 2012 , 47, 1298-303	2.4	13
27	Pancreatic autoantibodies and autoantibodies against goblet cells in pediatric patients with inflammatory bowel disease. <i>Journal of Pediatric Gastroenterology and Nutrition</i> , 2012 , 55, 429-35	2.8	26
26	Intestinal alkaline phosphatase in the colonic mucosa of children with inflammatory bowel disease. <i>World Journal of Gastroenterology</i> , 2012 , 18, 3254-9	5.6	37
25	Increased levels of platelet activation markers are positively associated with carotid wall thickness and other atherosclerotic risk factors in obese patients. <i>Thrombosis and Haemostasis</i> , 2011 , 106, 683-92	7	49
24	Serologic antiglycan antibodies in inflammatory bowel disease. <i>American Journal of Gastroenterology</i> , 2011 , 106, 406-12	0.7	31
23	Phenotypic polymorphism of haptoglobin: a novel risk factor for the development of infection in liver cirrhosis. <i>Human Immunology</i> , 2011 , 72, 348-54	2.3	9
22	Serum anti-Saccharomyces cerevisiae antibodies in Greek patients with Behcet's disease. <i>Yonsei Medical Journal</i> , 2011 , 52, 347-50	3	2
21	Cytokine (IL-10, IL-28B and LT-A) gene polymorphisms in chronic hepatitis C virus infection. <i>Orvosi Hetilap</i> , 2011 , 5, 9-19		10
20	Serum lipopolysaccharide-binding protein and soluble CD14 are markers of disease activity in patients with Crohn's disease. <i>Inflammatory Bowel Diseases</i> , 2011 , 17, 767-77	4.5	51
19	Duodenal ulceration in a patient with celiac disease and plasminogen I deficiency: coincidence or cofactors?. <i>Pediatrics</i> , 2011 , 128, e1302-6	7.4	4
18	A New Approach to Predict the Chance of Infectious Complications Following Autologous Hematopoietic Stem Cell Transplantation: Mannose-Binding Lectin ELISA. <i>Blood</i> , 2011 , 118, 4492-4492	2.2	
17	Presence of anti-microbial antibodies in liver cirrhosisa tell-tale sign of compromised immunity?. <i>PLoS ONE</i> , 2010 , 5, e12957	3.7	38
16	Association of adherence to therapy and complementary and alternative medicine use with demographic factors and disease phenotype in patients with inflammatory bowel disease. <i>Journal of Crohnps and Colitis</i> , 2010 , 4, 283-90	1.5	36
15	Mannose-binding lectin level and deficiency is not associated with inflammatory bowel diseases, disease phenotype, serology profile, and NOD2/CARD15 genotype in a large Hungarian cohort. <i>Human Immunology</i> , 2010 , 71, 407-13	2.3	11
14	Mannose-binding lectin deficiency confers risk for bacterial infections in a large Hungarian cohort of patients with liver cirrhosis. <i>Journal of Hepatology</i> , 2010 , 53, 484-91	13.4	22

LIST OF PUBLICATIONS

13	Early azathioprine/biological therapy is associated with decreased risk for first surgery and delays time to surgery but not reoperation in both smokers and nonsmokers with Crohn's disease, while smoking decreases the risk of colectomy in ulcerative colitis. <i>European Journal of Gastroenterology</i>	2.2	53
12	Mucosal expression of claudins 2, 3 and 4 in proximal and distal part of duodenum in children with coeliac disease. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 2010 , 456, 245-50	5.1	55
11	NKX2-3 and IRGM variants are associated with disease susceptibility to IBD in Eastern European patients. <i>World Journal of Gastroenterology</i> , 2010 , 16, 5233-40	5.6	15
10	Evaluation of the combined application of ethanol-fixed and formaldehyde-fixed neutrophil substrates for identifying atypical perinuclear antineutrophil cytoplasmic antibodies in inflammatory bowel disease. <i>Vaccine Journal</i> , 2009 , 16, 464-70		13
9	Pancreatic autoantibodies are associated with reactivity to microbial antibodies, penetrating disease behavior, perianal disease, and extraintestinal manifestations, but not with NOD2/CARD15 or TLR4 genotype in a Hungarian IBD cohort. <i>Inflammatory Bowel Diseases</i> , 2009 , 15, 365-74	4.5	51
8	The 3TUTR NFKBIA variant is associated with extensive colitis in Hungarian IBD patients. <i>Digestive Diseases and Sciences</i> , 2009 , 54, 351-9	4	13
7	Anti-microbial antibodies in celiac disease: trick or treat?. World Journal of Gastroenterology, 2009 , 15, 3891-900	5.6	20
6	Haptoglobin polymorphism: a novel genetic risk factor for celiac disease development and its clinical manifestations. <i>Clinical Chemistry</i> , 2008 , 54, 697-704	5.5	26
5	New serological markers for inflammatory bowel disease are associated with earlier age at onset, complicated disease behavior, risk for surgery, and NOD2/CARD15 genotype in a Hungarian IBD cohort. <i>American Journal of Gastroenterology</i> , 2008 , 103, 665-81	0.7	119
4	Seroreactivity to microbial components in Crohn's disease is associated with ileal involvement, noninflammatory disease behavior and NOD2/CARD15 genotype, but not with risk for surgery in a Hungarian cohort of IBD patients. <i>Inflammatory Bowel Diseases</i> , 2007 , 13, 984-92	4.5	59
3	Haptoglobin polymorphisms are associated with Crohn's disease, disease behavior, and extraintestinal manifestations in Hungarian patients. <i>Digestive Diseases and Sciences</i> , 2007 , 52, 1279-84	4	19
2	ATP-binding cassette transporter ABCG2 (BCRP) and ABCB1 (MDR1) variants are not associated with disease susceptibility, disease phenotype response to medical therapy or need for surgeryin Hungarian patients with inflammatory bowel diseases. <i>Scandinavian Journal of Gastroenterology</i> ,	2.4	34
1	DLG5 R30Q is not associated with IBD in Hungarian IBD patients but predicts clinical response to steroids in Crohn's disease. <i>Inflammatory Bowel Diseases</i> , 2006 , 12, 362-8	4.5	18