## Paulina Dumnicka

## List of Publications by Citations

Source: https://exaly.com/author-pdf/8937078/paulina-dumnicka-publications-by-citations.pdf

Version: 2024-04-17

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

635 58 15 22 h-index g-index citations papers 808 69 3.5 3.73 L-index avg, IF ext. papers ext. citations

#	Paper	IF	Citations
58	The Interplay between Inflammation, Coagulation and Endothelial Injury in the Early Phase of Acute Pancreatitis: Clinical Implications. <i>International Journal of Molecular Sciences</i> , <b>2017</b> , 18,	6.3	80
57	Malnutrition, Inflammation, Atherosclerosis Syndrome (MIA) and Diet Recommendations among End-Stage Renal Disease Patients Treated with Maintenance Hemodialysis. <i>Nutrients</i> , <b>2018</b> , 10,	6.7	46
56	Angiopoietin-2 Is an Early Indicator of Acute Pancreatic-Renal Syndrome in Patients with Acute Pancreatitis. <i>Mediators of Inflammation</i> , <b>2016</b> , 2016, 5780903	4.3	35
55	Markers of Glomerular and Tubular Damage in the Early Stage of Kidney Disease in Type 2 Diabetic Patients. <i>Mediators of Inflammation</i> , <b>2018</b> , 2018, 7659243	4.3	31
54	Interleukin-6 and neopterin levels in the serum and saliva of patients with Lichen planus and oral Lichen planus. <i>Journal of Oral Pathology and Medicine</i> , <b>2014</b> , 43, 734-9	3.3	26
53	Serum Concentrations of Angiopoietin-2 and Soluble fms-Like Tyrosine Kinase 1 (sFlt-1) Are Associated with Coagulopathy among Patients with Acute Pancreatitis. <i>International Journal of Molecular Sciences</i> , <b>2017</b> , 18,	6.3	25
52	Cardiovascular risk in chronic kidney disease patients: intima-media thickness predicts the incidence and severity of histologically assessed medial calcification in radial arteries. <i>BMC Nephrology</i> , <b>2015</b> , 16, 78	2.7	21
51	Sunitinib-induced hypothyroidism predicts progression-free survival in metastatic renal cell carcinoma patients. <i>Medical Oncology</i> , <b>2017</b> , 34, 68	3.7	20
50	Serum Soluble Fms-Like Tyrosine Kinase 1 (sFlt-1) Predicts the Severity of Acute Pancreatitis. <i>International Journal of Molecular Sciences</i> , <b>2016</b> , 17,	6.3	20
49	Osteoprotegerin and osteoprotegerin/TRAIL ratio are associated with cardiovascular dysfunction and mortality among patients with renal failure. <i>Advances in Medical Sciences</i> , <b>2016</b> , 61, 269-275	2.8	19
48	Potential Prognostic Markers of Acute Kidney Injury in the Early Phase of Acute Pancreatitis. <i>International Journal of Molecular Sciences</i> , <b>2019</b> , 20,	6.3	18
47	Relationship between aortic pulse wave velocity, selected proinflammatory cytokines, and vascular calcification parameters in peritoneal dialysis patients. <i>Journal of Hypertension</i> , <b>2014</b> , 32, 142-8	1.9	17
46	Vascular effects of advanced glycation end-products: content of immunohistochemically detected AGEs in radial artery samples as a predictor for arterial calcification and cardiovascular risk in asymptomatic patients with chronic kidney disease. <i>Disease Markers</i> , <b>2015</b> , 2015, 153978	3.2	16
45	Serum uromodulin concentrations correlate with glomerular filtration rate in patients with chronic kidney disease. <i>Polish Archives of Internal Medicine</i> , <b>2016</b> , 126, 995-1004	1.9	16
44	Co-occurring adverse events enable early prediction of progression-free survival in metastatic renal cell carcinoma patients treated with sunitinib: a hypothesis-generating study. <i>Tumori</i> , <b>2015</b> , 101, 555-9	1.7	15
43	Reference Values for TSH and Free Thyroid Hormones in Healthy Pregnant Women in Poland: A Prospective, Multicenter Study. <i>European Thyroid Journal</i> , <b>2017</b> , 6, 82-88	4.2	14
42	Serum Urokinase-Type Plasminogen Activator Receptor Does Not Outperform C-Reactive Protein and Procalcitonin as an Early Marker of Severity of Acute Pancreatitis. <i>Journal of Clinical Medicine</i> , <b>2018</b> , 7,	5.1	13

## (2016-2018)

41	Elevated Circulating Osteoprotegerin Levels in the Plasma of Hemodialyzed Patients With Severe Artery Calcification. <i>Therapeutic Apheresis and Dialysis</i> , <b>2018</b> , 22, 519-529	1.9	12	
40	Is Urinary NGAL Determination Useful for Monitoring Kidney Function and Assessment of Cardiovascular Disease? A 12-Month Observation of Patients with Type 2 Diabetes. <i>Disease Markers</i> , <b>2016</b> , 2016, 8489543	3.2	12	
39	Serum Uromodulin Levels in Prediction of Acute Kidney Injury in the Early Phase of Acute Pancreatitis. <i>Molecules</i> , <b>2017</b> , 22,	4.8	11	
38	Plasma pentraxin 3 concentrations in patients with acute pancreatitis. Clinical Laboratory, <b>2013</b> , 59, 100	0 <u>3</u> ⊵8	11	
37	Osteoprotegerin as a marker of cardiovascular risk in patients on peritoneal dialysis. <i>Polish Archives of Internal Medicine</i> , <b>2013</b> , 123, 149-55	1.9	10	
36	Pentraxin 3 as a new indicator of cardiovascular-related death in patients with advanced chronic kidney disease. <i>Polish Archives of Internal Medicine</i> , <b>2017</b> , 127, 170-177	1.9	10	
35	Urine NGAL is useful in the clinical evaluation of renal function in the early course of acute pancreatitis. <i>Folia Medica Cracoviensia</i> , <b>2016</b> , 56, 13-25	0.5	10	
34	Impaired fasting glucose and diabetes as predictors for radial artery calcification in end stage renal disease patients. <i>International Journal of Endocrinology</i> , <b>2013</b> , 2013, 969038	2.7	9	
33	Asymmetric dimethylarginine as a useful risk marker of radial artery calcification in patients with advanced kidney disease. <i>Polish Archives of Internal Medicine</i> , <b>2018</b> , 128, 157-165	1.9	9	
32	Transforming growth factor beta 1 as a risk factor for cardiovascular diseases in end-stage renal disease patients treated with peritoneal dialysis. <i>Clinical Laboratory</i> , <b>2014</b> , 60, 1163-8	2	9	
31	Endothelial injury is closely related to osteopontin and TNF receptor-mediated inflammation in end-stage renal disease. <i>Cytokine</i> , <b>2019</b> , 121, 154729	4	8	
30	Does the Automatic Measurement of Interleukin 6 Allow for Prediction of Complications during the First 48 h of Acute Pancreatitis?. <i>International Journal of Molecular Sciences</i> , <b>2018</b> , 19,	6.3	8	
29	Radiological Response and Neutrophil-to-Lymphocyte Ratio as Predictive Factors for Progression-Free and Overall Survival in Metastatic Renal Cell Carcinoma Patients Treated with Sunitinib. <i>Advances in Experimental Medicine and Biology</i> , <b>2019</b> , 1153, 31-45	3.6	7	
28	The Diagnostic Usefulness of Serum Total Bile Acid Concentrations in the Early Phase of Acute Pancreatitis of Varied Etiologies. <i>International Journal of Molecular Sciences</i> , <b>2017</b> , 18,	6.3	7	
27	Determination of serum neutrophil gelatinase-associated lipocalin at the early stage of acute pancreatitis. <i>Folia Medica Cracoviensia</i> , <b>2016</b> , 56, 5-16	0.5	7	
26	The correlation between the incidence of adverse events and progression-free survival in patients treated with cabozantinib for metastatic renal cell carcinoma (mRCC). <i>Medical Oncology</i> , <b>2019</b> , 36, 19	3.7	5	
25	Sterile leukocyturia affects urine neutrophil gelatinase-associated lipocalin concentration in type 2 diabetic patients. <i>Archives of Medical Science</i> , <b>2017</b> , 13, 321-327	2.9	5	
24	Macrocytosis during sunitinib treatment predicts progression-free survival in patients with metastatic renal cell carcinoma. <i>Medical Oncology</i> , <b>2016</b> , 33, 109	3.7	5	

23	Hand-Foot Syndrome and Progression-Free Survival in Patients Treated with Sunitinib for Metastatic Clear Cell Renal Cell Carcinoma. <i>Advances in Experimental Medicine and Biology</i> , <b>2019</b> , 1133, 35-40	3.6	4
22	Lower confidence limits for critical systematic errors. Clinical Biochemistry, 2007, 40, 1317-20	3.5	4
21	The Marker of Tubular Injury, Kidney Injury Molecule-1 (KIM-1), in Acute Kidney Injury Complicating Acute Pancreatitis: A Preliminary Study. <i>Journal of Clinical Medicine</i> , <b>2020</b> , 9,	5.1	3
20	Urinary Neutrophil Gelatinase-Associated Lipocalin Is Complementary to Albuminuria in Diagnosis of Early-Stage Diabetic Kidney Disease in Type 2 Diabetes. <i>BioMed Research International</i> , <b>2017</b> , 2017, 4691389	3	3
19	Proteoglycan/glycosaminoglycan and collagen content in the arterial wall of patients with end-stage renal disease: new indicators of vascular disease. <i>Polish Archives of Internal Medicine</i> , <b>2019</b> , 129, 781-789	1.9	3
18	Urinary NGAL Measured after the First Year Post Kidney Transplantation Predicts Changes in Glomerular Filtration over One-Year Follow-Up. <i>Journal of Clinical Medicine</i> , <b>2020</b> , 10,	5.1	3
17	Diagnostic Significance of Serum Galectin-3 in Hospitalized Patients with COVID-19-A Preliminary Study. <i>Biomolecules</i> , <b>2021</b> , 11,	5.9	3
16	Redox Balance Correlates with Nutritional Status among Patients with End-Stage Renal Disease Treated with Maintenance Hemodialysis. <i>Oxidative Medicine and Cellular Longevity</i> , <b>2019</b> , 2019, 6309465	5 <sup>6.7</sup>	2
15	Acute Phase Proteins and Vitamin D Seasonal Variation in End-Stage Renal Disease Patients. Journal of Clinical Medicine, <b>2020</b> , 9,	5.1	2
14	Does Beta-Trace Protein (BTP) Outperform Cystatin C as a Diagnostic Marker of Acute Kidney Injury Complicating the Early Phase of Acute Pancreatitis?. <i>Journal of Clinical Medicine</i> , <b>2020</b> , 9,	5.1	2
13	Carboxylated and intact osteocalcin predict adiponectin concentration in hemodialyzed patients. <i>Renal Failure</i> , <b>2016</b> , 38, 451-7	2.9	2
12	Effect of hemodialysis on acid leukocyte-type ribonuclease, alkaline ribonuclease and polymorphonuclear elastase serum levels in patients with end-stage renal disease. <i>Nephron Clinical Practice</i> , <b>2009</b> , 112, c248-54		2
11	Interplay of nitric oxide metabolites and markers of endothelial injury, inflammation, and vascular disease in the spectrum of advanced chronic kidney disease. <i>Kardiologia Polska</i> , <b>2020</b> , 78, 51-58	0.9	2
10	Complement Components in the Diagnosis and Treatment after Kidney Transplantation-Is There a Missing Link?. <i>Biomolecules</i> , <b>2021</b> , 11,	5.9	2
9	Osteoprotegerin, trail and osteoprotegerin/trail ratio in patients at early phase of acute pancreatitis. <i>Folia Medica Cracoviensia</i> , <b>2014</b> , 54, 17-26	0.5	2
8	Renal Impairment Detectors: IGFBP-7 and NGAL as Tubular Injury Markers in Multiple Myeloma Patients <i>Medicina (Lithuania)</i> , <b>2021</b> , 57,	3.1	2
7	A Multicenter Prospective Study on Efficacy and Safety of Ponatinib in Patients with Chronic Myeloid Leukemia Resistant or Intolerant to Previous Therapy: A 3-Year Report. <i>Blood</i> , <b>2020</b> , 136, 40-41	2.2	1
6	Evaluating the Relationship of GDF-15 with Clinical Characteristics, Cardinal Features, and Survival in Multiple Myeloma. <i>Mediators of Inflammation</i> , <b>2020</b> , 2020, 5657864	4.3	1

## LIST OF PUBLICATIONS

5	Sleep problems among children with Fetal Alcohol Spectrum Disorders (FASD)- an explorative study. <i>Italian Journal of Pediatrics</i> , <b>2021</b> , 47, 113	3.2	О
4	MP500BODY COMPOSITION ASSESSED BY BIOIMPEDANCE CORRELATES WITH LEFT VENTRICULAR MASS INDEX IN STAGE 5 CKD PATIENTS TREATED WITH PERITONEAL DIALYSIS. <i>Nephrology Dialysis Transplantation</i> , <b>2016</b> , 31, i507-i507	4.3	
3	FP514RELATIONSHIP BETWEEN SELECTED EXTRACELLULAR MATRIX COMPONENTS ASSESSED HISTOLOGICALLY IN RADIAL ARTERIES AND CARDIOVASCULAR COMORBIDITIES IN ESRD PATIENTS. <i>Nephrology Dialysis Transplantation</i> , <b>2018</b> , 33, i211-i212	4.3	
2	SP539GASTRIC INHIBITORY POLYPEPTIDE (GIP) AS A NOVEL INDICATOR OF EARLY MORPHOLOGICAL CHANGES IN RADIAL ARTERIES OF CKD STAGE 5 PATIENTS. <i>Nephrology Dialysis Transplantation</i> , <b>2018</b> , 33, i530-i530	4.3	
1	Determination of hepatocyte growth factor at early phase of acute pancreatitis. <i>Folia Medica Cracoviensia</i> , <b>2013</b> , 53, 87-95	0.5	