

Hany Onsy Habashy

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8945496/publications.pdf>

Version: 2024-02-01

15
papers

1,161
citations

759233

12
h-index

1058476

14
g-index

15
all docs

15
docs citations

15
times ranked

2170
citing authors

#	ARTICLE	IF	CITATIONS
1	RRM2 expression in different molecular subtypes of breast cancer and its prognostic significance. <i>Diagnostic Pathology</i> , 2022, 17, 1.	2.0	15
2	Electron microscopic and pathological changes of lung cancer after intratumoral injection of sodium bicarbonate. <i>Current Cancer Therapy Reviews</i> , 2021, 17, .	0.3	0
3	Expression levels of β -catenin and galectin-3 in meningioma and their effect on brain invasion and recurrence: a tissue microarray study. <i>Cancer Biology and Medicine</i> , 2017, 14, 319.	3.0	10
4	DACH1: Its Role as a Classifier of Long Term Good Prognosis in Luminal Breast Cancer. <i>PLoS ONE</i> , 2014, 9, e84428.	2.5	24
5	Immunoglobulin free light chains are biomarkers of poor prognosis in basal-like breast cancer and are potential targets in tumor-associated inflammation. <i>Oncotarget</i> , 2014, 5, 3159-3167.	1.8	34
6	The oestrogen receptor coactivator CARM1 has an oncogenic effect and is associated with poor prognosis in breast cancer. <i>Breast Cancer Research and Treatment</i> , 2013, 140, 307-316.	2.5	40
7	A review of the biological and clinical characteristics of luminal-like oestrogen receptor-positive breast cancer. <i>Histopathology</i> , 2012, 60, 854-863.	2.9	22
8	REG (Ras-like, oestrogen-regulated, growth-inhibitor) expression in breast cancer: a marker of ER-positive luminal-like subtype. <i>Breast Cancer Research and Treatment</i> , 2011, 128, 315-326.	2.5	41
9	FOXO3a nuclear localisation is associated with good prognosis in luminal-like breast cancer. <i>Breast Cancer Research and Treatment</i> , 2011, 129, 11-21.	2.5	69
10	Alpha- and beta-adrenergic receptor (AR) protein expression is associated with poor clinical outcome in breast cancer: an immunohistochemical study. <i>Breast Cancer Research and Treatment</i> , 2011, 130, 457-463.	2.5	87
11	Transferrin receptor (CD71) is a marker of poor prognosis in breast cancer and can predict response to tamoxifen. <i>Breast Cancer Research and Treatment</i> , 2010, 119, 283-293.	2.5	193
12	The prognostic significance of PELP1 expression in invasive breast cancer with emphasis on the ER-positive luminal-like subtype. <i>Breast Cancer Research and Treatment</i> , 2010, 120, 603-612.	2.5	83
13	Investigating AP-2 and YY1 protein expression as a cause of high HER2 gene transcription in breast cancers with discordant HER2 gene amplification. <i>Breast Cancer Research</i> , 2009, 11, R90.	5.0	42
14	Triple-Negative Breast Cancer: Distinguishing between Basal and Nonbasal Subtypes. <i>Clinical Cancer Research</i> , 2009, 15, 2302-2310.	7.0	422
15	Forkhead-box A1 (FOXA1) expression in breast cancer and its prognostic significance. <i>European Journal of Cancer</i> , 2008, 44, 1541-1551.	2.8	79