

Mikkel stergaard

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/8947778/mikkel-ostergaard-publications-by-year.pdf>

Version: 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

365
papers

12,532
citations

64
h-index

99
g-index

428
ext. papers

14,928
ext. citations

3.7
avg, IF

6.15
L-index

#	Paper	IF	Citations
365	Prediction of flare following remission and treatment withdrawal in early rheumatoid arthritis: post hoc analysis of a phase IIIb trial with abatacept.. <i>Arthritis Research and Therapy</i> , 2022 , 24, 47	5.7	2
364	The Programmed Death-1 Pathway Counter-Regulates Inflammation-Induced Osteoclast Activity in Clinical and Experimental Settings.. <i>Frontiers in Immunology</i> , 2022 , 13, 773946	8.4	2
363	Correspondence on 'No efficacy of anti-IL-23 therapy for axial spondyloarthritis in randomised controlled trials but in post-hoc analyses of psoriatic arthritis-related 'physician-reported spondylitis'? by Braun and Landewijck <i>Annals of the Rheumatic Diseases</i> , 2022 ,	2.4	1
362	Enthesitis in patients with psoriatic arthritis and axial spondyloarthritis - data from the Danish nationwide DANBIO registry.. <i>Seminars in Arthritis and Rheumatism</i> , 2021 , 52, 151948	5.3	0
361	Criterion validity of ultrasound in the identification of calcium pyrophosphate crystal deposits at the knee: an OMERACT ultrasound study. <i>Annals of the Rheumatic Diseases</i> , 2021 , 80, 261-267	2.4	9
360	Spinal Radiographic Progression and Predictors of Progression in Patients With Radiographic Axial Spondyloarthritis Receiving Ixekizumab Over 2 Years. <i>Journal of Rheumatology</i> , 2021 ,	4.1	3
359	Doppler ultrasound predicts successful discontinuation of biological DMARDs in rheumatoid arthritis patients in clinical remission. <i>Rheumatology</i> , 2021 , 60, 5549-5559	3.9	3
358	Response to: 'Correspondence on 'MRI lesions in the sacroiliac joints of patients with spondyloarthritis: an update of definitions and validation by the ASAS MRI working group" by Jibri. <i>Annals of the Rheumatic Diseases</i> , 2021 ,	2.4	
357	Development and Validation of 3 Preliminary MRI Sacroiliac Joint Composite Structural Damage Scores in a 5-year Longitudinal Axial Spondyloarthritis Study. <i>Journal of Rheumatology</i> , 2021 , 48, 1537-1546	4.1	0
356	POS0056 ANXIETY AND CONCERNS RELATED TO THE WORK SITUATION DURING THE COVID-19 PANDEMIC IN >5,000 PATIENTS WITH INFLAMMATORY RHEUMATIC DISEASE FOLLOWED IN THE DANISH DANBIO REGISTRY, RESULTS FROM A NATIONWIDE QUESTIONNAIRE. <i>Annals of the Rheumatic Diseases</i> , 2021 , 80, 224-234	2.4	
355	POS0918 EVALUATION OF SPINAL RADIOGRAPHIC PROGRESSION IN PATIENTS WITH RADIOGRAPHIC AXIAL SPONDYLOARTHRITIS RECEIVING IXEKIZUMAB THERAPY OVER 2 YEARS. <i>Annals of the Rheumatic Diseases</i> , 2021 , 80, 720.1-720	2.4	0
354	POS0454 COMPARISON OF MBDA SCORE, PATIENT GLOBAL ASSESSMENT AND EVALUATOR GLOBAL ASSESSMENT FOR PREDICTING RISK OF RADIOGRAPHIC PROGRESSION. <i>Annals of the Rheumatic Diseases</i> , 2021 , 80, 457-458	2.4	
353	POS1040 IMPLEMENTATION OF THE OMERACT PSAMRIS IN A PHASE IIB, RANDOMISED PLACEBO-CONTROLLED STUDY OF ABATACEPT IN PSORIATIC ARTHRITIS. <i>Annals of the Rheumatic Diseases</i> , 2021 , 80, 794-794	2.4	
352	AB0594 IMPROVING OSTEOARTHRITIS CARE BY AUTOMATIC MEASUREMENT OF HIP EFFUSION USING AI. <i>Annals of the Rheumatic Diseases</i> , 2021 , 80, 1334.1-1334	2.4	
351	POS0027 SECULAR TRENDS IN BASELINE CHARACTERISTICS, TREATMENT RETENTION AND RESPONSE RATES IN 27189 BIO-NATIVE AXIAL SPONDYLOARTHRITIS PATIENTS INITIATING TNFI INFLAMMATORY RHEUMATIC DISEASE. RESULTS FROM THE EUROSPA COLLABORATION. <i>Annals of the Rheumatic Diseases</i> , 2021 , 80, 217-218	2.4	0
350	Anxiety and concerns related to the work situation during the second wave of the COVID-19 pandemic in >5000 patients with inflammatory rheumatic disease followed in the DANBIO registry. <i>RMD Open</i> , 2021 , 7,	5.9	1
349	POS0443 EXPLORING NOVEL TENOSYNOVITIS AND COMBINED INFLAMMATION IMAGING OUTCOMES: RESULTS FROM A RANDOMISED CONTROLLED TRIAL IN EARLY RHEUMATOID ARTHRITIS. <i>Annals of the Rheumatic Diseases</i> , 2021 , 80, 450-451	2.4	

348	OP0159-HPR IMPACT OF NON-PHARMACOLOGICAL INTERVENTIONS TARGETING SLEEP DISTURBANCES OR DISORDERS IN PATIENTS WITH INFLAMMATORY ARTHRITIS: A SYSTEMATIC REVIEW AND META-ANALYSIS OF RANDOMISED TRIALS. <i>Annals of the Rheumatic Diseases</i> , 2021 , 80, 95-96	2.4	
347	Effectiveness and treatment retention of TNF inhibitors when used as monotherapy versus comedication with csDMARDs in 15 332 patients with psoriatic arthritis. Data from the EuroSpA collaboration. <i>Annals of the Rheumatic Diseases</i> , 2021 , 80, 1410-1418	2.4	6
346	Volumetric quantitative measurement of hip effusions by manual versus automated artificial intelligence techniques: An OMERACT preliminary validation study. <i>Seminars in Arthritis and Rheumatism</i> , 2021 , 51, 623-626	5.3	2
345	Consensus-based semi-quantitative ultrasound scoring system for gout lesions: Results of an OMERACT Delphi process and web-reliability exercise. <i>Seminars in Arthritis and Rheumatism</i> , 2021 , 51, 644-649	5.3	4
344	Rheumatoid Arthritis Relapse and Remission - Advancing Our Predictive Capability Using Modern Imaging. <i>Journal of Inflammation Research</i> , 2021 , 14, 2547-2555	4.8	2
343	Morphological characteristics of sacroiliac joint MRI lesions in axial spondyloarthritis and control subjects. <i>Rheumatology</i> , 2021 ,	3.9	3
342	Imaging in psoriatic arthritis: Status and recent advances. <i>Best Practice and Research in Clinical Rheumatology</i> , 2021 , 35, 101690	5.3	3
341	Impact of the COVID-19 pandemic on treat-to-target strategies and physical consultations in >7000 patients with inflammatory arthritis. <i>Rheumatology</i> , 2021 , 60, SI3-SI12	3.9	2
340	Whole-body Magnetic Resonance Imaging in Psoriatic Arthritis, Rheumatoid Arthritis, and Healthy Controls: Interscan, Intrareader, and Interreader Agreement and Distribution of Lesions. <i>Journal of Rheumatology</i> , 2021 , 48, 198-206	4.1	2
339	Which ultrasound lesions contribute to dactylitis in psoriatic arthritis and their reliability in a clinical setting. <i>Clinical Rheumatology</i> , 2021 , 40, 1061-1067	3.9	1
338	Ultrasound for the diagnosis of gout-the value of gout lesions as defined by the Outcome Measures in Rheumatology ultrasound group. <i>Rheumatology</i> , 2021 , 60, 239-249	3.9	10
337	Anatomic Distribution of Sacroiliac Joint Lesions on Magnetic Resonance Imaging in Patients With Axial Spondyloarthritis and Control Subjects: A Prospective Cross-Sectional Study, Including Postpartum Women, Patients With Disc Herniation, Cleaning Staff, Runners, and Healthy Individuals. <i>Arthritis Care and Research</i> , 2021 , 73, 749-754	4.7	7
336	Gradual reduction of tophaceous deposits during urate-lowering therapy. <i>Joint Bone Spine</i> , 2021 , 88, 105049	2.9	
335	Axial involvement in patients with early peripheral spondyloarthritis: a prospective MRI study of sacroiliac joints and spine. <i>Annals of the Rheumatic Diseases</i> , 2021 , 80, 103-108	2.4	7
334	Predictors of joint damage progression and stringent remission in patients with established rheumatoid arthritis in clinical remission. <i>Rheumatology</i> , 2021 , 60, 380-391	3.9	4
333	EULAR recommendations for the reporting of ultrasound studies in rheumatic and musculoskeletal diseases (RMDs). <i>Annals of the Rheumatic Diseases</i> , 2021 , 80, 840-847	2.4	10
332	Self-protection strategies and health behaviour in patients with inflammatory rheumatic diseases during the COVID-19 pandemic: results and predictors in more than 12 000 patients with inflammatory rheumatic diseases followed in the Danish DANBIO registry. <i>RMD Open</i> , 2021 , 7,	5.9	17
331	Risk of serious infections in arthritis patients treated with biological drugs: a matched cohort study and development of prediction model. <i>Rheumatology</i> , 2021 , 60, 3834-3844	3.9	2

330	Using a DAS28-CRP-steered treat-to-target strategy does not eliminate subclinical inflammation as assessed by ultrasonography in rheumatoid arthritis patients in longstanding clinical remission. <i>Arthritis Research and Therapy</i> , 2021 , 23, 48	5.7	6
329	Plasma interferon-alpha is associated with double-positivity for autoantibodies but is not a predictor of remission in early rheumatoid arthritis-a spin-off study of the NORD-STAR randomized clinical trial. <i>Arthritis Research and Therapy</i> , 2021 , 23, 189	5.7	0
328	Arthritis and enthesitis in the hip and pelvis region in spondyloarthritis - OMERACT validation of two whole-body MRI methods. <i>Seminars in Arthritis and Rheumatism</i> , 2021 , 51, 940-945	5.3	2
327	Impact of filgotinib on sacroiliac joint MRI structural lesions at 12 weeks in patients with active ankylosing spondylitis (TORTUGA trial). <i>Rheumatology</i> , 2021 ,	3.9	2
326	Reliability and agreement of proton density-weighted vs. gadolinium-enhanced T1-weighted MRI in hand osteoarthritis. An OMERACT MRI special interest group reliability exercise. <i>Seminars in Arthritis and Rheumatism</i> , 2021 , 51, 929-932	5.3	0
325	Joint and enthesal inflammation in the knee region in spondyloarthritis - reliability and responsiveness of two OMERACT whole-body MRI scores. <i>Seminars in Arthritis and Rheumatism</i> , 2021 , 51, 933-939	5.3	0
324	The Value of Magnetic Resonance Imaging for Assessing Disease Extent and Prediction of Relapse in Early Peripheral Spondyloarthritis. <i>Arthritis and Rheumatology</i> , 2021 , 73, 2044-2051	9.5	0
323	Diffusion-weighted MR imaging in chronic non-bacterial osteitis: Proof-of-concept of the apparent diffusion coefficient as an outcome measure. <i>Acta Radiologica Open</i> , 2021 , 10, 20584601211044478	1.2	
322	Improving domain definition and outcome instrument selection: Lessons learned for OMERACT from imaging. <i>Seminars in Arthritis and Rheumatism</i> , 2021 , 51, 1125-1133	5.3	3
321	Validation of the adjusted multi-biomarker disease activity score as a prognostic test for radiographic progression in rheumatoid arthritis: a combined analysis of multiple studies. <i>Arthritis Research and Therapy</i> , 2021 , 23, 1	5.7	11
320	Similar lipid level changes in early rheumatoid arthritis patients following 1-year treat-to-target strategy with adalimumab plus methotrexate versus placebo plus methotrexate: secondary analyses from the randomised controlled OPERA trial. <i>Rheumatology International</i> , 2021 , 41, 543-549	3.6	2
319	Assessing the sensitivity to change of the OMERACT ultrasound structural gout lesions during urate-lowering therapy. <i>RMD Open</i> , 2020 , 6,	5.9	10
318	Cognitive behavioural therapy for insomnia in patients with rheumatoid arthritis: protocol for the randomised, single-blinded, parallel-group Sleep-RA trial. <i>Trials</i> , 2020 , 21, 440	2.8	3
317	MRI of the sacroiliac joints: what is and what is not sacroiliitis?. <i>Current Opinion in Rheumatology</i> , 2020 , 32, 357-364	5.3	4
316	Test-retest repeatability of the apparent diffusion coefficient in sacroiliac joint MRI in patients with axial spondyloarthritis and healthy individuals. <i>Acta Radiologica Open</i> , 2020 , 9, 2058460120906015	1.2	3
315	Future use of musculoskeletal ultrasonography and magnetic resonance imaging in rheumatoid arthritis. <i>Current Opinion in Rheumatology</i> , 2020 , 32, 264-272	5.3	5
314	Outcomes and Findings of the International Rheumatoid Arthritis (RA) BIODAM Cohort for Validation of Soluble Biomarkers in RA. <i>Journal of Rheumatology</i> , 2020 , 47, 796-808	4.1	1
313	Whole-Body Magnetic Resonance Imaging Assessment of Joint Inflammation in Rheumatoid Arthritis-Agreement With Ultrasonography and Clinical Evaluation. <i>Frontiers in Medicine</i> , 2020 , 7, 285	4.9	

312	Magnetic Resonance Imaging of Enthesitis in Spondyloarthritis, Including Psoriatic Arthritis-Status and Recent Advances. <i>Frontiers in Medicine</i> , 2020 , 7, 296	4.9	9
311	Serial magnetic resonance imaging and ultrasound examinations demonstrate differential inflammatory lesion patterns in soft tissue and bone upon patient-reported flares in rheumatoid arthritis. <i>Arthritis Research and Therapy</i> , 2020 , 22, 19	5.7	6
310	How do people with rheumatoid arthritis experience participation in a smoking cessation trial: a qualitative study. <i>International Journal of Qualitative Studies on Health and Well-being</i> , 2020 , 15, 1725997		2
309	Comparison of the Effects of Secukinumab and Adalimumab Biosimilar on Radiographic Progression in Patients with Ankylosing Spondylitis: Design of a Randomized, Phase IIIb Study (SURPASS). <i>Clinical Drug Investigation</i> , 2020 , 40, 269-278	3.2	16
308	Impact of discordance between patient's and evaluator's global assessment on treatment outcomes in 14'868 patients with spondyloarthritis. <i>Rheumatology</i> , 2020 , 59, 2455-2461	3.9	4
307	Association between MRI findings and patient-reported outcomes in patients with rheumatoid arthritis in clinical remission and at relapse. <i>International Journal of Rheumatic Diseases</i> , 2020 , 23, 488-498 ^{2,3}		3
306	Impact of season on the association between vitamin D levels at diagnosis and one-year remission in early Rheumatoid Arthritis. <i>Scientific Reports</i> , 2020 , 10, 7371	4.9	1
305	The utility of magnetic resonance imaging lesion combinations in the sacroiliac joints for diagnosing patients with axial spondyloarthritis. A prospective study of 204 participants including post-partum women, patients with disc herniation, cleaning staff, runners and healthy persons.	3.9	8
304	Novel whole-body magnetic resonance imaging response and remission criteria document diminished inflammation during golimumab treatment in axial spondyloarthritis. <i>Rheumatology</i> , 2020 , 59, 3358-3368	3.9	4
303	OP0177 PROPERTIES AND LOCATIONS OF COLOR-CODED DUAL ENERGY CT LESIONS IN GOUT PATIENTS [A SYSTEMATIC EVALUATION]. <i>Annals of the Rheumatic Diseases</i> , 2020 , 79, 110.1-110	2.4	
302	SAT0548 DEVELOPMENT AND VALIDATION OF THREE PRELIMINARY MRI SACROILIAC JOINT COMPOSITE STRUCTURAL DAMAGE SCORES IN A 5-YEAR LONGITUDINAL STUDY OF PATIENTS WITH AXIAL SPONDYLOARTHRITIS. <i>Annals of the Rheumatic Diseases</i> , 2020 , 79, 1231.1-1231	2.4	
301	GRAPPA 2019 Project Report. <i>Journal of Rheumatology</i> , 2020 , 96, 53-57	4.1	6
300	AB1107 DACTYLITIS IN PSORIATIC ARTHRITIS [PREVALENCE AND RELIABILITY OF ULTRASOUND PATHOLOGIES]. <i>Annals of the Rheumatic Diseases</i> , 2020 , 79, 1843-1844	2.4	
299	AB0194 VITAMIN D TRAJECTORIES IN EARLY DIAGNOSED, AGGRESSIVELY TREATED RHEUMATOID ARTHRITIS PATIENTS: A 10 YEAR LONGITUDINAL COHORT STUDY BASED ON THE DANISH CIMESTRA TRIAL. <i>Annals of the Rheumatic Diseases</i> , 2020 , 79, 1397-1397	2.4	
298	THU0541 ANATOMICAL LOCATION OF SACROILIAC JOINT MRI LESIONS IN PATIENTS WITH AXIAL SPONDYLOARTHRITIS, POSTPARTUM WOMEN, PATIENTS WITH DISC HERNIATION, CLEANING STAFF, RUNNERS AND HEALTHY PERSONS. <i>Annals of the Rheumatic Diseases</i> , 2020 , 79, 510-511	2.4	
297	AB1243 TRAINING AND VALIDATION OF A MULTIVARIATE PREDICTOR OF RISK OF RADIOGRAPHIC PROGRESSION FOR PATIENTS WITH RHEUMATOID ARTHRITIS. <i>Annals of the Rheumatic Diseases</i> , 2020 , 79, 1913.1-1913	2.4	
296	AB1106 DO TENDER JOINTS IN PSORIATIC ARTHRITIS REFLECT INFLAMMATION ON ULTRASOUND?. <i>Annals of the Rheumatic Diseases</i> , 2020 , 79, 1842.1-1843	2.4	
295	Validation of assessment methods for the apparent diffusion coefficient in a clinical trial of axial spondyloarthritis patients treated with golimumab. <i>European Journal of Radiology Open</i> , 2020 , 7, 100285 ^{2.6}		1

294	Retention and response rates in 14 261 PsA patients starting TNF inhibitor treatment-results from 12 countries in EuroSpA. <i>Rheumatology</i> , 2020 , 59, 1640-1650	3.9	6
293	Scoring magnetic resonance imaging (MRI) inflammation and structural lesions in sacroiliac joints of patients with axial spondyloarthritis: assessment of all MRI slices of the cartilaginous compartment versus standardized six or five slices. <i>Scandinavian Journal of Rheumatology</i> , 2020 , 49, 200-209	1.9	1
292	Programmed death ligand 2 - A link between inflammation and bone loss in rheumatoid arthritis. <i>Journal of Translational Autoimmunity</i> , 2020 , 3, 100028	4.1	5
291	Ixekizumab for patients with non-radiographic axial spondyloarthritis (COAST-X): a randomised, placebo-controlled trial. <i>Lancet, The</i> , 2020 , 395, 53-64	4.0	74
290	Pain and Self-reported Swollen Joints Are Main Drivers of Patient-reported Flares in Rheumatoid Arthritis: Results from a 12-month Observational Study. <i>Journal of Rheumatology</i> , 2020 , 47, 1305-1313	4.1	5
289	Overall infection risk in rheumatoid arthritis during treatment with abatacept, rituximab and tocilizumab; an observational cohort study. <i>Rheumatology</i> , 2020 , 59, 1949-1956	3.9	10
288	Synovial hypertrophy without Doppler in the feet changes during treatment: results from a longitudinal study of rheumatoid arthritis patients initiating biological treatment. <i>Rheumatology</i> , 2020 , 59, 1765-1767	3.9	1
287	Response to: 'Correspondence on 'Ultrasound shows rapid reduction of crystal depositions during a treat-to-target approach in gout patients: 12-month results from the NOR-Gout study' by Hung. <i>Annals of the Rheumatic Diseases</i> , 2020 ,	2.4	
286	Imaging in the diagnosis and management of peripheral psoriatic arthritis. <i>Best Practice and Research in Clinical Rheumatology</i> , 2020 , 34, 101594	5.3	3
285	Ultrasound shows rapid reduction of crystal depositions during a treat-to-target approach in gout patients: 12-month results from the NOR-Gout study. <i>Annals of the Rheumatic Diseases</i> , 2020 , 79, 1500-1505	2.4	13
284	Secukinumab Provides Sustained Reduction in Fatigue in Patients with Ankylosing Spondylitis: Long-term Results of Two Phase III Randomized Controlled Trials. <i>Arthritis Care and Research</i> , 2020 ,	4.7	3
283	Active conventional treatment and three different biological treatments in early rheumatoid arthritis: phase IV investigator initiated, randomised, observer blinded clinical trial. <i>BMJ, The</i> , 2020 , 371, m4328	5.9	10
282	Peripheral Enthesitis Detected by Ultrasonography in Patients With Axial Spondyloarthritis-Anatomical Distribution, Morphology, and Response to Tumor Necrosis Factor-Inhibitor Therapy. <i>Frontiers in Medicine</i> , 2020 , 7, 341	4.9	2
281	Atlas of the OMERACT Heel Enthesitis MRI Scoring System (HEMRIS). <i>RMD Open</i> , 2020 , 6,	5.9	6
280	Drug retention, inactive disease and response rates in 1860 patients with axial spondyloarthritis initiating secukinumab treatment: routine care data from 13 registries in the EuroSpA collaboration. <i>RMD Open</i> , 2020 , 6,	5.9	10
279	Dual-energy CT in gout patients: Do all colour-coded lesions actually represent monosodium urate crystals?. <i>Arthritis Research and Therapy</i> , 2020 , 22, 212	5.7	14
278	Flares in rheumatoid arthritis: do patient-reported swollen and tender joints match clinical and ultrasonography findings?. <i>Rheumatology</i> , 2020 , 59, 129-136	3.9	9
277	Whole-body Magnetic Resonance Imaging Inflammation in Peripheral Joints and Enteses in Axial Spondyloarthritis: Distribution and Changes during Adalimumab Treatment. <i>Journal of Rheumatology</i> , 2020 , 47, 50-58	4.1	10

276	High versus standard magnetic resonance image resolution of the cervical spine in patients with axial spondyloarthritis. <i>Acta Radiologica</i> , 2020 , 61, 471-479	2	
275	Adherence to Treat-to-target Management in Rheumatoid Arthritis and Associated Factors: Data from the International RA BIODAM Cohort. <i>Journal of Rheumatology</i> , 2020 , 47, 809-819	4.1	7
274	Responsiveness of different dynamic contrast-enhanced magnetic resonance imaging approaches: a post-hoc analysis of a randomized controlled trial of certolizumab pegol in rheumatoid arthritis. <i>Scandinavian Journal of Rheumatology</i> , 2020 , 49, 105-111	1.9	1
273	Evolving the comprehensive management of rheumatoid arthritis: identification of unmet needs and development of practical and educational tools. <i>Clinical and Experimental Rheumatology</i> , 2020 , 38, 1056-1067	2.2	0
272	MRI lesions in the sacroiliac joints of patients with spondyloarthritis: an update of definitions and validation by the ASAS MRI working group. <i>Annals of the Rheumatic Diseases</i> , 2019 , 78, 1550-1558	2.4	81
271	Treatment response and drug retention rates in 24 195 biologic-naïve patients with axial spondyloarthritis initiating TNFi treatment: routine care data from 12 registries in the EuroSpA collaboration. <i>Annals of the Rheumatic Diseases</i> , 2019 , 78, 1536-1544	2.4	23
270	The need for comparative data in spondyloarthritis. <i>Arthritis Research and Therapy</i> , 2019 , 21, 32	5.7	3
269	Utility of Magnetic Resonance Imaging in Diagnosis and Monitoring Enthesitis in Patients with Spondyloarthritis: An OMERACT Systematic Literature Review. <i>Journal of Rheumatology</i> , 2019 , 46, 1207-1214	4.1	7
268	The OMERACT MRI in Enthesitis Initiative: Definitions of Key Pathologies, Suggested MRI Sequences, and a Novel Heel Enthesitis Scoring System. <i>Journal of Rheumatology</i> , 2019 , 46, 1232-1238	4.1	25
267	OMERACT Definitions for Ultrasonographic Pathologies and Elementary Lesions of Rheumatic Disorders 15 Years On. <i>Journal of Rheumatology</i> , 2019 , 46, 1388-1393	4.1	57
266	Effect of Magnetic Resonance Imaging vs Conventional Treat-to-Target Strategies on Disease Activity Remission and Radiographic Progression in Rheumatoid Arthritis: The IMAGINE-RA Randomized Clinical Trial. <i>JAMA - Journal of the American Medical Association</i> , 2019 , 321, 461-472	27.4	41
265	Musculoskeletal Ultrasound in Systemic Lupus Erythematosus: Systematic Literature Review by the Lupus Task Force of the OMERACT Ultrasound Working Group. <i>Journal of Rheumatology</i> , 2019 , 46, 1379-1387	4.1	10
264	Morbidity and mortality in patients with rheumatoid arthritis compared with an age- and sex-matched control population: A nationwide register study. <i>Journal of Comorbidity</i> , 2019 , 9, 2235042X19853484	4.1	10
263	Efficacy and safety of cannabidiol followed by an open label add-on of tetrahydrocannabinol for the treatment of chronic pain in patients with rheumatoid arthritis or ankylosing spondylitis: protocol for a multicentre, randomised, placebo-controlled study. <i>BMJ Open</i> , 2019 , 9, e028197	3	8
262	The OMERACT Stepwise Approach to Select and Develop Imaging Outcome Measurement Instruments: The Musculoskeletal Ultrasound Example. <i>Journal of Rheumatology</i> , 2019 , 46, 1394-1400	4.1	15
261	Imaging in rheumatoid arthritis: the role of magnetic resonance imaging and computed tomography. <i>Radiologia Medica</i> , 2019 , 124, 1128-1141	6.5	25
260	Quality of life and contact with healthcare systems among patients with psoriasis and psoriatic arthritis: results from the NORdic PATient survey of Psoriasis and Psoriatic arthritis (NORPAPP). <i>Archives of Dermatological Research</i> , 2019 , 311, 351-360	3.3	13
259	Validity and Responsiveness of Combined Inflammation and Combined Joint Damage Scores Based on the OMERACT Rheumatoid Arthritis MRI Scoring System (RAMRIS). <i>Journal of Rheumatology</i> , 2019 , 46, 1222-1227	4.1	5

258	Extremely poor patient-reported outcomes are associated with lack of clinical response and decreased retention rate of tumour necrosis factor inhibitor treatment in patients with axial spondyloarthritis. <i>Scandinavian Journal of Rheumatology</i> , 2019 , 48, 128-132	1.9	5
257	Anti-cyclic citrullinated peptide antibodies, 28-joint Disease Activity Score, and magnetic resonance imaging bone oedema at baseline predict 11 years' functional and radiographic outcome in early rheumatoid arthritis. <i>Scandinavian Journal of Rheumatology</i> , 2019 , 48, 1-8	1.9	15
256	Predictive value of a multi-biomarker disease activity score for clinical remission and radiographic progression in patients with early rheumatoid arthritis: a post-hoc study of the OPERA trial. <i>Scandinavian Journal of Rheumatology</i> , 2019 , 48, 9-16	1.9	14
255	Magnetic Resonance Imaging of Lesions in the Sacroiliac Joints for Differentiation of Patients With Axial Spondyloarthritis From Control Subjects With or Without Pelvic or Buttock Pain: A Prospective, Cross-Sectional Study of 204 Participants. <i>Arthritis and Rheumatology</i> , 2019 , 71, 2034-2046	9.5	27
254	What Level of Inflammation Leads to Structural Damage in the Sacroiliac Joints? A Four-Year Magnetic Resonance Imaging Follow-Up Study of Low Back Pain Patients. <i>Arthritis and Rheumatology</i> , 2019 , 71, 2027-2033	9.5	8
253	Imaging of Common Rheumatic Joint Diseases Affecting the Upper Limbs. <i>Radiologic Clinics of North America</i> , 2019 , 57, 1001-1034	2.3	2
252	Determining MRI Inflammation Targets When Considering a Rheumatoid Arthritis Treat-to-Target Strategy: Results of a Randomized, Placebo-Controlled Trial. <i>Advances in Therapy</i> , 2019 , 36, 2384-2393	4.1	3
251	Very early MRI responses to therapy as a predictor of later radiographic progression in early rheumatoid arthritis. <i>Arthritis Research and Therapy</i> , 2019 , 21, 214	5.7	9
250	Development and Validation of an OMERACT MRI Whole-Body Score for Inflammation in Peripheral Joints and Entheses in Inflammatory Arthritis (MRI-WIPE). <i>Journal of Rheumatology</i> , 2019 , 46, 1215-1221	4.1	24
249	Assessment of structural damage progression in established rheumatoid arthritis by conventional radiography, computed tomography, and magnetic resonance imaging. <i>Best Practice and Research in Clinical Rheumatology</i> , 2019 , 33, 101481	5.3	4
248	Canada-Denmark MRI scoring system of the spine in patients with axial spondyloarthritis: updated definitions, scoring rules and inter-reader reliability in a multiple reader setting. <i>RMD Open</i> , 2019 , 5, e001057	5.0	9
247	Adjustment of the multi-biomarker disease activity score to account for age, sex and adiposity in patients with rheumatoid arthritis. <i>Rheumatology</i> , 2019 , 58, 874-883	3.9	12
246	Dose tapering and discontinuation of biological therapy in rheumatoid arthritis patients in routine care - 2-year outcomes and predictors. <i>Rheumatology</i> , 2019 , 58, 110-119	3.9	19
245	Longitudinal Reliability of the OMERACT Thumb Base Osteoarthritis Magnetic Resonance Imaging Scoring System (TOMS). <i>Journal of Rheumatology</i> , 2019 , 46, 1228-1231	4.1	2
244	Structural progression rate decreases over time on serial radiography and magnetic resonance imaging of sacroiliac joints and spine in a five-year follow-up study of patients with ankylosing spondylitis treated with tumour necrosis factor inhibitor. <i>Scandinavian Journal of Rheumatology</i> , 2019 , 48, 185-197	1.9	13
243	Risk of serious infections in patients with rheumatoid arthritis treated in routine care with abatacept, rituximab and tocilizumab in Denmark and Sweden. <i>Annals of the Rheumatic Diseases</i> , 2019 , 78, 320-327	2.4	31
242	Treatment use and satisfaction among patients with psoriasis and psoriatic arthritis: results from the NORdic Patient survey of Psoriasis and Psoriatic arthritis (NORPAPP). <i>Journal of the European Academy of Dermatology and Venereology</i> , 2019 , 33, 340-354	4.6	13
241	Inflammatory and structural changes in vertebral bodies and posterior elements of the spine in axial spondyloarthritis: construct validity, responsiveness and discriminatory ability of the anatomy-based CANDEN scoring system in a randomised placebo-controlled trial. <i>RMD Open</i> , 2018 , 4, e000624	5.9	20

240	Atlas for the OMERACT thumb base osteoarthritis MRI scoring system (TOMS). <i>RMD Open</i> , 2018 , 4, e000583	4	4
239	Whole-body Magnetic Resonance Imaging in Axial Spondyloarthritis: Reduction of Sacroiliac, Spinal, and Enthesal Inflammation in a Placebo-controlled Trial of Adalimumab. <i>Journal of Rheumatology</i> , 2018 , 45, 621-629	4.1	22
238	Circulating serum interleukin-6, serum chitinase-3-like protein-1, and plasma vascular endothelial growth factor are not predictive for remission and radiographic progression in patients with early rheumatoid arthritis: post-hoc explorative and validation studies based on the CIMESTRA and OPERA trials. <i>Scandinavian Journal of Rheumatology</i> , 2018 , 47, 259-269	1.9	2
237	The Use of the OMERACT Ultrasound Tenosynovitis Scoring System in Multicenter Clinical Trials. <i>Journal of Rheumatology</i> , 2018 , 45, 165-169	4.1	4
236	Identification of calcium pyrophosphate deposition disease (CPPD) by ultrasound: reliability of the OMERACT definitions in an extended set of joints-an international multiobserver study by the OMERACT Calcium Pyrophosphate Deposition Disease Ultrasound Subtask Force. <i>Annals of the Rheumatic Diseases</i> , 2018 , 77, 1194-1199	2.4	24
235	Development and Validation of MRI Sacroiliac Joint Scoring Methods for the Semiaxial Scan Plane Corresponding to the Berlin and SPARCC MRI Scoring Methods, and of a New Global MRI Sacroiliac Joint Method. <i>Journal of Rheumatology</i> , 2018 , 45, 70-77	4.1	13
234	Structural damage progression in patients with early rheumatoid arthritis treated with methotrexate, baricitinib, or baricitinib plus methotrexate based on clinical response in the phase 3 RA-BEGIN study. <i>Clinical Rheumatology</i> , 2018 , 37, 2381-2390	3.9	18
233	The D-vitamin metabolite 1,25(OH) D in serum is associated with disease activity and Anti-Citrullinated Protein Antibodies in active and treatment naïve, early Rheumatoid Arthritis Patients. <i>Scandinavian Journal of Immunology</i> , 2018 , 88, e12704	3.4	8
232	Risankizumab, an IL-23 inhibitor, for ankylosing spondylitis: results of a randomised, double-blind, placebo-controlled, proof-of-concept, dose-finding phase 2 study. <i>Annals of the Rheumatic Diseases</i> , 2018 , 77, 1295-1302	2.4	188
231	Options for Assessing Joints and Entheses in Psoriatic Arthritis by Ultrasonography and Magnetic Resonance Imaging: How to Move Forward. <i>Journal of Rheumatology</i> , 2018 , 94, 44-47	4.1	6
230	Plasma MicroRNA Profiles in Patients with Early Rheumatoid Arthritis Responding to Adalimumab plus Methotrexate vs Methotrexate Alone: A Placebo-controlled Clinical Trial. <i>Journal of Rheumatology</i> , 2018 , 45, 53-61	4.1	23
229	Minimal Disease Activity as a Treatment Target in Psoriatic Arthritis: A Review of the Literature. <i>Journal of Rheumatology</i> , 2018 , 45, 6-13	4.1	38
228	Is synovial hypertrophy without Doppler activity sensitive to change? Post-hoc analysis from a rheumatoid arthritis ultrasound study. <i>Arthritis Research and Therapy</i> , 2018 , 20, 224	5.7	10
227	Magnetic Resonance Imaging in Rheumatology. <i>Magnetic Resonance Imaging Clinics of North America</i> , 2018 , 26, 599-613	1.6	5
226	Update on imaging in rheumatic diseases. <i>Clinical and Experimental Rheumatology</i> , 2018 , 36 Suppl 114, 2	2.2	0
225	Monitoring total-body inflammation and damage in joints and entheses: the first follow-up study of whole-body magnetic resonance imaging in rheumatoid arthritis. <i>Scandinavian Journal of Rheumatology</i> , 2017 , 46, 253-262	1.9	14
224	Short-term changes on MRI predict long-term changes on radiography in rheumatoid arthritis: an analysis by an OMERACT Task Force of pooled data from four randomised controlled trials. <i>Annals of the Rheumatic Diseases</i> , 2017 , 76, 992-997	2.4	16
223	Is MRI a predictive biomarker for clinical response to biologics in rheumatoid arthritis?. <i>Annals of the Rheumatic Diseases</i> , 2017 , 76, e45	2.4	2

222	Definition and Reliability Assessment of Elementary Ultrasonographic Findings in Calcium Pyrophosphate Deposition Disease: A Study by the OMERACT Calcium Pyrophosphate Deposition Disease Ultrasound Subtask Force. <i>Journal of Rheumatology</i> , 2017 , 44, 1744-1749	4.1	38
221	Intramuscular versus ultrasound-guided intratenosynovial glucocorticoid injection for tenosynovitis in patients with rheumatoid arthritis: a randomised, double-blind, controlled study. <i>Annals of the Rheumatic Diseases</i> , 2017 , 76, 666-672	2.4	17
220	Association between baseline vitamin D metabolite levels and long-term cardiovascular events in patients with rheumatoid arthritis from the CIMESTRA trial: protocol for a cohort study with patient-record evaluated outcomes. <i>BMJ Open</i> , 2017 , 7, e014816	3	2
219	Development and Validation of the OMERACT Rheumatoid Arthritis Magnetic Resonance Tenosynovitis Scoring System in a Multireader Exercise. <i>Journal of Rheumatology</i> , 2017 , 44, 1688-1693	4.1	20
218	Head-to-head comparison of aggressive conventional therapy and three biological treatments and comparison of two de-escalation strategies in patients who respond to treatment: study protocol for a multicenter, randomized, open-label, blinded-assessor, phase 4 study. <i>Trials</i> , 2017 , 18, 161	2.8	11
217	Ankylosing Spondylitis versus Nonradiographic Axial Spondyloarthritis: Comparison of Tumor Necrosis Factor Inhibitor Effectiveness and Effect of HLA-B27 Status. An Observational Cohort Study from the Nationwide DANBIO Registry. <i>Journal of Rheumatology</i> , 2017 , 44, 59-69	4.1	30
216	What the Rheumatologist Is Looking for and What the Radiologist Should Know in Imaging for Rheumatoid Arthritis. <i>Radiologic Clinics of North America</i> , 2017 , 55, 905-916	2.3	3
215	Whole-body Magnetic Resonance Imaging in Inflammatory Arthritis: Systematic Literature Review and First Steps Toward Standardization and an OMERACT Scoring System. <i>Journal of Rheumatology</i> , 2017 , 44, 1699-1705	4.1	33
214	The efficacy of motivational counselling and SMS reminders on daily sitting time in patients with rheumatoid arthritis: a randomised controlled trial. <i>Annals of the Rheumatic Diseases</i> , 2017 , 76, 1603-1606	2.4	48
213	Soluble CD206 plasma levels in rheumatoid arthritis reflect decrease in disease activity. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2017 , 77, 385-389	2	6
212	Magnetic resonance imaging assessed inflammation in the wrist is associated with patient-reported physical impairment, global assessment of disease activity and pain in early rheumatoid arthritis: longitudinal results from two randomised controlled trials. <i>Annals of the Rheumatic Diseases</i> , 2017 , 76, 1707-1715	2.4	21
211	Obesity and rates of clinical remission and low MRI inflammation in rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2017 , 76, 1743-1746	2.4	24
210	Work ability in rheumatoid arthritis patients: a register study on the prospective risk of exclusion and probability of returning to work. <i>Rheumatology</i> , 2017 , 56, 1135-1143	3.9	8
209	Expression of soluble CD83 in plasma from early-stage rheumatoid arthritis patients is not modified by anti-TNF- α therapy. <i>Cytokine</i> , 2017 , 96, 1-7	4	16
208	Development and Reliability of the OMERACT Thumb Base Osteoarthritis Magnetic Resonance Imaging Scoring System. <i>Journal of Rheumatology</i> , 2017 , 44, 1694-1698	4.1	22
207	Hand bone loss in early rheumatoid arthritis during a methotrexate-based treat-to-target strategy with or without adalimumab-a substudy of the optimized treatment algorithm in early RA (OPERA) trial. <i>Clinical Rheumatology</i> , 2017 , 36, 781-789	3.9	7
206	Scoring ultrasound synovitis in rheumatoid arthritis: a EULAR-OMERACT ultrasound taskforce-Part 2: reliability and application to multiple joints of a standardised consensus-based scoring system. <i>RMD Open</i> , 2017 , 3, e000427	5.9	89
205	Galectin-3 is Persistently Increased in Early Rheumatoid Arthritis (RA) and Associates with Anti-CCP Seropositivity and MRI Bone Lesions, While Early Fibrosis Markers Correlate with Disease Activity. <i>Scandinavian Journal of Immunology</i> , 2017 , 86, 471-478	3.4	14

204	Ultrasound of the hand is sufficient to detect subclinical inflammation in rheumatoid arthritis remission: a post hoc longitudinal study. <i>Arthritis Research and Therapy</i> , 2017 , 19, 221	5.7	22
203	Scoring ultrasound synovitis in rheumatoid arthritis: a EULAR-OMERACT ultrasound taskforce Part 1: definition and development of a standardised, consensus-based scoring system. <i>RMD Open</i> , 2017 , 3, e000428	5.9	139
202	The 2017 EULAR standardised procedures for ultrasound imaging in rheumatology. <i>Annals of the Rheumatic Diseases</i> , 2017 , 76, 1974-1979	2.4	112
201	The OMERACT Rheumatoid Arthritis Magnetic Resonance Imaging (MRI) Scoring System: Updated Recommendations by the OMERACT MRI in Arthritis Working Group. <i>Journal of Rheumatology</i> , 2017 , 44, 1706-1712	4.1	66
200	Welfare costs in patients with rheumatoid arthritis and their partners compared with matched controls: a register-based study. <i>Clinical Rheumatology</i> , 2017 , 36, 517-525	3.9	6
199	Increased galectin-3 may serve as a serologic signature of pre-rheumatoid arthritis while markers of synovitis and cartilage do not differ between early undifferentiated arthritis subsets. <i>Arthritis Research and Therapy</i> , 2017 , 19, 80	5.7	10
198	Limited Reliability of Radiographic Assessment of Sacroiliac Joints in Patients with Suspected Early Spondyloarthritis. <i>Journal of Rheumatology</i> , 2017 , 44, 70-77	4.1	32
197	Effect of a treat-to-target strategy based on methotrexate and intra-articular betamethasone with or without additional cyclosporin on MRI-assessed synovitis, osteitis, tenosynovitis, bone erosion, and joint space narrowing in early rheumatoid arthritis: results from a 2-year randomized controlled trial. <i>Journal of Rheumatology</i> , 2017 , 44, 335-345	1.9	7
196	No diagnostic utility of antibody patterns against <i>Klebsiella pneumoniae</i> capsular serotypes in patients with axial spondyloarthritis vs. patients with non-specific low back pain: a cross-sectional study. <i>Scandinavian Journal of Rheumatology</i> , 2017 , 46, 296-302	1.9	6
195	Defining the optimal biological monotherapy in rheumatoid arthritis: A systematic review and meta-analysis of randomised trials. <i>Seminars in Arthritis and Rheumatism</i> , 2017 , 46, 699-708	5.3	16
194	Investigation of a multi-biomarker disease activity score in rheumatoid arthritis by comparison with magnetic resonance imaging, computed tomography, ultrasonography, and radiography parameters of inflammation and damage. <i>Scandinavian Journal of Rheumatology</i> , 2017 , 46, 353-358	1.9	13
193	The effect of an intensive smoking cessation intervention on disease activity in patients with rheumatoid arthritis: study protocol for a randomised controlled trial. <i>Trials</i> , 2017 , 18, 570	2.8	12
192	Development and validation of rheumatoid arthritis magnetic resonance imaging inflammation thresholds associated with lack of damage progression. <i>Clinical and Experimental Rheumatology</i> , 2017 , 35, 607-613	2.2	7
191	Clinical and radiographic outcome of a treat-to-target strategy using methotrexate and intra-articular glucocorticoids with or without adalimumab induction: a 2-year investigator-initiated, double-blinded, randomised, controlled trial (OPERA). <i>Annals of the Rheumatic Diseases</i> , 2017 , 75, 1415-1423	2.4	30
190	Prediction of treatment response to adalimumab: a double-blind placebo-controlled study of circulating microRNA in patients with early rheumatoid arthritis. <i>Pharmacogenomics Journal</i> , 2016 , 16, 141-6	3.5	44
189	Full dose, reduced dose or discontinuation of etanercept in rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2016 , 75, 52-8	2.4	88
188	MRI assessment of suppression of structural damage in patients with rheumatoid arthritis receiving rituximab: results from the randomised, placebo-controlled, double-blind RA-SCORE study. <i>Annals of the Rheumatic Diseases</i> , 2016 , 75, 170-7	2.4	36
187	Imaging: MRI in healthy volunteers - important to do, and do correctly. <i>Nature Reviews Rheumatology</i> , 2016 , 12, 563-4	8.1	9

186	Validity and sensitivity to change of the semi-quantitative OMERACT ultrasound scoring system for tenosynovitis in patients with rheumatoid arthritis. <i>Rheumatology</i> , 2016 , 55, 2156-2166	3.9	18
185	Imaging in the diagnosis and management of peripheral psoriatic arthritis-The clinical utility of magnetic resonance imaging and ultrasonography. <i>Best Practice and Research in Clinical Rheumatology</i> , 2016 , 30, 624-637	5.3	24
184	FRI0535 Hand Bone Loss in Early Rheumatoid Arthritis Is Independent of Adalimumab Treatment. A Substudy of The Optimized Treatment Algorithm in Early RA (Opera) Trial. <i>Annals of the Rheumatic Diseases</i> , 2016 , 75, 633.1-633	2.4	
183	AB0036 Soluble CD83 Plasma Levels Are Associated with Disease Activity and Course of Disease in Early Rheumatoid Arthritis. <i>Annals of the Rheumatic Diseases</i> , 2016 , 75, 909.1-909	2.4	
182	THU0635 Defining The Optimal Biological Monotherapy in Rheumatoid Arthritis: A Systematic Review and Network Meta-Analysis of Randomised Trials. <i>Annals of the Rheumatic Diseases</i> , 2016 , 75, 423.3-424	2.4	
181	AB0962 Validity and Sensitivity To Change of The Semi-Quantitative Omeract Ultrasound Scoring System for Tenosynovitis in Patients with Rheumatoid Arthrit. <i>Annals of the Rheumatic Diseases</i> , 2016 , 75, 1230.1-1230	2.4	
180	AB0663 Efficacy and Safety of Adalimumab in Patients with Axial Spondyloarthritis - An Investigator-Initiated Randomized Placebo-Controlled Trial. <i>Annals of the Rheumatic Diseases</i> , 2016 , 75, 1131.3-1132	2.4	
179	OP0125 A Randomised, Double-Blind, Controlled Study of Intramuscular versus Ultrasound Guided Peritendinous Betamethasone Injection for Tenosynovitis in Rheumatoid Arthritis Patients. <i>Annals of the Rheumatic Diseases</i> , 2016 , 75, 103.1-103	2.4	
178	AB0961 Comparison between Ultrasound B-Flow, Colour Doppler and Magnetic Resonance Imaging for Assessment of Tenosynovitis in The Hand of Rheumatoid Arthritis Patients - A Pilot Study. <i>Annals of the Rheumatic Diseases</i> , 2016 , 75, 1229.3-1230	2.4	
177	Establishment of age- and sex-adjusted reference data for hand bone mass and investigation of hand bone loss in patients with rheumatoid arthritis treated in clinical practice: an observational study from the DANBIO registry and the Copenhagen Osteoarthritis Study. <i>Arthritis Research and Therapy</i> , 2016 , 18, 53	5.7	4
176	The interleukin-20 receptor axis in early rheumatoid arthritis: novel links between disease-associated autoantibodies and radiographic progression. <i>Arthritis Research and Therapy</i> , 2016 , 18, 61	5.7	24
175	Defining active sacroiliitis on MRI for classification of axial spondyloarthritis: update by the ASAS MRI working group. <i>Annals of the Rheumatic Diseases</i> , 2016 , 75, 1958-1963	2.4	259
174	Validity of early MRI structural damage end points and potential impact on clinical trial design in rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2016 , 75, 1114-9	2.4	21
173	Three-dimensional Doppler ultrasound findings in healthy wrist and finger tendon sheaths - can feeding vessels lead to misinterpretation in Doppler-detected tenosynovitis?. <i>Arthritis Research and Therapy</i> , 2016 , 18, 70	5.7	2
172	Sensitivity and specificity of optical spectral transmission imaging in detecting joint inflammation in rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2016 , 75, 632-3	2.4	7
171	EFSUMB COMPASS for Rheumatologists dissemination and implementation--an international survey. <i>Medical Ultrasonography</i> , 2016 , 18, 42-6	1.4	1
170	FRI0548 Responsiveness of A New MRI Scoring Method Based on The Canada-Denmark Definitions of Lesions in The Spine and The SPARCC MRI Spine Inflammation Index in Patients with Axial Spondyloarthritis. <i>Annals of the Rheumatic Diseases</i> , 2016 , 75, 638.3-639	2.4	
169	SAT0610 The Prospective Risk for Long Term Sickness Absence, Unemployment, and Disability Pension, and The Probability for Return To Work in Patients with Rheumatoid Arthritis. <i>Annals of the Rheumatic Diseases</i> , 2016 , 75, 890.2-890	2.4	

168	Safety of Repeated Open-Label Treatment Courses of Intravenous Ofatumumab, a Human Anti-CD20 Monoclonal Antibody, in Rheumatoid Arthritis: Results from Three Clinical Trials. <i>PLoS ONE</i> , 2016 , 11, e0157961	3.7	14
167	A disintegrin and metalloprotease-17 and galectin-9 are important regulators of local 4-1BB activity and disease outcome in rheumatoid arthritis. <i>Rheumatology</i> , 2016 , 55, 1871-9	3.9	9
166	Course of Magnetic Resonance Imaging-Detected Inflammation and Structural Lesions in the Sacroiliac Joints of Patients in the Randomized, Double-Blind, Placebo-Controlled Danish Multicenter Study of Adalimumab in Spondyloarthritis, as Assessed by the Berlin and Spondyloarthritis Research Consortium of Canada Methods. <i>Arthritis and Rheumatology</i> , 2016 , 68, 418-29	9.5	32
165	SAT0559 The Anatomical Distribution of Inflammation, Fat, Erosion and New Bone Formation in The Spine Assessed According To The Canada-Denmark MRI Definitions in Patients with Axial Spondyloarthritis. <i>Annals of the Rheumatic Diseases</i> , 2016 , 75, 872.1-872	2.4	
164	Value of ultrasonography as a marker of early response to abatacept in patients with rheumatoid arthritis and an inadequate response to methotrexate: results from the APPRAISE study. <i>Annals of the Rheumatic Diseases</i> , 2016 , 75, 1763-9	2.4	54
163	Exploring a new ultrasound score as a clinical predictive tool in patients with rheumatoid arthritis starting abatacept: results from the APPRAISE study. <i>RMD Open</i> , 2016 , 2, e000237	5.9	44
162	AB0263 Agreement between Das28, Acr/eular, Sdai, Cdai and Ultrasound Remission in Patients with Rheumatoid Arthritis Receiving Biological Treatment in Routine Care. <i>Annals of the Rheumatic Diseases</i> , 2016 , 75, 989.2-990	2.4	
161	Novel algorithms for the pragmatic use of ultrasound in the management of patients with rheumatoid arthritis: from diagnosis to remission. <i>Annals of the Rheumatic Diseases</i> , 2016 , 75, 1902-1908	2.4	64
160	Efficacy of VX-509 (decernotinib) in combination with a disease-modifying antirheumatic drug in patients with rheumatoid arthritis: clinical and MRI findings. <i>Annals of the Rheumatic Diseases</i> , 2016 , 75, 1979-1983	2.4	36
159	Recommendations of the ESSR Arthritis Subcommittee on Ultrasonography in Inflammatory Joint Disease. <i>Seminars in Musculoskeletal Radiology</i> , 2016 , 20, 496-506	1.8	16
158	Comparing the effects of tofacitinib, methotrexate and the combination, on bone marrow oedema, synovitis and bone erosion in methotrexate-naive, early active rheumatoid arthritis: results of an exploratory randomised MRI study incorporating semiquantitative and quantitative techniques. <i>Annals of the Rheumatic Diseases</i> , 2016 , 75, 1024-33	2.4	72
157	FRI0521 Reliability of An Omeract Rheumatoid Arthritis Tenosynovitis Scoring System for Wrist and Hand. <i>Annals of the Rheumatic Diseases</i> , 2016 , 75, 627.3-628	2.4	
156	Effect of Rheumatoid Arthritis on Longterm Sickness Absence in 1994-2011: A Danish Cohort Study. <i>Journal of Rheumatology</i> , 2016 , 43, 707-15	4.1	15
155	Simulation-Based Abdominal Ultrasound Training - A Systematic Review. <i>Ultraschall in Der Medizin</i> , 2016 , 37, 253-61	3.8	19
154	Aiming for a simpler early arthritis MRI protocol: can Gd contrast administration be eliminated?. <i>European Radiology</i> , 2015 , 25, 1520-7	8	32
153	Head-to-toe whole-body MRI in psoriatic arthritis, axial spondyloarthritis and healthy subjects: first steps towards global inflammation and damage scores of peripheral and axial joints. <i>Rheumatology</i> , 2015 , 54, 1039-49	3.9	41
152	The efficacy of motivational counseling and SMS-reminders on daily sitting time in patients with rheumatoid arthritis: protocol for a randomized controlled trial. <i>Trials</i> , 2015 , 16, 23	2.8	12
151	Physical activity and the association with fatigue and sleep in Danish patients with rheumatoid arthritis. <i>Rheumatology International</i> , 2015 , 35, 1655-64	3.6	33

150	International Consensus for ultrasound lesions in gout: results of Delphi process and web-reliability exercise. <i>Rheumatology</i> , 2015 , 54, 1797-805	3.9	94
149	Sleep quality and correlates of poor sleep in patients with rheumatoid arthritis. <i>Clinical Rheumatology</i> , 2015 , 34, 2029-39	3.9	52
148	EULAR recommendations for the use of imaging in the diagnosis and management of spondyloarthritis in clinical practice. <i>Annals of the Rheumatic Diseases</i> , 2015 , 74, 1327-39	2.4	284
147	Enthesitis in patients with psoriatic arthritis, axial spondyloarthritis and healthy subjects assessed by 'head-to-toe' whole-body MRI and clinical examination. <i>Annals of the Rheumatic Diseases</i> , 2015 , 74, 823-9	2.4	81
146	Assessing Elementary Lesions in Gout by Ultrasound: Results of an OMERACT Patient-based Agreement and Reliability Exercise. <i>Journal of Rheumatology</i> , 2015 , 42, 2149-54	4.1	45
145	Does evaluation of the ligamentous compartment enhance diagnostic utility of sacroiliac joint MRI in axial spondyloarthritis?. <i>Arthritis Research and Therapy</i> , 2015 , 17, 246	5.7	11
144	Effectiveness and drug adherence of biologic monotherapy in routine care of patients with rheumatoid arthritis: a cohort study of patients registered in the Danish biologics registry. <i>Rheumatology</i> , 2015 , 54, 2156-65	3.9	26
143	The Longitudinal Reliability and Responsiveness of the OMERACT Hand Osteoarthritis Magnetic Resonance Imaging Scoring System (HOAMRIS). <i>Journal of Rheumatology</i> , 2015 , 42, 2486-91	4.1	20
142	Imaging in Psoriatic Arthritis. <i>Rheumatic Disease Clinics of North America</i> , 2015 , 41, 593-613	2.4	25
141	EULAR-PReS points to consider for the use of imaging in the diagnosis and management of juvenile idiopathic arthritis in clinical practice. <i>Annals of the Rheumatic Diseases</i> , 2015 , 74, 1946-57	2.4	79
140	Power and color Doppler ultrasound settings for inflammatory flow: impact on scoring of disease activity in patients with rheumatoid arthritis. <i>Arthritis and Rheumatology</i> , 2015 , 67, 386-95	9.5	66
139	Does spinal MRI add incremental diagnostic value to MRI of the sacroiliac joints alone in patients with non-radiographic axial spondyloarthritis?. <i>Annals of the Rheumatic Diseases</i> , 2015 , 74, 985-92	2.4	65
138	MOR103, a human monoclonal antibody to granulocyte-macrophage colony-stimulating factor, in the treatment of patients with moderate rheumatoid arthritis: results of a phase Ib/IIa randomised, double-blind, placebo-controlled, dose-escalation trial. <i>Annals of the Rheumatic Diseases</i> , 2015 , 74, 1058-64	2.4	106
137	Candidate lesion-based criteria for defining a positive sacroiliac joint MRI in two cohorts of patients with axial spondyloarthritis. <i>Annals of the Rheumatic Diseases</i> , 2015 , 74, 1976-82	2.4	65
136	SAT0222 Effects of Tofacitinib on MRI Endpoints in Methotrexate-Naive Early Rheumatoid Arthritis: A Phase 2 MRI Study with Semi-Quantitative and Quantitative Endpoints. <i>Annals of the Rheumatic Diseases</i> , 2015 , 74, 738-738	2.4	4
135	OP0256 ADAM17 And Galectin-9 are Critical Regulators of Local 4-1BB Activity and Disease Outcome in Rheumatoid Arthritis. <i>Annals of the Rheumatic Diseases</i> , 2015 , 74, 168.3-169	2.4	
134	FRI0194 Is There an Association Between Spondyloarthritis and Antibodies Towards Borrelia, Ehrlichia and Chlamydia Species?. <i>Annals of the Rheumatic Diseases</i> , 2015 , 74, 495.1-495	2.4	1
133	FRI0503 Eular-PreS Points to Consider for the Use of Imaging in the Diagnosis and Management of Juvenile Idiopathic Arthritis in Clinical Practice. <i>Annals of the Rheumatic Diseases</i> , 2015 , 74, 611.1-611	2.4	

132	FRI0598 3D Ultrasound Doppler Findings in Wrist Tendon Sheaths of Healthy Controls. <i>Annals of the Rheumatic Diseases</i> , 2015 , 74, 644.2-645	2.4	
131	OP0073 Polymorphisms in the FCN1 Gene Coding for M-Ficolin are Associated with Disease Activity, Radiographic Damage and are the Strongest Predictors of DAS28 Remission in 180 DMARD Naive Early Rheumatoid Arthritis Patients. <i>Annals of the Rheumatic Diseases</i> , 2015 , 74, 96.1-96	2.4	
130	SAT0003 The IL-20 Receptor Axis in Early Rheumatoid Arthritis: Novel Inflammation-Independent Links Between Rheumatoid Arthritis-Associated Autoantibodies and Radiographic Progression. <i>Annals of the Rheumatic Diseases</i> , 2015 , 74, 650.3-651	2.4	
129	THU0498 The Patients' Experience of Imaging: Views from a Group Convened to Support the Development of Points to Consider for the Use of Imaging in the Diagnosis and Management of Juvenile Idiopathic Arthritis. <i>Annals of the Rheumatic Diseases</i> , 2015 , 74, 380.2-380	2.4	
128	AB1076 Ultrasound Abnormalities at the Entheses are not an Uncommon Finding in Patients with Rheumatoid Arthritis. <i>Annals of the Rheumatic Diseases</i> , 2015 , 74, 1260.1-1260	2.4	
127	Adherence to methotrexate in rheumatoid arthritis: a danish nationwide cohort study. <i>Arthritis</i> , 2015 , 2015, 915142		10
126	MRI assessment of early response to certolizumab pegol in rheumatoid arthritis: a randomised, double-blind, placebo-controlled phase IIIb study applying MRI at weeks 0, 1, 2, 4, 8 and 16. <i>Annals of the Rheumatic Diseases</i> , 2015 , 74, 1156-63	2.4	23
125	Validation of the OMERACT Magnetic Resonance Imaging Joint Space Narrowing Score for the Wrist in a Multireader Longitudinal Trial. <i>Journal of Rheumatology</i> , 2015 , 42, 2480-5	4.1	23
124	The OMERACT MRI in Arthritis Working Group - Update on Status and Future Research Priorities. <i>Journal of Rheumatology</i> , 2015 , 42, 2470-2	4.1	10
123	Impact of a magnetic resonance imaging-guided treat-to-target strategy on disease activity and progression in patients with rheumatoid arthritis (the IMAGINE-RA trial): study protocol for a randomized controlled trial. <i>Trials</i> , 2015 , 16, 178	2.8	9
122	A treat-to-target strategy with methotrexate and intra-articular triamcinolone with or without adalimumab effectively reduces MRI synovitis, osteitis and tenosynovitis and halts structural damage progression in early rheumatoid arthritis: results from the OPERA randomised controlled trial. <i>Annals of the Rheumatic Diseases</i> , 2015 , 74, 867-75	2.4	44
121	Recommendations of the ESSR Arthritis Subcommittee for the Use of Magnetic Resonance Imaging in Musculoskeletal Rheumatic Diseases. <i>Seminars in Musculoskeletal Radiology</i> , 2015 , 19, 396-411	1.8	76
120	AB1084 Correlations Between Different Imaging Modalities in the Assessment of Acute Synovitis of the Wrist in Rheumatoid Arthritis: A Cross-Sectional Study. <i>Annals of the Rheumatic Diseases</i> , 2015 , 74, 1263.1-1263	2.4	
119	Validation of the OMERACT Psoriatic Arthritis Magnetic Resonance Imaging Score (PsAMRIS) for the Hand and Foot in a Randomized Placebo-controlled Trial. <i>Journal of Rheumatology</i> , 2015 , 42, 2473-9	4.1	41
118	Macrophage activity assessed by soluble CD163 in early rheumatoid arthritis: association with disease activity but different response patterns to synthetic and biologic DMARDs. <i>Clinical and Experimental Rheumatology</i> , 2015 , 33, 498-502	2.2	14
117	Magnetic resonance imaging for diagnosing, monitoring and prognostication in psoriatic arthritis. <i>Clinical and Experimental Rheumatology</i> , 2015 , 33, S66-9	2.2	11
116	Whole-body MRI assessment of disease activity and structural damage in rheumatoid arthritis: first step towards an MRI joint count. <i>Rheumatology</i> , 2014 , 53, 845-53	3.9	33
115	Development and validation of modified disease activity scores in rheumatoid arthritis: superior correlation with magnetic resonance imaging-detected synovitis and radiographic progression. <i>Arthritis and Rheumatology</i> , 2014 , 66, 794-802	9.5	43

114	Bone marrow oedema assessment by magnetic resonance imaging in rheumatoid arthritis wrist and metacarpophalangeal joints: the importance of field strength, coil type and image resolution. <i>Rheumatology</i> , 2014 , 53, 1446-51	3.9	10
113	Early MRI measures independently predict 1-year and 2-year radiographic progression in rheumatoid arthritis: secondary analysis from a large clinical trial. <i>Annals of the Rheumatic Diseases</i> , 2014 , 73, 1968-74	2.4	61
112	Determining a magnetic resonance imaging inflammatory activity acceptable state without subsequent radiographic progression in rheumatoid arthritis: results from a followup MRI study of 254 patients in clinical remission or low disease activity. <i>Journal of Rheumatology</i> , 2014 , 41, 398-406	4.1	61
111	Reply: To PMID 24757132. <i>Arthritis and Rheumatology</i> , 2014 , 66, 3245-6	9.5	
110	The OMERACT-RAMRIS rheumatoid arthritis magnetic resonance imaging joint space narrowing score: intrareader and interreader reliability and agreement with computed tomography and conventional radiography. <i>Journal of Rheumatology</i> , 2014 , 41, 392-7	4.1	31
109	Fat infiltration on magnetic resonance imaging of the sacroiliac joints has limited diagnostic utility in nonradiographic axial spondyloarthritis. <i>Journal of Rheumatology</i> , 2014 , 41, 75-83	4.1	36
108	SAT0216 Sensitivity and Specificity of Optical Spectral Transmission Imaging in Detecting Joint Inflammation in Rheumatoid Arthritis. <i>Annals of the Rheumatic Diseases</i> , 2014 , 73, 668.1-668	2.4	
107	FRI0183 Adherence to Methotrexate in Rheumatoid Arthritis Treated by Practice Vs. Hospital Based Rheumatologists: A Danish Nationwide Cohort Study. <i>Annals of the Rheumatic Diseases</i> , 2014 , 73, 448.2-449	2.4	
106	CXCL13 predicts disease activity in early rheumatoid arthritis and could be an indicator of the therapeutic 'window of opportunity'. <i>Arthritis Research and Therapy</i> , 2014 , 16, 434	5.7	47
105	THU0550 Changes in Soluble CD18 Reflect Latency in Restoration of the Immune System after Early Treatment of Rheumatoid Arthritis. <i>Annals of the Rheumatic Diseases</i> , 2014 , 73, 373.1-373	2.4	
104	Which factors influence radiographic progression during treatment with tumor necrosis factor inhibitors in clinical practice? Results from 930 patients with rheumatoid arthritis in the nationwide Danish DANBIO registry. <i>Journal of Rheumatology</i> , 2014 , 41, 2352-60	4.1	9
103	Adalimumab added to a treat-to-target strategy with methotrexate and intra-articular triamcinolone in early rheumatoid arthritis increased remission rates, function and quality of life. The OPERA Study: an investigator-initiated, randomised, double-blind, parallel-group, placebo-controlled trial. <i>Annals of the Rheumatic Diseases</i> , 2014 , 73, 454-461	2.4	84
102	Optimal use of MRI in clinical trials, clinical care and clinical registries of patients with rheumatoid arthritis. <i>Clinical and Experimental Rheumatology</i> , 2014 , 32, S-17-22	2.2	5
101	Reliability of a consensus-based ultrasound score for tenosynovitis in rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2013 , 72, 1328-34	2.4	170
100	EULAR recommendations for the use of imaging of the joints in the clinical management of rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2013 , 72, 804-14	2.4	376
99	The FAT Spondyloarthritis Spine Score (FASSS): development and validation of a new scoring method for the evaluation of fat lesions in the spine of patients with axial spondyloarthritis. <i>Arthritis Research and Therapy</i> , 2013 , 15, R216	5.7	6
98	Review: the utility of magnetic resonance imaging for assessing structural damage in randomized controlled trials in rheumatoid arthritis. <i>Arthritis and Rheumatism</i> , 2013 , 65, 2513-23		27
97	THU0435 Detection of synovitis, bone erosions, and bone marrow oedema in patients with inflammatory hand pain - a comparison of low field MRI and high field MRI. <i>Annals of the Rheumatic Diseases</i> , 2013 , 71, 302.2-302	2.4	

96	Power Doppler ultrasonography of painful Achilles tendons and entheses in patients with and without spondyloarthritis: a comparison with clinical examination and contrast-enhanced MRI. <i>Clinical Rheumatology</i> , 2013 , 32, 301-8	3.9	18
95	Clinical response, drug survival, and predictors thereof among 548 patients with psoriatic arthritis who switched tumor necrosis factor inhibitor therapy: results from the Danish Nationwide DANBIO Registry. <i>Arthritis and Rheumatism</i> , 2013 , 65, 1213-23		127
94	MRI comes of age in RA clinical trials. <i>Annals of the Rheumatic Diseases</i> , 2013 , 72, 794-6	2.4	35
93	AB0316 Patient global assessment of general health by vas at baseline predicts ACR/EULAR remission after 3, 6 and 12 months of efficient treatment in DMARD- and steroid naïve early rheumatoid arthritis patients. The danish cimestra-study. <i>Annals of the Rheumatic Diseases</i> , 2013 , 72, A666.2-A667	2.4	
92	SAT0250 Validation of a New Scoring Method for the Evaluation of Fat Lesions in the Spine of Patients with Axial Spondyloarthritis: The Canada-Denmark Fat Spondyloarthritis Spine Score (FASSS). <i>Annals of the Rheumatic Diseases</i> , 2013 , 72, A666.2-A667	2.4	
91	OP0168 Clinical response, drug survival and predictors thereof in 432 patients with ankylosing spondylitis switching anti tumor necrosis factor therapy: Results from the danish nationwide danbio registry. <i>Annals of the Rheumatic Diseases</i> , 2013 , 71, 111.2-111	2.4	3
90	OP0223 M-ficolin, an activator of the complement system, predicts DAS28 remission in early DMARD naïve rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2013 , 71, 131.1-131	2.4	1
89	SAT0130 Defining the Optimal Biological Monotherapy in Rheumatoid Arthritis (RA): Network Meta-Analysis of Randomized Trials. <i>Annals of the Rheumatic Diseases</i> , 2013 , 72, A625.1-A625	2.4	
88	SAT0523 Prediction of Clinical Response to Biologics in Rheumatoid Arthritis by Baseline Magnetic Resonance Imaging (MRI) of the Wrist: a Cohort Study Associating MRI Synovitis Score with Clinical Outcome. <i>Annals of the Rheumatic Diseases</i> , 2013 , 72, A758.4-A759	2.4	
87	AB0304 Predictors of gain in quality adjusted life years in RA patients treated with biologics for one year. <i>Annals of the Rheumatic Diseases</i> , 2013 , 71, 655.1-655	2.4	
86	Imaging in rheumatology: new tools for use in clinical practice in 2012. Preface. <i>Best Practice and Research in Clinical Rheumatology</i> , 2012 , 26, 743-9	5.3	2
85	The diagnostic utility of MRI in spondyloarthritis. <i>Best Practice and Research in Clinical Rheumatology</i> , 2012 , 26, 751-66	5.3	13
84	Can erosions on MRI of the sacroiliac joints be reliably detected in patients with ankylosing spondylitis? - A cross-sectional study. <i>Arthritis Research and Therapy</i> , 2012 , 14, R124	5.7	71
83	CD6 and syntaxin binding protein 6 variants and response to tumor necrosis factor alpha inhibitors in Danish patients with rheumatoid arthritis. <i>PLoS ONE</i> , 2012 , 7, e38539	3.7	13
82	Descriptions of spinal MRI lesions and definition of a positive MRI of the spine in axial spondyloarthritis: a consensual approach by the ASAS/OMERACT MRI study group. <i>Annals of the Rheumatic Diseases</i> , 2012 , 71, 1278-88	2.4	162
81	Imaging in ankylosing spondylitis. <i>Therapeutic Advances in Musculoskeletal Disease</i> , 2012 , 4, 301-11	3.8	34
80	Dynamic gadolinium-enhanced magnetic resonance imaging allows accurate assessment of the synovial inflammatory activity in rheumatoid arthritis knee joints: a comparison with synovial histology. <i>Scandinavian Journal of Rheumatology</i> , 2012 , 41, 89-94	1.9	55
79	Practice of ultrasound-guided arthrocentesis and joint injection, including training and implementation, in Europe: results of a survey of experts and scientific societies. <i>Rheumatology</i> , 2012 , 51, 184-90	3.9	35

78	Can imaging be used for inflammatory arthritis screening?. <i>Seminars in Musculoskeletal Radiology</i> , 2012 , 16, 401-9	1.8	12
77	Development of standardized approaches to reporting of minimal residual disease data using a reporting software package designed within the European LeukemiaNet. <i>Leukemia</i> , 2011 , 25, 1168-73	10.7	20
76	The impact of MRI on the clinical management of inflammatory arthritides. <i>Skeletal Radiology</i> , 2011 , 40, 1153-73	2.7	33
75	Bone edema on magnetic resonance imaging is an independent predictor of rheumatoid arthritis development in patients with early undifferentiated arthritis. <i>Arthritis and Rheumatism</i> , 2011 , 63, 2192-202		79
74	Significant improvement in synovitis, osteitis, and bone erosion following golimumab and methotrexate combination therapy as compared with methotrexate alone: a magnetic resonance imaging study of 318 methotrexate-naïve rheumatoid arthritis patients. <i>Arthritis and Rheumatism</i> , 2011 , 63, 2719-22		71
73	Radiographic progression is associated with resolution of systemic inflammation in patients with axial spondylarthritis treated with tumor necrosis factor inhibitors: a study of radiographic progression, inflammation on magnetic resonance imaging, and circulating biomarkers of inflammation. <i>Arthritis and Rheumatism</i> , 2011 , 63, 2789-800		67
72	Synovitis and osteitis are very frequent in rheumatoid arthritis clinical remission: results from an MRI study of 294 patients in clinical remission or low disease activity state. <i>Journal of Rheumatology</i> , 2011 , 38, 2039-44	4.1	85
71	Recent advances in imaging in psoriatic arthritis. <i>Therapeutic Advances in Musculoskeletal Disease</i> , 2011 , 3, 43-53	3.8	19
70	MRI in early rheumatoid arthritis: synovitis and bone marrow oedema are independent predictors of subsequent radiographic progression. <i>Annals of the Rheumatic Diseases</i> , 2011 , 70, 428-33	2.4	108
69	Development and preliminary validation of a magnetic resonance imaging joint space narrowing score for use in rheumatoid arthritis: potential adjunct to the OMERACT RA MRI scoring system. <i>Journal of Rheumatology</i> , 2011 , 38, 2045-50	4.1	64
68	The OMERACT Psoriatic Arthritis Magnetic Resonance Imaging Score (PsAMRIS) is reliable and sensitive to change: results from an OMERACT workshop. <i>Journal of Rheumatology</i> , 2011 , 38, 2034-8	4.1	55
67	Resolution of inflammation following treatment of ankylosing spondylitis is associated with new bone formation. <i>Journal of Rheumatology</i> , 2011 , 38, 1349-54	4.1	78
66	Patterns of magnetic resonance imaging bone erosion in rheumatoid arthritis--which bones are most frequently involved and show the most change?. <i>Journal of Rheumatology</i> , 2011 , 38, 2014-7	4.1	12
65	Radiographic progression and remission rates in early rheumatoid arthritis - MRI bone oedema and anti-CCP predicted radiographic progression in the 5-year extension of the double-blind randomised CIMESTR trial. <i>Annals of the Rheumatic Diseases</i> , 2010 , 69, 1789-95	2.4	136
64	Responsiveness of the Ankylosing Spondylitis Disease Activity Score (ASDAS) and clinical and MRI measures of disease activity in a 1-year follow-up study of patients with axial spondyloarthritis treated with tumour necrosis factor alpha inhibitors. <i>Annals of the Rheumatic Diseases</i> , 2010 , 69, 1065-71	2.4	88
63	Current state of musculoskeletal ultrasound training and implementation in Europe: results of a survey of experts and scientific societies. <i>Rheumatology</i> , 2010 , 49, 2438-43	3.9	47
62	Circulating levels of interleukin-6, vascular endothelial growth factor, YKL-40, matrix metalloproteinase-3, and total aggrecan in spondyloarthritis patients during 3 years of treatment with TNF inhibitors. <i>Clinical Rheumatology</i> , 2010 , 29, 1301-9	3.9	52
61	Magnetic resonance imaging in spondyloarthritis--how to quantify findings and measure response. <i>Best Practice and Research in Clinical Rheumatology</i> , 2010 , 24, 637-57	5.3	32

60	Assessment of structural lesions in sacroiliac joints enhances diagnostic utility of magnetic resonance imaging in early spondylarthritis. <i>Arthritis Care and Research</i> , 2010 , 62, 1763-71	4.7	93
59	Ofatumumab, a human anti-CD20 monoclonal antibody, for treatment of rheumatoid arthritis with an inadequate response to one or more disease-modifying antirheumatic drugs: results of a randomized, double-blind, placebo-controlled, phase I/II study. <i>Arthritis and Rheumatism</i> , 2010 , 62, 2227-38		73
58	The diagnostic utility of magnetic resonance imaging in spondylarthritis: an international multicenter evaluation of one hundred eighty-seven subjects. <i>Arthritis and Rheumatism</i> , 2010 , 62, 3048-58		204
57	Monitoring anti-TNFalpha treatment in rheumatoid arthritis: responsiveness of magnetic resonance imaging and ultrasonography of the dominant wrist joint compared with conventional measures of disease activity and structural damage. <i>Annals of the Rheumatic Diseases</i> , 2009 , 68, 1572-9	2.4	72
56	MRI bone oedema is the strongest predictor of subsequent radiographic progression in early rheumatoid arthritis. Results from a 2-year randomised controlled trial (CIMESTRA). <i>Annals of the Rheumatic Diseases</i> , 2009 , 68, 384-90	2.4	299
55	Testing an OMERACT MRI scoring system for peripheral psoriatic arthritis in cross-sectional and longitudinal settings. <i>Journal of Rheumatology</i> , 2009 , 36, 1811-5	4.1	26
54	Proposal for levels of evidence schema for validation of a soluble biomarker reflecting damage endpoints in rheumatoid arthritis, psoriatic arthritis, and ankylosing spondylitis, and recommendations for study design. <i>Journal of Rheumatology</i> , 2009 , 36, 1792-9	4.1	30
53	The OMERACT psoriatic arthritis magnetic resonance imaging scoring system (PsAMRIS): definitions of key pathologies, suggested MRI sequences, and preliminary scoring system for PsA Hands. <i>Journal of Rheumatology</i> , 2009 , 36, 1816-24	4.1	138
52	Reappraisal of OMERACT 8 draft validation criteria for a soluble biomarker reflecting structural damage endpoints in rheumatoid arthritis, psoriatic arthritis, and spondyloarthritis: the OMERACT 9 v2 criteria. <i>Journal of Rheumatology</i> , 2009 , 36, 1785-91	4.1	35
51	Testing of the OMERACT 8 draft validation criteria for a soluble biomarker reflecting structural damage in rheumatoid arthritis: a systematic literature search on 5 candidate biomarkers. <i>Journal of Rheumatology</i> , 2009 , 36, 1769-84	4.1	15
50	Erosive progression is minimal, but erosion healing rare, in patients with rheumatoid arthritis treated with adalimumab. A 1 year investigator-initiated follow-up study using high-resolution computed tomography as the primary outcome measure. <i>Annals of the Rheumatic Diseases</i> , 2009 , 68, 1585-90	2.4	64
49	Reducing invasiveness, duration, and cost of magnetic resonance imaging in rheumatoid arthritis by omitting intravenous contrast injection -- Does it change the assessment of inflammatory and destructive joint changes by the OMERACT RAMRIS?. <i>Journal of Rheumatology</i> , 2009 , 36, 1806-10	4.1	82
48	Inflammatory lesions of the spine on magnetic resonance imaging predict the development of new syndesmophytes in ankylosing spondylitis: evidence of a relationship between inflammation and new bone formation. <i>Arthritis and Rheumatism</i> , 2009 , 60, 93-102		254
47	Imaging in rheumatoid arthritis--status and recent advances for magnetic resonance imaging, ultrasonography, computed tomography and conventional radiography. <i>Best Practice and Research in Clinical Rheumatology</i> , 2008 , 22, 1019-44	5.3	100
46	Circulating levels of osteopontin, osteoprotegerin, total soluble receptor activator of nuclear factor-kappa B ligand, and high-sensitivity C-reactive protein in patients with active rheumatoid arthritis randomized to etanercept alone or in combination with methotrexate. <i>Scandinavian Journal of Rheumatology</i> , 2008 , 37, 241-7	1.9	38
45	Detection of bone erosions in rheumatoid arthritis wrist joints with magnetic resonance imaging, computed tomography and radiography. <i>Arthritis Research and Therapy</i> , 2008 , 10, R25	5.7	104
44	Magnetic resonance imaging findings in 84 patients with early rheumatoid arthritis: bone marrow oedema predicts erosive progression. <i>Annals of the Rheumatic Diseases</i> , 2008 , 67, 794-800	2.4	210
43	Diagnostic sensitivity and specificity of Doppler ultrasound in rheumatoid arthritis. <i>Journal of Rheumatology</i> , 2008 , 35, 49-53	4.1	39

42	Low remission rates but long drug survival in rheumatoid arthritis patients treated with infliximab or etanercept: results from the nationwide Danish DANBIO database. <i>Scandinavian Journal of Rheumatology</i> , 2007 , 36, 151-4	1.9	34
41	Reply to Wilms' tumour 1 mutations are associated with FLT3-ITD and failure of standard induction chemotherapy in patients with normal karyotype AML by Summers et al.. <i>Leukemia</i> , 2007 , 21, 552-552	10.7	
40	Established rheumatoid arthritis - new imaging modalities. <i>Best Practice and Research in Clinical Rheumatology</i> , 2007 , 21, 841-56	5.3	43
39	No erosive progression revealed by MRI in rheumatoid arthritis patients treated with etanercept, even in patients with persistent MRI and clinical signs of joint inflammation. <i>Clinical Rheumatology</i> , 2007 , 26, 1857-61	3.9	22
38	Ultrasonography, magnetic resonance imaging, radiography, and clinical assessment of inflammatory and destructive changes in fingers and toes of patients with psoriatic arthritis. <i>Arthritis Research and Therapy</i> , 2007 , 9, R119	5.7	136
37	Biological variation and reference intervals for circulating osteopontin, osteoprotegerin, total soluble receptor activator of nuclear factor kappa B ligand and high-sensitivity C-reactive protein. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2007 , 67, 821-35	2	26
36	The OMERACT Magnetic Resonance Imaging Inflammatory Arthritis Group - advances and priorities. <i>Journal of Rheumatology</i> , 2007 , 34, 852-3	4.1	8
35	Developing a magnetic resonance imaging scoring system for peripheral psoriatic arthritis. <i>Journal of Rheumatology</i> , 2007 , 34, 859-61	4.1	34
34	Scoring inflammatory activity of the spine by magnetic resonance imaging in ankylosing spondylitis: a multireader experiment. <i>Journal of Rheumatology</i> , 2007 , 34, 862-70	4.1	80
33	Ultrasonography and magnetic resonance imaging in early rheumatoid arthritis: recent advances. <i>Current Rheumatology Reports</i> , 2006 , 8, 378-85	4.9	14
32	Combination treatment with methotrexate, cyclosporine, and intraarticular betamethasone compared with methotrexate and intraarticular betamethasone in early active rheumatoid arthritis: an investigator-initiated, multicenter, randomized, double-blind, parallel-group, placebo-controlled study. <i>Arthritis and Rheumatism</i> , 2006 , 54, 1401-9		129
31	Does digital X-ray radiogrammetry help assess joint damage in patients with rheumatoid arthritis?. <i>Nature Clinical Practice Rheumatology</i> , 2006 , 2, 18-9		
30	Are bone erosions detected by magnetic resonance imaging and ultrasonography true erosions? A comparison with computed tomography in rheumatoid arthritis metacarpophalangeal joints. <i>Arthritis Research and Therapy</i> , 2006 , 8, R110	5.7	168
29	Drug survival and reasons for discontinuation of intramuscular methotrexate: a study of 212 consecutive patients switching from oral methotrexate. <i>Scandinavian Journal of Rheumatology</i> , 2006 , 35, 102-6	1.9	13
28	Plasma IL-6, plasma VEGF, and serum YKL-40: relationship with disease activity and radiographic progression in rheumatoid arthritis patients treated with infliximab and methotrexate. <i>Scandinavian Journal of Rheumatology</i> , 2006 , 35, 489-91	1.9	24
27	Cardiac output--pulse contour analysis vs. pulmonary artery thermodilution. <i>Acta Anaesthesiologica Scandinavica</i> , 2006 , 50, 1044-9	1.9	31
26	An introduction to the EULAR-OMERACT rheumatoid arthritis MRI reference image atlas. <i>Annals of the Rheumatic Diseases</i> , 2005 , 64 Suppl 1, i3-7	2.4	202
25	Can magnetic resonance imaging differentiate undifferentiated arthritis?. <i>Arthritis Research and Therapy</i> , 2005 , 7, 243-5	5.7	7

24	Routine database registration of biological therapy increases the reporting of adverse events twentyfold in clinical practice. First results from the Danish Database (DANBIO). <i>Scandinavian Journal of Rheumatology</i> , 2005 , 34, 40-4	1.9	38
23	Precision of bolus thermodilution cardiac output measurements in patients with atrial fibrillation. <i>Acta Anaesthesiologica Scandinavica</i> , 2005 , 49, 366-72	1.9	27
22	Imaging in early rheumatoid arthritis: roles of magnetic resonance imaging, ultrasonography, conventional radiography and computed tomography. <i>Best Practice and Research in Clinical Rheumatology</i> , 2005 , 19, 91-116	5.3	81
21	The smallest detectable difference and sensitivity to change of magnetic resonance imaging and radiographic scoring of structural joint damage in rheumatoid arthritis finger, wrist, and toe joints: a comparison of the OMERACT rheumatoid arthritis magnetic resonance imaging score applied to different joint combinations and the Sharp/van der Heide radiographic score. <i>Arthritis and Rheumatism</i> , 2005 , 52, 3860-7		99
20	Reliability and sensitivity to change of the OMERACT rheumatoid arthritis magnetic resonance imaging score in a multireader, longitudinal setting. <i>Arthritis and Rheumatism</i> , 2005 , 52, 3860-7		149
19	The EULAR-OMERACT rheumatoid arthritis MRI reference image atlas: the wrist joint. <i>Annals of the Rheumatic Diseases</i> , 2005 , 64 Suppl 1, i23-47	2.4	98
18	Magnetic resonance imaging for accelerated assessment of drug effect and prediction of subsequent radiographic progression in rheumatoid arthritis: a study of patients receiving combined anakinra and methotrexate treatment. <i>Annals of the Rheumatic Diseases</i> , 2005 , 64, 1503-6	2.4	29
17	Magnetic resonance imaging of peripheral joints in rheumatic diseases. <i>Best Practice and Research in Clinical Rheumatology</i> , 2004 , 18, 861-79	5.3	22
16	Magnetic resonance imaging of wrist and finger joints in healthy subjects occasionally shows changes resembling erosions and synovitis as seen in rheumatoid arthritis. <i>Arthritis and Rheumatism</i> , 2004 , 50, 1097-106		123
15	Ultrasonography in rheumatoid arthritis: a very promising method still needing more validation. <i>Current Opinion in Rheumatology</i> , 2004 , 16, 223-30	5.3	27
14	New radiographic bone erosions in the wrists of patients with rheumatoid arthritis are detectable with magnetic resonance imaging a median of two years earlier. <i>Arthritis and Rheumatism</i> , 2003 , 48, 2128-31		128
13	OMERACT Rheumatoid Arthritis Magnetic Resonance Imaging Studies. Core set of MRI acquisitions, joint pathology definitions, and the OMERACT RA-MRI scoring system. <i>Journal of Rheumatology</i> , 2003 , 30, 1385-6	4.1	424
12	Power Doppler ultrasonography for assessment of synovitis in the metacarpophalangeal joints of patients with rheumatoid arthritis: a comparison with dynamic magnetic resonance imaging. <i>Arthritis and Rheumatism</i> , 2001 , 44, 2018-23		320
11	Importance of timing of post-contrast MRI in rheumatoid arthritis: what happens during the first 60 minutes after IV gadolinium-DTPA?. <i>Annals of the Rheumatic Diseases</i> , 2001 , 60, 1050-4	2.4	60
10	Low field dedicated magnetic resonance imaging in untreated rheumatoid arthritis of recent onset. <i>Annals of the Rheumatic Diseases</i> , 2001 , 60, 770-6	2.4	67
9	Magnetic Resonance Imaging in Rheumatoid Arthritis. Quantitative methods for assessment of the inflammatory process in peripheral joints: Summary of Thesis. <i>Scandinavian Journal of Rheumatology</i> , 1999 , 28, 265-265	1.9	
8	Quantification of synovitis by MRI: correlation between dynamic and static gadolinium-enhanced magnetic resonance imaging and microscopic and macroscopic signs of synovial inflammation. <i>Magnetic Resonance Imaging</i> , 1998 , 16, 743-54	3.3	207
7	Intra-articular corticosteroids in arthritic disease: a guide to treatment. <i>BioDrugs</i> , 1998 , 9, 95-103	7.9	52

6	Magnetic resonance imaging-determined synovial membrane and joint effusion volumes in rheumatoid arthritis and osteoarthritis: comparison with the macroscopic and microscopic appearance of the synovium. <i>Arthritis and Rheumatism</i> , 1997 , 40, 1856-67		206
5	The accuracy of MRI-determined synovial membrane and joint effusion volumes in arthritis. A comparison of pre- and post-aspiration volumes. <i>Scandinavian Journal of Rheumatology</i> , 1995 , 24, 305-11 ^{1.9}		29
4	Effect of intraarticular osmic acid on synovial membrane volume and inflammation, determined by magnetic resonance imaging. <i>Scandinavian Journal of Rheumatology</i> , 1995 , 24, 5-12	1.9	25
3	Scoring of synovial membrane hypertrophy and bone erosions by MR imaging in clinically active and inactive rheumatoid arthritis of the wrist. <i>Scandinavian Journal of Rheumatology</i> , 1995 , 24, 212-8	1.9	81
2	Intra-articular and circulating levels of type I and III collagen markers in inflammatory and degenerative joint disease. <i>Acta Orthopaedica</i> , 1995 , 66, 147-148		
1	Synovial volume--a marker of disease severity in rheumatoid arthritis? Quantification by MRI. <i>Scandinavian Journal of Rheumatology</i> , 1994 , 23, 197-202	1.9	65