

# Gabriela R Lauretti

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8948429/publications.pdf>

Version: 2024-02-01

54  
papers

1,826  
citations

236833

25  
h-index

265120

42  
g-index

55  
all docs

55  
docs citations

55  
times ranked

1138  
citing authors

#	ARTICLE	IF	CITATIONS
1	Oral Ketamine and Transdermal Nitroglycerin as Analgesic Adjuvants to Oral Morphine Therapy for Cancer Pain Management. <i>Anesthesiology</i> , 1999, 90, 1528-1533..	1.3	202
2	Mecanismos envolvidos na analgesia da lidocaína por via venosa. <i>Revista Brasileira De Anestesiologia</i> , 2008, 58, 280-286.	0.6	114
3	Comparison of sustained-release morphine with sustained-release oxycodone in advanced cancer patients. <i>British Journal of Cancer</i> , 2003, 89, 2027-2030.	2.9	110
4	Study of Three Different Doses of Epidural Neostigmine Coadministered with Lidocaine for Postoperative Analgesia. <i>Anesthesiology</i> , 1999, 90, 1534-1538..	1.3	93
5	Dose-Response Study of Intrathecal Morphine Versus Intrathecal Neostigmine, Their Combination, or Placebo for Postoperative Analgesia in Patients Undergoing Anterior and Posterior Vaginoplasty. <i>Anesthesia and Analgesia</i> , 1996, 82, 1182-1187.	1.1	92
6	The activity of opioid analgesics in seizure models utilizing N-methyl-dl-aspartic acid, kainic acid, bicuculline and pentylentetrazole. <i>Neuropharmacology</i> , 1994, 33, 155-160.	2.0	77
7	Low doses of epidural ketamine or neostigmine, but not midazolam, improve morphine analgesia in epidural terminal cancer pain therapy. <i>Journal of Clinical Anesthesia</i> , 1999, 11, 663-668.	0.7	74
8	Dose-Response Study of Intrathecal Morphine Versus Intrathecal Neostigmine, Their Combination, or Placebo for Postoperative Analgesia in Patients Undergoing Anterior and Posterior Vaginoplasty. <i>Anesthesia and Analgesia</i> , 1996, 82, 1182-1187.	1.1	68
9	High and Low Frequency TENS Reduce Postoperative Pain Intensity After Laparoscopic Tubal Ligation. <i>Clinical Journal of Pain</i> , 2009, 25, 12-19.	0.8	66
10	Double-blind evaluation of transdermal nitroglycerine as adjuvant to oral morphine for cancer pain management. <i>Journal of Clinical Anesthesia</i> , 2002, 14, 83-86.	0.7	62
11	Epidural Morphine and Neostigmine for Postoperative Analgesia After Orthopedic Surgery. <i>Anesthesia and Analgesia</i> , 2002, 95, 1698-1701.	1.1	61
12	The Effects of Intrathecal Neostigmine on Somatic and Visceral Pain. <i>Anesthesia and Analgesia</i> , 1996, 82, 617-620.	1.1	60
13	Intrathecal neostigmine for postoperative analgesia after orthopedic surgery. <i>Journal of Clinical Anesthesia</i> , 1997, 9, 473-477.	0.7	52
14	Antinociceptive Effect of Low-Dose Intrathecal Neostigmine Combined with Intrathecal Morphine following Gynecologic Surgery. <i>Anesthesiology</i> , 2003, 98, 495-498.	1.3	51
15	Transdermal Ketamine as an Adjuvant for Postoperative Analgesia After Abdominal Gynecological Surgery Using Lidocaine Epidural Blockade. <i>Anesthesia and Analgesia</i> , 2000, 91, 1479-1482.	1.1	42
16	Efficacy of the Greater Occipital Nerve Block for Cervicogenic Headache: Comparing Classical and Subcompartmental Techniques. <i>Pain Practice</i> , 2015, 15, 654-661.	0.9	41
17	Postoperative analgesia by intraarticular and epidural neostigmine following knee surgery. <i>Journal of Clinical Anesthesia</i> , 2000, 12, 444-448.	0.7	39
18	Efficacy of the use of two simultaneously TENS devices for fibromyalgia pain. <i>Rheumatology International</i> , 2013, 33, 2117-2122.	1.5	35

#	ARTICLE	IF	CITATIONS
19	Intravenous Ketamine or Fentanyl Prolongs Postoperative Analgesia After Intrathecal Neostigmine. <i>Anesthesia and Analgesia</i> , 1996, 83, 766-770.	1.1	33
20	Enantioselective analysis of unbound tramadol, O-desmethyltramadol and N-desmethyltramadol in plasma by ultrafiltration and LC-MS/MS: Application to clinical pharmacokinetics. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2012, 880, 140-147.	1.2	32
21	The New Portable Transcutaneous Electrical Nerve Stimulation Device Was Efficacious in the Control of Primary Dysmenorrhea Cramp Pain. <i>Neuromodulation</i> , 2015, 18, 522-527.	0.4	32
22	Intravenous Ketamine or Fentanyl Prolongs Postoperative Analgesia After Intrathecal Neostigmine. <i>Anesthesia and Analgesia</i> , 1996, 83, 766-770.	1.1	27
23	Postoperative analgesia and antiemetic efficacy after subarachnoid neostigmine in orthopedic surgery. <i>Regional Anesthesia and Pain Medicine</i> , 1997, 22, 337-342.	1.1	27
24	Combined intrathecal fentanyl and neostigmine. <i>Journal of Clinical Anesthesia</i> , 1998, 10, 291-296.	0.7	27
25	The Effect of Transdermal Nitroglycerin on Spinal S(+)-Ketamine Antinociception Following Orthopedic Surgery. <i>Journal of Clinical Anesthesia</i> , 2001, 13, 576-581.	0.7	26
26	Epidural Nonsteroidal Antiinflammatory Drugs for Cancer Pain. <i>Anesthesia and Analgesia</i> , 1998, 86, 117-118.	1.1	25
27	Intrathecal ketorolac enhances intrathecal morphine analgesia following total knee arthroplasty. <i>Journal of Anaesthesiology Clinical Pharmacology</i> , 2013, 29, 503.	0.2	23
28	Low-dose intrathecal clonidine combined with sufentanil as analgesic drugs in abdominal gynecological surgery. <i>Journal of Clinical Anesthesia</i> , 2000, 12, 357-362.	0.7	21
29	The evolution of spinal/epidural neostigmine in clinical application: Thoughts after two decades. <i>Saudi Journal of Anaesthesia</i> , 2015, 9, 71.	0.2	21
30	Epidural methadone results in dose-dependent analgesia in cancer pain, further enhanced by epidural dexamethasone. <i>British Journal of Cancer</i> , 2013, 108, 259-264.	2.9	18
31	The Antinociceptive Effect of Local or Systemic Parecoxib Combined with Lidocaine/Clonidine Intravenous Regional Analgesia for Complex Regional Pain Syndrome Type I in the Arm. <i>Anesthesia and Analgesia</i> , 2005, 101, 807-811.	1.1	17
32	Tramadol and beta-cyclodextrin piroxicam: Effective multimodal balanced analgesia for the intra- and postoperative period. <i>Regional Anesthesia and Pain Medicine</i> , 1997, 22, 243-248.	1.1	16
33	The SLCO1A2 -189_-188InsA polymorphism reduces clearance of rocuronium in patients submitted to elective surgeries. <i>European Journal of Clinical Pharmacology</i> , 2017, 73, 957-963.	0.8	14
34	MODIFICATION BY DRUGS USED IN ANAESTHESIA OF CNS STIMULATION INDUCED IN MICE BY LAUDANOSINE AND STRYCHNINE. <i>British Journal of Anaesthesia</i> , 1991, 67, 608-613.	1.5	13
35	Pharmacogenomics in chronic pain therapy: from disease to treatment and challenges for clinical practice. <i>Pharmacogenomics</i> , 2019, 20, 971-982.	0.6	13
36	Propofol decreases ocular pressure in outpatients undergoing trabeculectomy. <i>Journal of Clinical Anesthesia</i> , 1997, 9, 289-292.	0.7	11

#	ARTICLE	IF	CITATIONS
37	Clonidine as coadjuvant in eye surgery: comparison of peribulbar versus oral administration. <i>Journal of Clinical Anesthesia</i> , 2002, 14, 140-145.	0.7	11
38	The combination of epidural clonidine and S(+)-ketamine did not enhance analgesic efficacy beyond that for each individual drug in adult orthopedic surgery. <i>Journal of Clinical Anesthesia</i> , 2005, 17, 79-84.	0.7	10
39	Pharmacokinetics and pharmacodynamics of rocuronium in young adult and elderly patients undergoing elective surgery. <i>Journal of Pharmacy and Pharmacology</i> , 2016, 68, 1351-1358.	1.2	10
40	Cetirizine Reduces Gabapentin Plasma Concentrations and Effect: Role of Renal Drug Transporters for Organic Cations. <i>Journal of Clinical Pharmacology</i> , 2020, 60, 1076-1086.	1.0	8
41	Spinal Alcohol Neurolysis for Intractable Thoracic Postherpetic Neuralgia after Test Bupivacaine Spinal Analgesia. <i>Anesthesiology</i> , 2004, 101, 244-247.	1.3	7
42	Analysis of rocuronium in human plasma by liquid chromatography-tandem mass spectrometry with application in clinical pharmacokinetics. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2014, 90, 180-185.	1.4	7
43	The Effects of Intrathecal Neostigmine on Somatic and Visceral Pain. <i>Anesthesia and Analgesia</i> , 1996, 82, 617-620.	1.1	6
44	Impact of fraction unbound, <sc>CYP</sc>3A, and <sc>CYP</sc>2D6 in vivo activities, and other potential covariates to the clearance of tramadol enantiomers in patients with neuropathic pain. <i>Fundamental and Clinical Pharmacology</i> , 2016, 30, 153-161.	1.0	5
45	Highlights in opioid agonists and antagonists. <i>Expert Review of Neurotherapeutics</i> , 2006, 6, 613-622.	1.4	4
46	Population pharmacokinetics of gabapentin in patients with neuropathic pain: Lack of effect of diabetes or glycaemic control. <i>British Journal of Clinical Pharmacology</i> , 2021, 87, 1981-1989.	1.1	4
47	AvaliaçŁo clĀnica comparativa entre a cetamina e a clonidina por via peridural no tratamento da dor crĀnica neuropĀtica. <i>Revista Brasileira De Anestesiologia</i> , 2002, 52, 34-40.	0.6	3
48	AvaliaçŁo duplamente encoberta da hipnose em fibromialgia. <i>Sao Paulo Medical Journal</i> , 2005, 123, 14-14.	0.4	3
49	AvaliaçŁo do efeito antinociceptivo do fentanil transdĀrmico no controle da dor lombar pĀs-operatĀria. <i>Coluna/ Columna</i> , 2009, 8, 412-416.	0.0	2
50	Analgesia after Epidural Dexamethasone is Further Enhanced by IV Dipyronne, but Not IV Parecoxibe Following Minor Orthopedic Surgery. <i>Korean Journal of Pain</i> , 2014, 27, 345-352.	0.8	2
51	Comparison between the intravenous and caudal routes of sufentanil in children undergoing orchidopexy and further evaluation of the association of caudal adrenaline and neostigmine. <i>Saudi Journal of Anaesthesia</i> , 2014, 8, 345.	0.2	1
52	The clinical and laboratorial evaluation of transdermal ketamine, fentanyl, clonidine or their combination in chronic low back pain. <i>Coluna/ Columna</i> , 2009, 8, 434-440.	0.0	1
53	Chronic visceral pain secondary to ventral disc herniation: Development of visceral complex regional pain syndrome. <i>Saudi Journal of Anaesthesia</i> , 2015, 9, 314.	0.2	1
54	Spinal and intravenous midazolam anesthetic effects on fentanyl/ lidocaine regional anesthesia following back minor orthopedic surgery. <i>Coluna/ Columna</i> , 2010, 9, 30-34.	0.0	0