

Manel Mata-Cases

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/8949199/manel-mata-cases-publications-by-year.pdf>
Version: 2024-04-11

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.
The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

71 papers	1,301 citations	19 h-index	34 g-index
93 ext. papers	1,786 ext. citations	4.1 avg, IF	4.33 L-index

#	Paper	IF	Citations
71	Compliance with the DPP-4 inhibitors dose adjustment recommendations based on renal function in a population database.. <i>Endocrinología Diabetes Y Nutrición (English Ed)</i> , 2022 , 69, 83-91	0.1	
70	Mortality trends in type 1 diabetes: a multicountry analysis of six population-based cohorts.. <i>Diabetologia</i> , 2022 , 1	10.3	4
69	Heterogeneity in the association between prediabetes categories and reduction on glomerular filtration rate in a 5-year follow-up.. <i>Scientific Reports</i> , 2022 , 12, 7373	4.9	0
68	Trends in the Degree of Control and Treatment of Cardiovascular Risk Factors in People With Type 2 Diabetes in a Primary Care Setting in Catalonia During 2007-2018.. <i>Frontiers in Endocrinology</i> , 2021 , 12, 810757	5.7	1
67	Trends in the incidence of diagnosed diabetes: a multicountry analysis of aggregate data from 22 million diagnoses in high-income and middle-income settings. <i>Lancet Diabetes and Endocrinology</i> , 2021 , 9, 203-211	18.1	20
66	The heterogeneity of reversion to normoglycemia according to prediabetes type is not explained by lifestyle factors. <i>Scientific Reports</i> , 2021 , 11, 9667	4.9	0
65	327-OR: Multicountry Analysis of Trends in All-Cause Mortality among People with Type 1 Diabetes. <i>Diabetes</i> , 2021 , 70, 327-OR	0.9	
64	Compliance with the DPP-4 inhibitors dose adjustment recommendations based on renal function in a population database. <i>Endocrinología, Diabetes Y Nutrición</i> , 2021 , 69, 83-83	1.3	
63	Evaluation of clinical and antidiabetic treatment characteristics of different sub-groups of patients with type 2 diabetes: Data from a Mediterranean population database. <i>Primary Care Diabetes</i> , 2021 , 15, 588-595	2.4	4
62	Risk factors for severe outcomes in people with diabetes hospitalised for COVID-19: a cross-sectional database study. <i>BMJ Open</i> , 2021 , 11, e051237	3	0
61	Analysis of the Adherence and Safety of Second Oral Glucose-Lowering Therapy in Routine Practice From the Mediterranean Area: A Retrospective Cohort Study. <i>Frontiers in Endocrinology</i> , 2021 , 12, 708372	5.7	2
60	A disease state approach to the pharmacological management of Type 2 diabetes in primary care: A position statement by Primary Care Diabetes Europe. <i>Primary Care Diabetes</i> , 2021 , 15, 31-51	2.4	13
59	Analysis of the effectiveness of second oral glucose-lowering therapy in routine clinical practice from the mediterranean area: A retrospective cohort study. <i>Diabetes Research and Clinical Practice</i> , 2021 , 171, 108616	7.4	1
58	Cardiovascular and mortality benefits of sodium-glucose co-transporter-2 inhibitors in patients with type 2 diabetes mellitus: CVD-Real Catalonia. <i>Cardiovascular Diabetology</i> , 2021 , 20, 139	8.7	2
57	Incidence and characteristics of diabetic foot ulcers in subjects with type 2 diabetes in Catalanian primary care centres: An observational multicentre study. <i>Primary Care Diabetes</i> , 2021 , 15, 1033-1039	2.4	0
56	Mediterranean diet and diabetes risk in a cohort study of individuals with prediabetes: propensity score analyses.. <i>Diabetic Medicine</i> , 2021 , e14768	3.5	0
55	Prevalence of pre-diabetes and undiagnosed diabetes in the Mollerussa prospective observational cohort study in a semi-rural area of Catalonia. <i>BMJ Open</i> , 2020 , 10, e033332	3	5

54	The Association Between Poor Glycemic Control and Health Care Costs in People With Diabetes: A Population-Based Study. <i>Diabetes Care</i> , 2020 , 43, 751-758	14.6	18
53	Impact on guidelines: The general practitioner point of view. <i>Diabetes Research and Clinical Practice</i> , 2020 , 166, 108091	7.4	3
52	Half of patients with type 2 diabetes mellitus are at very high cardiovascular risk according to the ESC/EASD: data from a large Mediterranean population. <i>European Journal of Preventive Cardiology</i> , 2020 ,	3.9	5
51	Therapeutic Inertia: Still a Long Way to Go That Cannot Be Postponed. <i>Diabetes Spectrum</i> , 2020 , 33, 50-57.	5.9	2
50	miR-10b and miR-223-3p in serum microvesicles signal progression from prediabetes to type 2 diabetes. <i>Journal of Endocrinological Investigation</i> , 2020 , 43, 451-459	5.2	21
49	Prediabetes Is Independently Associated with Subclinical Carotid Atherosclerosis: An Observational Study in a Non-Urban Mediterranean Population. <i>Journal of Clinical Medicine</i> , 2020 , 9,	5.1	4
48	Improving management of glycaemic control in people with T2DM in primary care: estimation of the impact on the clinical complications and associated costs. <i>BMC Health Services Research</i> , 2020 , 20, 803	2.9	3
47	Glucagon-Like Peptide-1 Receptor Agonists in Patients with Type 2 Diabetes: Prescription According to Reimbursement Constraints and Guideline Recommendations in Catalonia. <i>Journal of Clinical Medicine</i> , 2019 , 8,	5.1	4
46	High levels of fasting glucose and glycosylated hemoglobin values are associated with hyperfiltration in a Spanish prediabetes cohort. The PREDAPS Study. <i>PLoS ONE</i> , 2019 , 14, e0222848	3.7	3
45	Lactic acidosis associated with metformin in patients with moderate to severe chronic kidney disease: study protocol for a multicenter population-based case-control study using health databases. <i>BMC Nephrology</i> , 2019 , 20, 193	2.7	1
44	Glucagon-like peptide-1 receptor agonists in patients with type 2 diabetes:real-world evidence from a Mediterranean area. <i>Current Medical Research and Opinion</i> , 2019 , 35, 1735-1744	2.5	7
43	Glycaemic control after treatment intensification in patients with type 2 diabetes uncontrolled on two or more non-insulin antidiabetic drugs in a real-world setting. <i>Diabetes, Obesity and Metabolism</i> , 2019 , 21, 1373-1380	6.7	9
42	INTEGRA study protocol: primary care intervention in type 2 diabetes patients with poor glycaemic control. <i>BMC Family Practice</i> , 2019 , 20, 25	2.6	0
41	Prevalence and coprevalence of chronic comorbid conditions in patients with type 2 diabetes in Catalonia: a population-based cross-sectional study. <i>BMJ Open</i> , 2019 , 9, e031281	3	22
40	How Many Patients with Type 2 Diabetes Meet the Inclusion Criteria of the Cardiovascular Outcome Trials with SGLT2 Inhibitors? Estimations from a Population Database in a Mediterranean Area. <i>Journal of Diabetes Research</i> , 2019 , 2019, 2018374	3.9	9
39	Consensus on insulin treatment in type 2 diabetes. <i>Endocrinología y Nutrición (English Ed)</i> , 2018 , 65, 1-8	0.1	
38	Decreased iron stores are associated with cardiovascular disease in patients with type 2 diabetes both cross-sectionally and longitudinally. <i>Atherosclerosis</i> , 2018 , 272, 193-199	3.1	6
37	Therapeutic inertia in patients treated with two or more antidiabetics in primary care: Factors predicting intensification of treatment. <i>Diabetes, Obesity and Metabolism</i> , 2018 , 20, 103-112	6.7	38

36	Fatty liver index is a predictor of incident diabetes in patients with prediabetes: The PREDAPS study. <i>PLoS ONE</i> , 2018 , 13, e0198327	3.7	23
35	Diabetes Treatment in Patients with Renal Disease: How Many Are Receiving Contraindicated Drugs?. <i>Diabetes</i> , 2018 , 67, 1647-P	0.9	
34	Multiple Chronic Comorbidities in a T2DM Mediterranean Population. <i>Diabetes</i> , 2018 , 67, 1646-P	0.9	
33	Spanish Diabetes Society (SED) recommendations for the pharmacologic treatment of hyperglycemia in type 2 diabetes: 2018 Update. <i>Endocrinología, Diabetes Y Nutrición</i> , 2018 , 65, 611-624	1.3	19
32	Prevalence, clinical features and risk assessment of pre-diabetes in Spain: the prospective Mollerussa cohort study. <i>BMJ Open</i> , 2017 , 7, e015158	3	14
31	Impact of implementing electronic clinical practice guidelines for the diagnosis, control and treatment of cardiovascular risk factors: A pre-post controlled study. <i>Atencion Primaria</i> , 2017 , 49, 389-398	3.6	4
30	Is quality of life different between diabetic and non-diabetic people? The importance of cardiovascular risks. <i>PLoS ONE</i> , 2017 , 12, e0189505	3.7	9
29	Prevalence and control of type 2 diabetes mellitus among primary care physicians in Spain. PRISMA Study. <i>Endocrinología, Diabetes Y Nutrición</i> , 2017 , 64, 265-271	1.3	2
28	Clinical characteristics of type 2 diabetic patients on basal insulin therapy with adequate fasting glucose control who do not achieve HbA1c targets. <i>Journal of Diabetes</i> , 2017 , 9, 34-44	3.8	14
27	Attitudes towards insulin initiation in type 2 diabetes patients among healthcare providers: A survey research. <i>Diabetes Research and Clinical Practice</i> , 2016 , 122, 46-53	7.4	13
26	Glycaemic control and antidiabetic treatment trends in primary care centres in patients with type 2 diabetes mellitus during 2007-2013 in Catalonia: a population-based study. <i>BMJ Open</i> , 2016 , 6, e012463	3	49
25	Epidemiology and clinical management of type 2 diabetes mellitus and associated comorbidities in Spain (e-Management study). <i>Medicina Clínica</i> , 2016 , 147 Suppl 1, 1-7	1	8
24	Direct medical costs attributable to type 2 diabetes mellitus: a population-based study in Catalonia, Spain. <i>European Journal of Health Economics</i> , 2016 , 17, 1001-1010	3.6	58
23	Understanding the physical, social, and emotional experiences of people with uncontrolled Type 2 diabetes: a qualitative study. <i>Patient Preference and Adherence</i> , 2016 , 10, 2323-2332	2.4	9
22	Noninsulin Antidiabetic Drugs for Patients with Type 2 Diabetes Mellitus: Are We Respecting Their Contraindications?. <i>Journal of Diabetes Research</i> , 2016 , 2016, 7502489	3.9	13
21	Is diabetes mellitus correctly registered and classified in primary care? A population-based study in Catalonia, Spain. <i>Endocrinología Y Nutrición: Organo De La Sociedad Espanola De Endocrinología Y Nutrición</i> , 2016 , 63, 440-448		12
20	Risk of Cause-Specific Death in Individuals With Diabetes: A Competing Risks Analysis. <i>Diabetes Care</i> , 2016 , 39, 1987-1995	14.6	165
19	The impact of a programme to improve quality of care for people with type 2 diabetes on hard to reach groups: The GEDAPS study. <i>Primary Care Diabetes</i> , 2015 , 9, 211-8	2.4	12

18	Consenso sobre la detección y el manejo de la prediabetes. Grupo de Trabajo de Consensos y Guías Clínicas de la Sociedad Española de Diabetes. <i>Avances En Diabetología</i> , 2015 , 31, 89-101		3
17	Metabolic control and cardiovascular risk factors in type 2 diabetes mellitus patients according to diabetes duration. <i>Family Practice</i> , 2015 , 32, 27-34	1.9	15
16	Older type 2 diabetic patients are more likely to achieve glycaemic and cardiovascular risk factors targets than younger patients: analysis of a primary care database. <i>International Journal of Clinical Practice</i> , 2015 , 69, 1486-95	2.9	38
15	Prevalence of diabetic retinopathy in individuals with type 2 diabetes who had recorded diabetic retinopathy from retinal photographs in Catalonia (Spain). <i>British Journal of Ophthalmology</i> , 2015 , 99, 1628-33	5.5	25
14	Consensus on the detection and management of prediabetes. Consensus and Clinical Guidelines Working Group of the Spanish Diabetes Society. <i>Revista Clinica Espanola</i> , 2015 , 215, 117-29	0.7	8
13	Differences in the Cardiometabolic Control in Type 2 Diabetes according to Gender and the Presence of Cardiovascular Disease: Results from the eControl Study. <i>International Journal of Endocrinology</i> , 2014 , 2014, 131709	2.7	22
12	Investigar en diabetes desde una base de datos de atención primaria: la experiencia del Sistema de Información para el Desarrollo de la Investigación en Atención Primaria (SIDIAP). <i>Avances En Diabetología</i> , 2013 , 29, 169-174		5
11	Clinical inertia in the treatment of hyperglycemia in type 2 diabetes patients in primary care. <i>Current Medical Research and Opinion</i> , 2013 , 29, 1495-502	2.5	42
10	Estudio observacional de eficacia y seguridad del cambio de insulina NPH a glargina en atención primaria. Estudio LAURA. <i>Avances En Diabetología</i> , 2013 , 29, 137-144		4
9	Inercia terapéutica en el control glucémico de la diabetes mellitus tipo 2: inconvenientes y ventajas. <i>Hipertension Y Riesgo Vascular</i> , 2012 , 29, 34-40	0.5	1
8	Chronic kidney disease in the type 2 diabetic patients: prevalence and associated variables in a random sample of 2642 patients of a Mediterranean area. <i>BMC Nephrology</i> , 2012 , 13, 87	2.7	39
7	Trends in chronic complications of type 2 diabetic patients from Spanish primary health care centres (GEDAPS study): ten year-implementation of St. Vincent recommendations. <i>Primary Care Diabetes</i> , 2012 , 6, 11-8	2.4	20
6	Fifteen years of continuous improvement of quality care of type 2 diabetes mellitus in primary care in Catalonia, Spain. <i>International Journal of Clinical Practice</i> , 2012 , 66, 289-98	2.9	47
5	Control of glycemia and cardiovascular risk factors in patients with type 2 diabetes in primary care in Catalonia (Spain). <i>Diabetes Care</i> , 2012 , 35, 774-9	14.6	151
4	Incidence of complications and mortality in a type 2 diabetes patient cohort study followed up from diagnosis in a primary healthcare centre. <i>International Journal of Clinical Practice</i> , 2011 , 65, 299-307	2.9	18
3	La aplicación de la HbA1c como único criterio podrá retrasar el diagnóstico de la diabetes mellitus tipo 2. <i>Avances En Diabetología</i> , 2010 , 26, 419-423		1
2	Evaluation of risk factors for development of complications in Type II diabetes in Europe. <i>Diabetologia</i> , 2002 , 45, S23-8	10.3	155
1	Criterios y pautas de terapia combinada en la diabetes tipo 2. <i>Endocrinología Y Nutrición: Organo De La Sociedad Espanola De Endocrinología Y Nutrición</i> , 2001 , 48, 82-97		5

