

# Muhammad Safder

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8951517/publications.pdf>

Version: 2024-02-01

15  
papers

121  
citations

1307594

7  
h-index

1281871

11  
g-index

16  
all docs

16  
docs citations

16  
times ranked

189  
citing authors

#	ARTICLE	IF	CITATIONS
1	The Metabolites of the Genus <i>Onopordum</i> (Asteraceae): Chemistry and Biological Properties. <i>Current Organic Chemistry</i> , 2011, 15, 888-927.	1.6	24
2	Artalbic acid, a sesquiterpene with an unusual skeleton from <i>Artemisia alba</i> (Asteraceae) from Sicily. <i>Tetrahedron Letters</i> , 2011, 52, 4543-4545.	1.4	18
3	Cytotoxic geranylflavonoids from <i>Bonannia graeca</i> . <i>Phytochemistry</i> , 2011, 72, 942-945.	2.9	14
4	Extraction, optimization, and characterization of lipids from spent hens: An unexploited sustainable bioresource. <i>Journal of Cleaner Production</i> , 2019, 206, 622-630.	9.3	13
5	Asphorodin, a potent lipoxygenase inhibitory triterpene diglycoside from <i>Asphodelus tenuifolius</i> . <i>Journal of Asian Natural Products Research</i> , 2009, 11, 945-950.	1.4	11
6	New Secondary Metabolites from <i>Asphodelus tenuifolius</i> . <i>Helvetica Chimica Acta</i> , 2012, 95, 144-151.	1.6	8
7	Eremosides A, New Iridoid Glucosides from <i>Eremostachys loasifolia</i> . <i>Helvetica Chimica Acta</i> , 2012, 95, 586-593.	1.6	7
8	Lipid-derived hybrid bionanocomposites from spent hens. <i>Materials Today Communications</i> , 2020, 25, 101327.	1.9	7
9	Bio-composites from spent hen derived lipids grafted on CNC and reinforced with nanoclay. <i>Carbohydrate Polymers</i> , 2022, 281, 119082.	10.2	7
10	Spiraeamide, new sphingolipid from <i>Spiraea brahuica</i> . <i>Journal of Asian Natural Products Research</i> , 2012, 14, 601-606.	1.4	5
11	Supercritical CO <sub>2</sub> extraction and solvent-free rapid alternative bioepoxy production from spent hens. <i>Journal of CO<sub>2</sub> Utilization</i> , 2019, 34, 335-342.	6.8	5
12	Structural determination of crotamides A and B, the new amides from <i>Croton sparsiflorus</i> . <i>Journal of Asian Natural Products Research</i> , 2010, 12, 662-665.	1.4	2
13	6-Bromo-4-(2-cyclohexylidenehydrazin-1-ylidene)-1-methyl-2,2-dioxo-3,4-dihydro-1 <i>H</i> -2,6,1-benzothiazine. <i>Acta Crystallographica Section E: Structure Reports Online</i> , 2012, 68, o2088-o2089.	0.2	0
14	{2-[(3-Bromobenzylidene)amino]-5-chlorophenyl}(phenyl)methanone. <i>Acta Crystallographica Section E: Structure Reports Online</i> , 2012, 68, o645-o645.	0.2	0
15	1-Ethyl-4-{2-[1-(4-methylphenyl)ethylidene]hydrazinylidene}-3,4-dihydro-1 <i>H</i> -2,6,1-benzothiazine-2,2-dione. <i>Acta Crystallographica Section E: Structure Reports Online</i> , 2013, 69, o200-o200.	0.2	0