

# Stuart J Pearse

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8970164/publications.pdf>

Version: 2024-02-01

20  
papers

2,697  
citations

706676

14  
h-index

889612

19  
g-index

20  
all docs

20  
docs citations

20  
times ranked

3701  
citing authors

#	ARTICLE	IF	CITATIONS
1	Interactions among cluster-root investment, leaf phosphorus concentration, and relative growth rate in two <i>Lupinus</i> species. <i>American Journal of Botany</i> , 2015, 102, 1529-1537.	0.8	5
2	Mineral nutrition of <i>campos rupestres</i> plant species on contrasting nutrient-impoverished soil types. <i>New Phytologist</i> , 2015, 205, 1183-1194.	3.5	149
3	Low levels of ribosomal rRNA partly account for the very high photosynthetic phosphorus-use efficiency of Proteaceae species. <i>Plant, Cell and Environment</i> , 2014, 37, 1276-1298.	2.8	121
4	Nutrient limitation along the Jurien Bay dune chronosequence: response to Uren & Parsons (). <i>Journal of Ecology</i> , 2013, 101, 1088-1092.	1.9	14
5	Cluster-root formation and carboxylate release in three <i>Lupinus</i> species as dependent on phosphorus supply, internal phosphorus concentration and relative growth rate. <i>Annals of Botany</i> , 2013, 112, 1449-1459.	1.4	18
6	Phosphorus nutrition of phosphorus-sensitive Australian native plants: threats to plant communities in a global biodiversity hotspot. <i>Journal of Ecology</i> , 2013, 101, 1088-1092.		76
7	Downregulation of net phosphorus-uptake capacity is inversely related to leaf phosphorus-resorption proficiency in four species from a phosphorus-impoverished environment. <i>Annals of Botany</i> , 2013, 111, 445-454.	1.4	67
8	<i>Viminaria juncea</i> does not vary its shoot phosphorus concentration and only marginally decreases its mycorrhizal colonization and cluster-root dry weight under a wide range of phosphorus supplies. <i>Annals of Botany</i> , 2013, 111, 801-809.	1.4	13
9	Proteaceae from severely phosphorus-impoverished soils extensively replace phospholipids with galactolipids and sulfolipids during leaf development to achieve a high photosynthetic phosphorus-use efficiency. <i>New Phytologist</i> , 2012, 196, 1098-1108.	3.5	225
10	Experimental assessment of nutrient limitation along a 2-million-year dune chronosequence in the south-western Australia biodiversity hotspot. <i>Journal of Ecology</i> , 2012, 100, 631-642.	1.9	189
11	Why does the musketeer approach to phosphorus acquisition from sparingly soluble forms fail: All for one, but not one for all?. <i>Plant and Soil</i> , 2011, 348, 81-83.	1.8	4
12	An enzymatic fluorescent assay for the quantification of phosphite in a microtiter plate format. <i>Analytical Biochemistry</i> , 2011, 412, 74-78.	1.1	14
13	Phosphorus Nutrition of Proteaceae in Severely Phosphorus-Impoverished Soils: Are There Lessons To Be Learned for Future Crops?. <i>Plant Physiology</i> , 2011, 156, 1058-1066.	2.3	176
14	Biological nitrification inhibition by <i>Brachiaria humidicola</i> roots varies with soil type and inhibits nitrifying bacteria, but not other major soil microorganisms. <i>Soil Science and Plant Nutrition</i> , 2009, 55, 725-733.	0.8	47
15	Detection, isolation and characterization of a root-exuded compound, methyl 3-(4-hydroxyphenyl) propionate, responsible for biological nitrification inhibition by sorghum ( <i>Sorghum bicolor</i> ). <i>New Phytologist</i> , 2008, 180, 442-451.	3.5	148
16	Rhizosphere processes do not explain variation in P acquisition from sparingly soluble forms among <i>Lupinus albus</i> accessions. <i>Australian Journal of Agricultural Research</i> , 2008, 59, 616.	1.5	8
17	Carboxylate composition of root exudates does not relate consistently to a crop species'™ ability to use phosphorus from aluminium, iron or calcium phosphate sources. <i>New Phytologist</i> , 2007, 173, 181-190.	3.5	175
18	<i>Triticum aestivum</i> shows a greater biomass response to a supply of aluminium phosphate than <i>Lupinus albus</i> , despite releasing fewer carboxylates into the rhizosphere. <i>New Phytologist</i> , 2006, 169, 515-524.	3.5	67

#	ARTICLE	IF	CITATIONS
19	Carboxylate release of wheat, canola and 11 grain legume species as affected by phosphorus status. <i>Plant and Soil</i> , 2006, 288, 127-139.	1.8	169
20	Root Structure and Functioning for Efficient Acquisition of Phosphorus: Matching Morphological and Physiological Traits. <i>Annals of Botany</i> , 2006, 98, 693-713.	1.4	1,012