
Elżbieta A WÅ,odarczyk

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/8970410/publications.pdf

Version: 2024-02-01



#	Article	IF	CITATIONS
1	Development of central auditory processes in Polish children and adolescents at the age from 7 to 16Âyears. Current Psychology, 2023, 42, 1789-1806.	1.7	1
2	The COVID-19 pandemic and upgrades of CI speech processors for children: part l—procedure of speech processor upgrade. European Archives of Oto-Rhino-Laryngology, 2022, , 1.	0.8	0
3	The COVID-19 pandemic and upgrades of CI speech processors for children: part Il–hearing outcomes. European Archives of Oto-Rhino-Laryngology, 2022, , 1.	0.8	2
4	A simple qualitative scale for diagnosis of laryngopharyngeal reflux: high correlations with pH measurements and disease severity. The usefulness of the Warsaw Scale in LPR diagnostics compared to other diagnostic tools. European Archives of Oto-Rhino-Laryngology, 2021, 278, 4883-4892.	0.8	1
5	A revised grading system for the Tinnitus Handicap Inventory based on a large clinical population. International Journal of Audiology, 2020, 59, 61-67.	0.9	24
6	Audiological and psychological profiles of children with tinnitus. Hearing, Balance and Communication, 2020, 18, 90-97.	0.1	5
7	Three-year audiological outcomes of the latest generation middle ear transducer (MET) implant. European Archives of Oto-Rhino-Laryngology, 2020, 277, 3013-3019.	0.8	1
8	Diagnosis of laryngopharyngeal reflux in children with voice disorders using 24-hour pharyngeal pH monitoring. International Journal of Pediatric Otorhinolaryngology, 2019, 121, 188-196.	0.4	12
9	The application of 24-hour pharyngeal pH-monitoring and Reflux Finding Score and Reflux Symptom Index questionnaires in the diagnostics of laryngopharyngeal reflux. Przeglad Gastroenterologiczny, 2019, 14, 274-282.	0.3	5
10	Visual Analogue Scales as a Tool for Initial Assessment of Tinnitus Severity: Psychometric Evaluation in a Clinical Population. Audiology and Neuro-Otology, 2018, 23, 229-237.	0.6	27
11	The use of RSI and RFS questionnaires in the Polish language version. Otolaryngologia Polska, 2018, 72, 1-5.	0.2	4