
Hans-Christian Pape

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/897173/publications.pdf Version: 2024-02-01



#	Article	lF	CITATIONS
1	Functional deletion of neuropeptide Y receptors type 2 in local synaptic networks of anteroventral BNST facilitates recall and increases return of fear. Molecular Psychiatry, 2021, 26, 2900-2911.	7.9	6
2	Brain-Derived Neurotrophic Factor/Tropomyosin Receptor Kinase B Signaling Controls Excitability and Long-Term Depression in Oval Nucleus of the BNST. Journal of Neuroscience, 2021, 41, 435-445.	3.6	8
3	Neuropeptide S Receptor Stimulation Excites Principal Neurons in Murine Basolateral Amygdala through a Calcium-Dependent Decrease in Membrane Potassium Conductance. Pharmaceuticals, 2021, 14, 519.	3.8	1
4	Human-Specific Neuropeptide S Receptor Variants Regulate Fear Extinction in the Basal Amygdala of Male and Female Mice Depending on Threat Salience. Biological Psychiatry, 2021, 90, 145-155.	1.3	10
5	Seizure prediction in genetic rat models of absence epilepsy: improved performance through multiple-site cortico-thalamic recordings combined with machine learning. ENeuro, 2021, , ENEURO.0160-21.2021.	1.9	0
6	Stimulation of 5-HT receptors in anterodorsal BNST guides fear to predictable and unpredictable threat. European Neuropsychopharmacology, 2020, 39, 56-69.	0.7	4
7	The µ-opioid system in midline thalamic nuclei modulates defence strategies towards a conditioned fear stimulus in male mice. Journal of Psychopharmacology, 2020, 34, 1280-1288.	4.0	10
8	Spikeâ€wave discharges in absence epilepsy: segregation of electrographic components reveals distinct pathways of seizure activity. Journal of Physiology, 2020, 598, 2397-2414.	2.9	25
9	On the objectivity, reliability, and validity of deep learning enabled bioimage analyses. ELife, 2020, 9, .	6.0	24
10	Regional specificity of cortico-thalamic coupling strength and directionality during waxing and waning of spike and wave discharges. Scientific Reports, 2019, 9, 2100.	3.3	23
11	Single stimulation of Y2 receptors in BNSTav facilitates extinction and dampens reinstatement of fear. Psychopharmacology, 2019, 236, 281-291.	3.1	9
12	Neuropeptide Y2 receptors in anteroventral BNST control remote fear memory depending on extinction training. Neurobiology of Learning and Memory, 2018, 149, 144-153.	1.9	14
13	24 Integrative Funktionen des Gehirns. , 2018, , .		0
14	Physiological Profile of Neuropeptide Y-Expressing Neurons in Bed Nucleus of Stria Terminalis in Mice: State of High Excitability. Frontiers in Cellular Neuroscience, 2018, 12, 393.	3.7	10
15	Impairment of frequency-specific responses associated with altered electrical activity patterns in auditory thalamus following focal and general demyelination. Experimental Neurology, 2018, 309, 54-66.	4.1	15
16	Neuropeptide S precursor knockout mice display memory and arousal deficits. European Journal of Neuroscience, 2017, 46, 1689-1700.	2.6	23
17	The quality of cortical network function recovery depends on localization and degree of axonal demyelination. Brain, Behavior, and Immunity, 2017, 59, 103-117.	4.1	25
18	Impact of Life History on Fear Memory and Extinction. Frontiers in Behavioral Neuroscience, 2016, 10, 185.	2.0	11

#	Article	IF	CITATIONS
19	Dynorphin-Dependent Reduction of Excitability and Attenuation of Inhibitory Afferents of NPS Neurons in the Pericoerulear Region of Mice. Frontiers in Cellular Neuroscience, 2016, 10, 61.	3.7	4
20	Neuronal correlates of sustained fear in the anterolateral part of the bed nucleus of stria terminalis. Neurobiology of Learning and Memory, 2016, 131, 137-146.	1.9	32
21	Distinct state anxiety after predictable and unpredictable fear training in mice. Behavioural Brain Research, 2016, 304, 20-23.	2.2	16
22	mGluR-mediated calcium signalling in the thalamic reticular nucleus. Cell Calcium, 2016, 59, 312-323.	2.4	13
23	Shortâ€ŧerm depression of gap junctional coupling in reticular thalamic neurons of absence epileptic rats. Journal of Physiology, 2016, 594, 5695-5710.	2.9	18
24	Neuronal Expression of the Human Neuropeptide S Receptor NPSR1 Identifies NPS-Induced Calcium Signaling Pathways. PLoS ONE, 2015, 10, e0117319.	2.5	25
25	Â-Opioid Receptor-Mediated Inhibition of Intercalated Neurons and Effect on Synaptic Transmission to the Central Amygdala. Journal of Neuroscience, 2015, 35, 7317-7325.	3.6	43
26	Increased GABAergic Efficacy of Central Amygdala Projections to Neuropeptide S Neurons in the Brainstem During Fear Memory Retrieval. Neuropsychopharmacology, 2015, 40, 2753-2763.	5.4	24
27	Expression of freezing and fearâ€potentiated startle during sustained fear in mice. Genes, Brain and Behavior, 2015, 14, 281-291.	2.2	45
28	Glutamic Acid Decarboxylase 65: A Link Between GABAergic Synaptic Plasticity in the Lateral Amygdala and Conditioned Fear Generalization. Neuropsychopharmacology, 2014, 39, 2211-2220.	5.4	51
29	Spreading depression triggers ictaform activity in partially disinhibited neuronal tissues. Experimental Neurology, 2014, 253, 1-15.	4.1	46
30	Cannabinoid CB1 Receptor in Dorsal Telencephalic Glutamatergic Neurons: Distinctive Sufficiency for Hippocampus-Dependent and Amygdala-Dependent Synaptic and Behavioral Functions. Journal of Neuroscience, 2013, 33, 10264-10277.	3.6	108
31	Neuropeptide S receptor gene is associated with cortisol responses to social stress in humans. Biological Psychology, 2013, 93, 304-307.	2.2	45
32	Single dose of <scp>l</scp> -dopa makes extinction memories context-independent and prevents the return of fear. Proceedings of the National Academy of Sciences of the United States of America, 2013, 110, E2428-36.	7.1	169
33	Neuronale Schaltkreise von FurchtgedÃehtnis und Furchtextinktion. E-Neuroforum, 2013, 19, 92-103.	0.1	5
34	Directional Theta Coherence in Prefrontal Cortical to Amygdalo-Hippocampal Pathways Signals Fear Extinction. PLoS ONE, 2013, 8, e77707.	2.5	86
35	Differential regulation of glutamic acid decarboxylase gene expression after extinction of a recent memory vs. intermediate memory. Learning and Memory, 2012, 19, 194-200.	1.3	17
36	Activation of neuropeptide Sâ€expressing neurons in the locus coeruleus by corticotropinâ€releasing factor. Journal of Physiology, 2012, 590, 3701-3717.	2.9	38

#	Article	IF	CITATIONS
37	Prevention of Stress-Impaired Fear Extinction Through Neuropeptide S Action in the Lateral Amygdala. Neuropsychopharmacology, 2012, 37, 1588-1599.	5.4	88
38	A special issue on sleep. Pflugers Archiv European Journal of Physiology, 2012, 463, 1-2.	2.8	3
39	Impaired extinction of fear and maintained amygdalaâ€hippocampal theta synchrony in a mouse model of temporal lobe epilepsy. Epilepsia, 2011, 52, 337-346.	5.1	34
40	Short-Term Adaptation of Conditioned Fear Responses Through Endocannabinoid Signaling in the Central Amygdala. Neuropsychopharmacology, 2011, 36, 652-663.	5.4	84
41	Abnormalities in GABAergic synaptic transmission of intralaminar thalamic neurons in a genetic rat model of absence epilepsy. Molecular and Cellular Neurosciences, 2011, 46, 444-451.	2.2	10
42	Increased in vivo release of neuropeptide S in the amygdala of freely moving rats after local depolarisation and emotional stress. Amino Acids, 2011, 41, 991-996.	2.7	46
43	Seizure-related activity of intralaminar thalamic neurons in a genetic model of absence epilepsy. Neurobiology of Disease, 2011, 43, 266-274.	4.4	43
44	Neuropeptide S Enhances Memory During the Consolidation Phase and Interacts with Noradrenergic Systems in the Brain. Neuropsychopharmacology, 2011, 36, 744-752.	5.4	105
45	Patterns of Coupled Theta Activity in Amygdala-Hippocampal-Prefrontal Cortical Circuits during Fear Extinction. PLoS ONE, 2011, 6, e21714.	2.5	220
46	Social Defeat: Impact on Fear Extinction and Amygdala-Prefrontal Cortical Theta Synchrony in 5-HTT Deficient Mice. PLoS ONE, 2011, 6, e22600.	2.5	97
47	Alteration of NMDA receptor–mediated synaptic interactions in the lateral amygdala associated with seizure activity in a mouse model of chronic temporal lobe epilepsy. Epilepsia, 2010, 51, 1754-1762.	5.1	8
48	GABAergic Interneurons in the Mouse Lateral Amygdala: A Classification Study. Journal of Neurophysiology, 2010, 104, 617-626.	1.8	69
49	Petrified or Aroused with Fear: The Central Amygdala Takes the Lead. Neuron, 2010, 67, 527-529.	8.1	11
50	Neuropeptide S: A transmitter system in the brain regulating fear and anxiety. Neuropharmacology, 2010, 58, 29-34.	4.1	125
51	Plastic Synaptic Networks of the Amygdala for the Acquisition, Expression, and Extinction of Conditioned Fear. Physiological Reviews, 2010, 90, 419-463.	28.8	871
52	Neuropeptid S: Ein neues Transmittersystem im Gehirn. E-Neuroforum, 2009, 15, 56-62.	0.1	0
53	Deficiency of the 65 kDa Isoform of Glutamic Acid Decarboxylase Impairs Extinction of Cued But Not Contextual Fear Memory. Journal of Neuroscience, 2009, 29, 15713-15720.	3.6	90
54	Burst discharges in neurons of the thalamic reticular nucleus are shaped by calcium-induced calcium release. Cell Calcium, 2009, 46, 333-346.	2.4	49

#	Article	IF	CITATIONS
55	Correlation of T-channel coding gene expression, IT, and the low threshold Ca2+ spike in the thalamus of a rat model of absence epilepsy. Molecular and Cellular Neurosciences, 2008, 39, 384-399.	2.2	73
56	Neuropeptide Y activates a G-protein-coupled inwardly rectifying potassium current and dampens excitability in the lateral amygdala. Molecular and Cellular Neurosciences, 2008, 39, 491-498.	2.2	55
57	Neuropeptide S-Mediated Control of Fear Expression and Extinction: Role of Intercalated GABAergic Neurons in the Amygdala. Neuron, 2008, 59, 298-310.	8.1	271
58	Role of the somatostatin system in contextual fear memory and hippocampal synaptic plasticity. Learning and Memory, 2008, 15, 252-260.	1.3	35
59	Critical role of the 65-kDa isoform of glutamic acid decarboxylase in consolidation and generalization of Pavlovian fear memory. Learning and Memory, 2008, 15, 163-171.	1.3	95
60	Theta resynchronization during reconsolidation of remote contextual fear memory. NeuroReport, 2007, 18, 1107-1111.	1.2	55
61	Postsynaptic mechanisms underlying responsiveness of amygdaloid neurons to cholecystokinin are mediated by a transient receptor potential-like current. Molecular and Cellular Neurosciences, 2007, 35, 356-367.	2.2	49
62	Specific expression of low-voltage-activated calcium channel isoforms and splice variants in thalamic local circuit interneurons. Molecular and Cellular Neurosciences, 2007, 36, 132-145.	2.2	42
63	Plasticity of inhibitory synaptic network interactions in the lateral amygdala upon fear conditioning in mice. European Journal of Neuroscience, 2007, 25, 1205-1211.	2.6	38
64	Dissociated theta phase synchronization in amygdalo- hippocampal circuits during various stages of fear memory. European Journal of Neuroscience, 2007, 25, 1823-1831.	2.6	98
65	Molecular and functional properties of neurons in the human lateral amygdala. Molecular and Cellular Neurosciences, 2006, 31, 210-217.	2.2	14
66	Classification of projection neurons and interneurons in the rat lateral amygdala based upon cluster analysis. Molecular and Cellular Neurosciences, 2006, 33, 57-67.	2.2	64
67	Mechanisms of somatostatin-evoked responses in neurons of the rat lateral amygdala. European Journal of Neuroscience, 2005, 21, 755-762.	2.6	29
68	Theta activity in neurons and networks of the amygdala related to long-term fear memory. Hippocampus, 2005, 15, 874-880.	1.9	129
69	Impaired Regulation of Thalamic Pacemaker Channels through an Imbalance of Subunit Expression in Absence Epilepsy. Journal of Neuroscience, 2005, 25, 9871-9882.	3.6	103
70	Novel vistas of calcium-mediated signalling in the thalamus. Pflugers Archiv European Journal of Physiology, 2004, 448, 131-138.	2.8	49
71	Absence epilepsy and sinus dysrhythmia in mice lacking the pacemaker channel HCN2. EMBO Journal, 2003, 22, 216-224.	7.8	471
72	Amygdalar and Hippocampal Theta Rhythm Synchronization During Fear Memory Retrieval. Science, 2003, 301, 846-850.	12.6	662

#	Article	IF	CITATIONS
73	Generalisation of conditioned fear and its behavioural expression in mice. Behavioural Brain Research, 2003, 145, 89-98.	2.2	137
74	Contribution of NR2B Subunits to Synaptic Transmission in Amygdaloid Interneurons. Journal of Neuroscience, 2003, 23, 2549-2556.	3.6	46
75	Contribution of GABA _A and GABA _B Receptors to Thalamic Neuronal Activity during Spontaneous Absence Seizures in Rats. Journal of Neuroscience, 2001, 21, 1378-1384.	3.6	69
76	Contribution of intralaminar thalamic nuclei to spike-and-wave-discharges during spontaneous seizures in a genetic rat model of absence epilepsy. European Journal of Neuroscience, 2001, 13, 1537-1546.	2.6	71
77	Identification of Genes Expressed in the Amygdala During the Formation of Fear Memory. Learning and Memory, 2001, 8, 209-219.	1.3	73
78	Input-Specific Long-Term Depression in the Lateral Amygdala Evoked by Theta Frequency Stimulation. Journal of Neuroscience, 2000, 20, RC68-RC68.	3.6	60
79	Putative Cortical and Thalamic Inputs Elicit Convergent Excitation in a Population of GABAergic Interneurons of the Lateral Amygdala. Journal of Neuroscience, 2000, 20, 8909-8915.	3.6	105
80	Mechanisms and functional significance of a slow inhibitory potential in neurons of the lateral amygdala. European Journal of Neuroscience, 1998, 10, 853-867.	2.6	38
81	Relations between cortical and thalamic cellular activities during absence seizures in rats. European Journal of Neuroscience, 1998, 10, 1103-1112.	2.6	135
82	Two Types of Intrinsic Oscillations in Neurons of the Lateral and Basolateral Nuclei of the Amygdala. Journal of Neurophysiology, 1998, 79, 205-216.	1.8	108
83	Ionic Mechanisms of Intrinsic Oscillations in Neurons of the Basolateral Amygdaloid Complex. Journal of Neurophysiology, 1998, 79, 217-226.	1.8	91
84	Postsynaptic Mechanisms Underlying Responsiveness of Amygdaloid Neurons to Nociceptin/Orphanin FQ. Journal of Neuroscience, 1998, 18, 8133-8144.	3.6	72

6