

Perry L Wood Jr

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8984695/publications.pdf>

Version: 2024-02-01

109
papers

2,386
citations

304368

22
h-index

315357

38
g-index

117
all docs

117
docs citations

117
times ranked

1286
citing authors

#	ARTICLE	IF	CITATIONS
1	Integrated Analyses Resolve Conflicts over Squamate Reptile Phylogeny and Reveal Unexpected Placements for Fossil Taxa. PLoS ONE, 2015, 10, e0118199.	1.1	229
2	Phylogeny of bent-toed geckos (<i>Cyrtodactylus</i>) reveals a west to east pattern of diversification. Molecular Phylogenetics and Evolution, 2012, 65, 992-1003.	1.2	109
3	A phylogeny and taxonomy of the Thai-Malay Peninsula Bent-toed Geckos of the <i>Cyrtodactylus pulchellus</i> complex (Squamata: Gekkonidae): combined morphological and molecular analyses with descriptions of seven new species. Zootaxa, 2012, 3520, 1.	0.2	85
4	Systematics and natural history of Southeast Asian Rock Geckos (genus <i>Cnemaspis</i> Strauch, 1887) with descriptions of eight new species from Malaysia, Thailand, and Indonesia. Zootaxa, 2014, 3880, 1-147.	0.2	80
5	Twelve new species of <i>Cyrtodactylus</i> Gray (Squamata: Gekkonidae) from isolated limestone habitats in east-central and southern Myanmar demonstrate high localized diversity and unprecedented microendemism. Zoological Journal of the Linnean Society, 2018, 182, 862-959.	1.0	71
6	Identification of 104 rapidly-evolving nuclear protein-coding markers for amplification across scaled reptiles using genomic resources. Conservation Genetics Resources, 2012, 4, 1-10.	0.4	70
7	Integrative taxonomy uncovers high levels of cryptic species diversity in <i>Hemiphyllodactylus</i> Bleeker, 1860 (Squamata: Gekkonidae) and the description of a new species from Peninsular Malaysia. Zoological Journal of the Linnean Society, 2013, 169, 849-880.	1.0	54
8	Gene flow creates a mirage of cryptic species in a Southeast Asian spotted stream frog complex. Molecular Ecology, 2020, 29, 3970-3987.	2.0	49
9	Phylogeography, geographic variation, and taxonomy of the Bent-toed Gecko <i>Cyrtodactylus quadrivirgatus</i> Taylor, 1962 from Peninsular Malaysia with the description of a new swamp dwelling species. Zootaxa, 2012, 3406, 39.	0.2	45
10	A new species of karst forest Bent-toed Gecko (genus <i>Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 392 Td</i>) summary of Peninsular Malaysia's endemic karst forest herpetofauna and the need for its conservation. Zootaxa, 2016, 4061, 1.	0.2	42
11	Integrative taxonomy and preliminary assessment of species limits in the <i>Liolaemus walkeri</i> complex (Squamata, Liolaemidae) with descriptions of three new species from Peru. ZooKeys, 2013, 364, 47-91.	0.5	40
12	The phylogenetic relationships of three new species of the <i>Cyrtodactylus pulchellus</i> complex (Squamata: Gekkonidae) from poorly explored regions in northeastern Peninsular Malaysia. Zootaxa, 2014, 3786, 359.	0.2	37
13	The systematics and independent evolution of cave ecomorphology in distantly related clades of Bent-toed Geckos (Genus <i>Cyrtodactylus</i> Gray, 1827) from the Mekong Delta and islands in the Gulf of Thailand. Zootaxa, 2015, 3980, 106.	0.2	37
14	Three more new species of <i>Cyrtodactylus</i> (Squamata: Gekkonidae) from the Salween Basin of eastern Myanmar underscore the urgent need for the conservation of karst habitats. Journal of Natural History, 2018, 52, 1243-1294.	0.2	35
15	Evolution of habitat preference in 243 species of Bent-toed geckos (Genus <i>Cyrtodactylus</i> Gray,) Tj ETQq1 1,0,784314,rgBT /Overlock 10 Tf 50 392 Td	0.8	33
16	Six new species of the <i>Cyrtodactylus intermedius</i> complex (Squamata: Gekkonidae) from the Cardamom Mountains and associated highlands of Southeast Asia. Zootaxa, 2019, 4554, 1-62.	0.2	31
17	Parachute geckos free fall into synonymy: Gekko phylogeny, and a new subgeneric classification, inferred from thousands of ultraconserved elements. Molecular Phylogenetics and Evolution, 2020, 146, 106731.	1.2	31
18	Multiple origins of parthenogenesis, and a revised species phylogeny for the Southeast Asian butterfly lizards, <i>Leiolepis</i> . Biological Journal of the Linnean Society, 2014, 113, 1080-1093.	0.7	30

#	ARTICLE	IF	CITATIONS
19	Larger, unfiltered datasets are more effective at resolving phylogenetic conflict: Introns, exons, and UCEs resolve ambiguities in Golden-backed frogs (Anura: Ranidae; genus Hylarana). <i>Molecular Phylogenetics and Evolution</i> , 2020, 151, 106899.	1.2	30
20	Gene Flow Increases Phylogenetic Structure and Inflates Cryptic Species Estimations: A Case Study on Widespread Philippine Puddle Frogs (<i>Occidozyga laevis</i>). <i>Systematic Biology</i> , 2021, 71, 40-57.	2.7	30
21	Karstic Landscapes Are Foci of Species Diversity in the World's Third-Largest Vertebrate Genus <i>Cyrtodactylus</i> Gray, 1827 (Reptilia: Squamata; Gekkonidae). <i>Diversity</i> , 2021, 13, 183.	0.7	29
22	A new species of karst forest-adapted Bent-toed Gecko (genus <i>Cyrtodactylus</i> Gray, 1827) belonging to the <i>C. sworderi</i> complex from a threatened karst forest in Perak, Peninsular Malaysia. <i>Zootaxa</i> , 2014, 3755, 434-46.	0.2	28
23	<i>Cyrtodactylus</i> in the city: A new Bent-toed Gecko (Genus <i>Cyrtodactylus</i>) is the only endemic species of vertebrate from Batu Caves, Selangor, Peninsular Malaysia. <i>Zootaxa</i> , 2014, 3774, 381.	0.2	28
24	An integrative taxonomic review of the agamid genus <i>Bronchocela</i> (Kuhl, 1820) from Peninsular Malaysia with descriptions of new montane and insular endemics. <i>Zootaxa</i> , 2015, 3948, 1.	0.2	28
25	Two new diminutive species of <i>Cnemaspis</i> Strauch 1887 (Squamata: Gekkonidae) from Peninsular Malaysia. <i>Zootaxa</i> , 2009, 2019, 40-56.	0.2	27
26	A new <i>Cyrtodactylus</i> Gray, 1827 (Squamata, Gekkonidae) from the Shan Hills and the biogeography of Bent-toed Geckos from eastern Myanmar. <i>Zootaxa</i> , 2018, 4446, 477-500.	0.2	27
27	A new species of upland Stream Toad of the genus <i>Ansonia</i> Stoliczka, 1870 (Anura: Bufonidae) from northeastern Peninsular Malaysia. <i>Zootaxa</i> , 2014, 3764, 427.	0.2	26
28	Two new Bent-toed Geckos of the <i>Cyrtodactylus pulchellus</i> complex from Peninsular Malaysia and multiple instances of convergent adaptation to limestone forest ecosystems. <i>Zootaxa</i> , 2016, 4105, 401-29.	0.2	26
29	Phylogenetic partitioning of the third-largest vertebrate genus in the world, <i>Cyrtodactylus</i> Gray, 1827 (Reptilia; Squamata; Gekkonidae) and its relevance to taxonomy and conservation. <i>Vertebrate Zoology</i> , 2021, 71, 101-154.	2.0	25
30	Two New Species of Torrent-dwelling Toads <i>Ansonia</i> Stoliczka, 1870 (Anura: Bufonidae) from Peninsular Malaysia. <i>Herpetologica</i> , 2008, 64, 321-340.	0.2	24
31	The distribution, taxonomy, and redescription of the geckos <i>Cnemaspis affinis</i> (Stoliczka 1887) and <i>C. flavolineata</i> (Nicholls 1949) with descriptions of a new montane species and two new lowland, karst-dwelling species from Peninsular Malaysia. <i>Zootaxa</i> , 2008, 1931, 1-24.	0.2	24
32	Cryptic speciation within <i>Asthenodipsas vertebralis</i> (Boulenger, 1900) (Squamata: Tj ETQq0 0 0 rBT /Overlock 10 Tf). <i>Zootaxa</i> , 2013, 3664, 505-24.	0.2	24
33	Phylogenetic relationships and description of a new upland species of Bent-toed Gecko (<i>Cyrtodactylus</i>) Tj ETQq1 1 0.784314 rBT /Overlock 10 Tf. <i>Zootaxa</i> , 2013, 3664, 239-52.	0.2	23
34	Repeated evolution of sympatric, palaeoendemic species in closely related, co-distributed lineages of <i>Hemiphyllodactylus</i> Bleeker, 1860 (Squamata: Gekkonidae) across a sky-island archipelago in Peninsular Malaysia. <i>Zoological Journal of the Linnean Society</i> , 2015, 174, 859-876.	1.0	22
35	A new species of <i>Hemiphyllodactylus</i> Bleeker, 1860 (Squamata: Tj ETQq1 1 0.784314 rBT /Overlock 10 Tf). <i>Zootaxa</i> , 2013, 3664, 239-52.	0.2	21
36	Taxonomic and conservation implications of population genetic admixture, mito-nuclear discordance, and male-biased dispersal of a large endangered snake, <i>Drymarchon couperi</i> . <i>PLoS ONE</i> , 2019, 14, e0214439.	1.1	21

#	ARTICLE	IF	CITATIONS
37	Sex Chromosome Turnover in Bent-Toed Geckos (<i>Cyrtodactylus</i>). <i>Genes</i> , 2021, 12, 116.	1.0	21
38	A revision and redescription of the rock gecko <i>Cnemaspis siamensis</i> (Taylor 1925) (Squamata: Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 70	0.2	20
39	A new cryptic species of <i>Acanthosaura</i> Gray, 1831 (Squamata: Agamidae) from Thailand and Cambodia. <i>Zootaxa</i> , 2010, 2488, 22.	0.2	19
40	Phylogenetic relationships of geckos of the <i>Hemiphyllodactylus harterti</i> group, a new species from Penang Island, Peninsular Malaysia, and a likely case of true cryptic speciation. <i>Zootaxa</i> , 2016, 4107, 367-80.	0.2	19
41	Out of Borneo, again and again: biogeography of the Stream Toad genus <i>Ansonia</i> (Anura: Bufonidae) and the discovery of the first limestone cave-dwelling species. <i>Biological Journal of the Linnean Society</i> , 2016, , .	0.7	19
42	Dragons in the mist: three new species of <i>Pseudocalotes</i> Fitzinger (Squamata: Agamidae) from the sky island archipelago of Peninsular Malaysia. <i>Zootaxa</i> , 2016, 4136, 461-90.	0.2	19
43	<p class="HeadingRunIn">A new species of lowland karst dwelling <i>Cnemaspis</i> Strauch 1887 (Squamata: Gekkonidae) from northwestern Peninsular Malaysia</p>. <i>Zootaxa</i> , 2013, 3691, 538.	0.2	18
44	Fossil-calibrated phylogeny and historical biogeography of Southeast Asian water monitors (<i>Varanus</i>) Tj ETQq0 0 0 rgBT /Overlock 10 Tf	1.2	18
45	Systematics, ecomorphology, cryptic speciation and biogeography of the lizard genus <i>Tytthoscincus</i> Linkem, Diesmos & Brown (Squamata: Scincidae) from the sky-island archipelago of Peninsular Malaysia. <i>Zoological Journal of the Linnean Society</i> , 2018, 183, 635-671.	1.0	18
46	Two new montane species of <i>Acanthosaura</i> Gray, 1831 (Squamata: Agamidae) from Peninsular Malaysia. <i>Zootaxa</i> , 2009, 2012, 28-46.	0.2	18
47	A new insular species of Rock Gecko (<i>Cnemaspis</i> Boulenger) from Pulau Langkawi, Kedah, Peninsular Malaysia. <i>Zootaxa</i> , 2015, 3985, 203-18.	0.2	17
48	The fourth Bent-toed Gecko of the genus <i>Cyrtodactylus</i> (Squamata:) Tj ETQq0 0 0 rgBT /Overlock 1	0.2	17
49	Evolution of virulence in <i>Photorhabdus</i> spp., entomopathogenic nematode symbionts. <i>Systematic and Applied Microbiology</i> , 2016, 39, 173-179.	1.2	17
50	Target-capture phylogenomics provide insights on gene and species tree discordances in Old World treefrogs (Anura: Rhacophoridae). <i>Proceedings of the Royal Society B: Biological Sciences</i> , 2020, 287, 20202102.	1.2	17
51	FrogCap: A modular sequence capture probe–set for phylogenomics and population genetics for all frogs, assessed across multiple phylogenetic scales. <i>Molecular Ecology Resources</i> , 2022, 22, 1100-1119.	2.2	17
52	Peninsular Malaysia–s first limbless lizard: a new species of skink of the genus <i>Larutia</i> (BÄthme) from Pulau Pinang with a phylogeny of the genus. <i>Zootaxa</i> , 2011, 2799, .	0.2	16
53	A diminutive new species of cave-dwelling Wolf Snake (Colubridae: <i>Lycodon</i> Boie, 1826) from Peninsular Malaysia. <i>Zootaxa</i> , 2014, 3815, 51-67.	0.2	16
54	A new species of <i>Hemiphyllodactylus</i> Bleeker, 1860 (Squamata:) Tj ETQq0 0 0 rgBT /Overlock 10 Tf	0.2	16

#	ARTICLE	IF	CITATIONS
55	A re-description of <i>Cyrtodactylus chrysopylos</i> Bauer (Squamata: Gekkonidae) with comments on the adaptive significance of orange coloration in hatchlings and descriptions of two new species from eastern Myanmar (Burma). <i>Zootaxa</i> , 2018, 4527, 151.	0.2	16
56	Two more new species of <i>Hemiphyllodactylus</i> Bleeker (Squamata: Gekkonidae) from the Shan Hills of eastern Myanmar (Burma). <i>Zootaxa</i> , 2018, 4483, 295-316.	0.2	16
57	Two new species of <i>Cyrtodactylus</i> Gray, 1827 (Squamata: Gekkonidae) from a karstic archipelago in the Salween Basin of southern Myanmar (Burma). <i>Zootaxa</i> , 2020, 4718, zootaxa.4718.2.1.	0.2	16
58	Phylogenetic partitioning of the third-largest vertebrate genus in the world, <i>Cyrtodactylus</i> Gray, 1827 (Reptilia; Squamata; Gekkonidae) and its relevance to taxonomy and conservation. <i>Vertebrate Zoology</i> , 0, 71, 101-154.	2.0	16
59	A phylogenetic taxonomy of the <i>Cyrtodactylus peguensis</i> group (Reptilia: Squamata: Gekkonidae) with descriptions of two new species from Myanmar. <i>PeerJ</i> , 2018, 6, e5575.	0.9	16
60	A new species of insular Rock Gecko (Genus <i>Cnemaspis</i> Strauch, 1887) from the Bidong Archipelago, Terengganu, Peninsular Malaysia. <i>Zootaxa</i> , 2014, 3755, 447-56.	0.2	15
61	The first riparian skink (Genus: <i>Sphenomorphus</i> Strauch, 1887) from Peninsular Malaysia and its relationship to other Indochinese and Sundaic species. <i>Zootaxa</i> , 2016, 4173, 29.	0.2	15
62	Three new karst-dwelling <i>Cnemaspis</i> Strauch, 1887 (Squamata; Gekkonidae) from Peninsular Thailand and the phylogenetic placement of <i>C. punctatunuchalis</i> and <i>C. vandeventeri</i> . <i>PeerJ</i> , 2017, 5, e2884.	0.9	15
63	A new species of karst-adapted <i>Cnemaspis</i> Strauch, 1887 (Squamata: Gekkonidae) from a threatened karst region in Pahang, Peninsular Malaysia. <i>Zootaxa</i> , 2013, 3746, 463.	0.2	14
64	Phylogenetic taxonomy of <i>Hemiphyllodactylus</i> Bleeker, 1860 (Squamata: Gekkonidae) with descriptions of three new species from Myanmar. <i>Journal of Natural History</i> , 2018, 52, 881-915.	0.2	14
65	Integrative taxonomy of the Asian skinks <i>Sphenomorphus stellatus</i> (Boulenger, 1900) and <i>S. praesignis</i> (Boulenger, 1900) with the resurrection of <i>S. annamiticus</i> (Boettger, 1901) and the description of a new species from Cambodia. <i>Zootaxa</i> , 2019, 4683, zootaxa.4683.3.4.	0.2	14
66	Three new species of <i>Cyrtodactylus</i> (Squamata: Gekkonidae) from Peninsular Malaysia. <i>Zootaxa</i> , 2008, 1921, 1-23.	0.2	14
67	A new species of Gekko Laurenti, 1768 (Squamata: Gekkonidae) from Dong Nai Province, Southeastern Vietnam. <i>Zootaxa</i> , 2009, 2238, 33-42.	0.2	13
68	The phylogenetic relationships of a new Stream Toad of the genus <i>Ansonia</i> Stoliczka, 1870 (Anura: Tj ETQq0 0 0 rgBT /Overlock 10 Tf 5	0.2	13
69	The shadow of the past: Convergence of young and old South American desert lizards as measured by head shape traits. <i>Ecology and Evolution</i> , 2018, 8, 11399-11409.	0.8	13
70	The discovery and description of a new species of Bent-toed Gecko of the <i>Cyrtodactylus pulchellus</i> complex (Squamata: Tj ETQq0 0 0 rgBT /Overlock 10 Tf 5	0.2	12
71	Four new Burmese species of <i>Hemiphyllodactylus</i> Bleeker (Squamata: Gekkonidae) from distantly related parapatric clades from the Shan Plateau and Salween Basin. <i>Zootaxa</i> , 2020, 4758, zootaxa.4758.1.2.	0.2	12
72	<i>Cyrtodactylus rosichonariefi</i> sp. nov. (Squamata: Gekkonidae), a new swamp-dwelling bent-toed gecko from Bunguran Island (Great Natuna), Indonesia. <i>Zootaxa</i> , 2015, 3964, 114-24.	0.2	11

#	ARTICLE	IF	CITATIONS
73	On the taxonomy and phylogeny of the skinks <i>Lipinia sekayuensis</i> Grismer, Ismail, Awang, Rizal, & Ahmad and <i>Lipinia surda</i> Boulenger from Peninsular Malaysia. <i>Zootaxa</i> , 2016, 4147, 59.	0.2	11
74	A new species of Bent-toed Gecko (Squamata, Gekkonidae, <i>Cyrtodactylus</i>) from the Shan Plateau in eastern Myanmar (Burma). <i>Zootaxa</i> , 2019, 4624, zootaxa.4624.3.1.	0.2	11
75	Two more new species of the <i>Cyrtodactylus peguensis</i> group (Squamata: Gekkonidae) from the fringes of the Ayeyarwady Basin, Myanmar. <i>Zootaxa</i> , 2019, 4577, 274.	0.2	11
76	The biogeography of bent-toed geckos, <i>Cyrtodactylus</i> (Squamata: Gekkonidae). <i>PeerJ</i> , 2022, 10, e13153.	0.9	11
77	A new species of forest-dwelling <i>Cyrtodactylus</i> Gray (Squamata: Gekkonidae) from the Indawgyi Wildlife Sanctuary, Kachin State, Myanmar. <i>Zootaxa</i> , 2019, 4623, 1-25.	0.2	10
78	A new species of crocodile newt <i>Tylototriton</i> (Caudata: Salamandridae) from northern Myanmar (Burma). <i>Journal of Natural History</i> , 2019, 53, 475-495.	0.2	10
79	Geographic structure of genetic variation in the Parachute Gecko <i>Ptychozoon lionotum</i> Annandale, 1905 across Indochina and Sundaland with descriptions of three new species. <i>Zootaxa</i> , 2019, 4638, 151-198.	0.2	9
80	Rivers of Indochina as potential drivers of lineage diversification in the spotted flying lizard (<i>Draco</i>) Tj ETQq0 0 0 rgBTj/Overlock 10 Tf 50	1.2	9
81	Different roads lead to Rome: Integrative taxonomic approaches lead to the discovery of two new lizard lineages in the <i>Liolaemus montanus</i> group (Squamata: Liolaemidae). <i>Biological Journal of the Linnean Society</i> , 2016, , .	0.7	8
82	A new species of frog of the genus <i>Abavorana</i> Oliver, Prendini, Kraus & Raxworthy 2015 (Anura: Ranidae) from Gunung Jerai, Kedah, northwestern Peninsular Malaysia. <i>Zootaxa</i> , 2017, 4320, .	0.2	8
83	Systematics and natural history of mountain reed snakes (genus <i>Macrocalamus</i> ; Calamariinae). <i>Zoological Journal of the Linnean Society</i> , 2020, 188, 1236-1276.	1.0	8
84	A new potentially endangered limestone-associated Bent-toed Gecko of the <i>Cyrtodactylus pulchellus</i> (Squamata: Gekkonidae) complex from northern Peninsular Malaysia. <i>Zootaxa</i> , 2020, 4751, zootaxa.4751.3.2.	0.2	8
85	Unexpected Discovery of Another New Species of Philippine False Gecko (Gekkonidae; <i>Pseudogekko</i>) from the Bicol Peninsula of Luzon Island. <i>Herpetologica</i> , 2020, 76, 315.	0.2	7
86	The transfer of two clades of Malaysian <i>Sphenomorphus</i> Fitzinger (Squamata: Scincidae) into the genus <i>Tythoscincus</i> Linkem, Diesmos, & Brown and the description of a new Malaysian swamp-dwelling species. <i>Zootaxa</i> , 2016, 4092, 231-42.	0.2	6
87	A new species of Mud Snake (Serpentes, Homalopsidae, <i>Gyiophis</i> Murphy & Voris, 2014) from Myanmar with a first molecular phylogenetic assessment of the genus. <i>Zootaxa</i> , 2017, 4238, 571.	0.2	6
88	Morphological and molecular analyses reveal two new insular species of <i>Cnemaspis</i> Strauch, 1887 (Squamata, Gekkonidae) from Satun Province, southern Thailand. <i>ZooKeys</i> , 2019, 858, 127-161.	0.5	6
89	Phylogenetic and multivariate analyses of <i>Gekko smithii</i> Gray, 1842 recover a new species from Peninsular Malaysia and support the resurrection of <i>G. albomaculatus</i> (Giebel, 1861) from Sumatra. <i>Vertebrate Zoology</i> , 0, 72, 47-80.	2.0	6
90	A new species of lizard in the genus <i>Caledoniscincus</i> (Reptilia: Scincidae) from southern New Caledonia and a review of <i>Caledoniscincus</i> <i>atropunctatus</i> (Roux) . <i>Zootaxa</i> , 2013, 3694, 501.	0.2	5

#	ARTICLE	IF	CITATIONS
91	<i>Acanthosaura phuketensis</i> (Squamata: Agamidae), a new long-horned tree agamid from southwestern Thailand. <i>Zootaxa</i> , 2015, 4020, 473-94.	0.2	5
92	Geographically structured genetic variation in <i>Ptychozoon lionotum</i> (Squamata: Gekkonidae) and a new species from an isolated volcano in Myanmar. <i>Zootaxa</i> , 2018, 4514, 202.	0.2	5
93	Two new <i>Tytthoscincus</i> Linkem, Diesmos, & Brown (Squamata: Scincidae) from Peninsular Malaysia and another case of microsyntopy between ecologically specialized, unrelated, leaf-litter species. <i>Zootaxa</i> , 2018, 4425, 87.	0.2	5
94	The rediscovery of Schaefer's Spine-jawed Snake (<i>Xenophidion schaeferi</i> Günther & Manthey), a new record of the genus from Sumatra, Indonesia. <i>Zootaxa</i> , 2018, 4441, 366-378.	0.2	5
95	The taxonomy and phylogeny of the <i>Cyrtodactylus brevipalmatus</i> group (Squamata: Gekkonidae) with emphasis on <i>C. interdigitalis</i> and <i>C. ngati</i> . <i>Vertebrate Zoology</i> , 0, 72, 245-269.	2.0	5
96	Taxonomy, phylogeny, and distribution of <i>Bronchocela rayaensis</i> (Squamata: Agamidae) on the Thai-Malay Peninsula. <i>Zootaxa</i> , 2016, 4092, 414-20.	0.2	4
97	On the taxonomy and phylogeny of the rare Selangor Mud Snake (<i>Raclitia indica</i>) Gray (Serpentes). <i>Tropical Zoology</i> , 2014, 49, 1-14.	0.2	4
98	Discovery of the westernmost population of the genus <i>Ansonia</i> (Anura, Bufonidae) with the description of a new species from the Shan Plateau of eastern Myanmar. <i>Zootaxa</i> , 2019, 4656, 545-571.	0.2	4
99	Integrative taxonomy of the rock-dwelling gecko <i>Cnemaspis siamensis</i> complex (Squamata, Gekkonidae) reveals a new species from Nakhon Si Thammarat Province, southern Thailand. <i>ZooKeys</i> , 2020, 932, 129-159.	0.5	4
100	A new species of lowland karst dwelling <i>Cnemaspis</i> Strauch 1887 (Squamata: Gekkonidae) from northwestern Peninsular Malaysia. <i>Zootaxa</i> , 2013, 3691, 538-58.	0.2	4
101	Generalizing Bayesian phylogenetics to infer shared evolutionary events. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2022, 119, .	3.3	4
102	Taxonomic reassessment and phylogenetic placement of <i>Cyrtodactylus phuketensis</i> (Reptilia). <i>Tropical Zoology</i> , 2018, 53, 1-10.	0.5	3
103	A new species of scincid lizard in the genus <i>Caledoniscincus</i> (Reptilia: Scincidae) from northwest New Caledonia. <i>Zootaxa</i> , 2012, 3229, 47.	0.2	2
104	A new species of lizard in the genus <i>Caledoniscincus</i> (Reptilia: Scincidae) from far northwest New Caledonia. <i>Zootaxa</i> , 2014, 3795, 45.	0.2	2
105	A new species of <i>Cnemaspis</i> Strauch 1887 (Squamata: Gekkonidae) from the Langkawi Archipelago, Kedah, Peninsular Malaysia with an updated checklist of the herpetofauna of Tuba Island. <i>Zootaxa</i> , 2020, 4767, zootaxa.4767.1.6.	0.2	2
106	A new insular species of the <i>Cyrtodactylus pulchellus</i> group (Reptilia, Gekkonidae) from Tarutao Island, southern Thailand revealed by morphological and genetic evidence. <i>ZooKeys</i> , 2021, 1070, 101-134.	0.5	2
107	Two new syntopic species of wolf snakes (genus <i>Lycodon</i> H. Boie in Fitzinger, 1826) from an imperiled ecosystem in the Song Giang River Valley of southern Vietnam (Squamata: Colubridae). <i>Vertebrate Zoology</i> , 0, 72, 371-384.	2.0	1
108	A new <i>Liopeltis</i> Fitzinger, 1843 (Squamata: Colubridae) from Pulau Tioman, Peninsular Malaysia. <i>Zootaxa</i> , 2020, 4766, 472-484.	0.2	0

#	ARTICLE	IF	CITATIONS
109	Herpetofauna of gunung pantl forest reserve, johor, peninsular malaysia. Tropical Life Sciences Research, 2010, 21, 71-82.	0.5	0