

# Perry L Wood Jr

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8984695/publications.pdf>

Version: 2024-02-01

109  
papers

2,386  
citations

304368

22  
h-index

315357

38  
g-index

117  
all docs

117  
docs citations

117  
times ranked

1286  
citing authors

#	ARTICLE	IF	CITATIONS
1	FrogCap: A modular sequence capture probe set for phylogenomics and population genetics for all frogs, assessed across multiple phylogenetic scales. <i>Molecular Ecology Resources</i> , 2022, 22, 1100-1119.	2.2	17
2	The biogeography of bent-toed geckos, <i>Cyrtodactylus</i> (Squamata: Gekkonidae). <i>PeerJ</i> , 2022, 10, e13153.	0.9	11
3	Generalizing Bayesian phylogenetics to infer shared evolutionary events. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2022, 119, .	3.3	4
4	Sex Chromosome Turnover in Bent-Toed Geckos ( <i>Cyrtodactylus</i> ). <i>Genes</i> , 2021, 12, 116.	1.0	21
5	Karstic Landscapes Are Foci of Species Diversity in the World's Third-Largest Vertebrate Genus <i>Cyrtodactylus</i> Gray, 1827 (Reptilia: Squamata; Gekkonidae). <i>Diversity</i> , 2021, 13, 183.	0.7	29
6	Taxonomic reassessment and phylogenetic placement of <i>Cyrtodactylus phuketensis</i> (Reptilia, Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 542	0.5	8
7	Gene Flow Increases Phylogenetic Structure and Inflates Cryptic Species Estimations: A Case Study on Widespread Philippine Puddle Frogs ( <i>Occidozyga laevis</i> ). <i>Systematic Biology</i> , 2021, 71, 40-57.	2.7	30
8	A new insular species of the <i>Cyrtodactylus pulchellus</i> group (Reptilia, Gekkonidae) from Tarutao Island, southern Thailand revealed by morphological and genetic evidence. <i>ZooKeys</i> , 2021, 1070, 101-134.	0.5	2
9	Systematics and natural history of mountain reed snakes (genus <i>Macrocalamus</i> ; Calamariinae). <i>Zoological Journal of the Linnean Society</i> , 2020, 188, 1236-1276.	1.0	8
10	Parachute geckos free fall into synonymy: Gekko phylogeny, and a new subgeneric classification, inferred from thousands of ultraconserved elements. <i>Molecular Phylogenetics and Evolution</i> , 2020, 146, 106731.	1.2	31
11	Two new species of <i>Cyrtodactylus</i> Gray, 1827 (Squamata: Gekkonidae) from a karstic archipelago in the Salween Basin of southern Myanmar (Burma). <i>Zootaxa</i> , 2020, 4718, zootaxa.4718.2.1.	0.2	16
12	Evolution of habitat preference in 243 species of Bent-toed geckos (Genus <i>Cyrtodactylus</i> Gray,) Tj ETQq0 0 0 rgBT /Overlock 10	0.8	33
13	Gene flow creates a mirage of cryptic species in a Southeast Asian spotted stream frog complex. <i>Molecular Ecology</i> , 2020, 29, 3970-3987.	2.0	49
14	Target-capture phylogenomics provide insights on gene and species tree discordances in Old World treefrogs (Anura: Rhacophoridae). <i>Proceedings of the Royal Society B: Biological Sciences</i> , 2020, 287, 20202102.	1.2	17
15	Rivers of Indochina as potential drivers of lineage diversification in the spotted flying lizard ( <i>Draco</i> ) Tj ETQq1 1 0.784314 rgBT /Overlock	1.2	14
16	Larger, unfiltered datasets are more effective at resolving phylogenetic conflict: Introns, exons, and UCEs resolve ambiguities in Golden-backed frogs (Anura: Ranidae; genus <i>Hylarana</i> ). <i>Molecular Phylogenetics and Evolution</i> , 2020, 151, 106899.	1.2	30
17	Four new Burmese species of <i>Hemiphyllodactylus</i> Bleeker (Squamata: Gekkonidae) from distantly related parapatric clades from the Shan Plateau and Salween Basin. <i>Zootaxa</i> , 2020, 4758, zootaxa.4758.1.2.	0.2	12
18	<strong>A new <i>Liopeltis</i> Fitzinger, 1843 (Squamata: Colubridae) from Pulau Tioman, Peninsular Malaysia</strong> . <i>Zootaxa</i> , 2020, 4766, 472-484.	0.2	0

#	ARTICLE	IF	CITATIONS
19	A new species of <i>Cnemaspis</i> Strauch 1887 (Squamata: Gekkonidae) from the Langkawi Archipelago, Kedah, Peninsular Malaysia with an updated checklist of the herpetofauna of Tuba Island. <i>Zootaxa</i> , 2020, 4767, zootaxa.4767.1.6.	0.2	2
20	A new potentially endangered limestone-associated Bent-toed Gecko of the <i>Cyrtodactylus pulchellus</i> (Squamata: Gekkonidae) complex from northern Peninsular Malaysia. <i>Zootaxa</i> , 2020, 4751, zootaxa.4751.3.2.	0.2	8
21	Unexpected Discovery of Another New Species of Philippine False Gecko (Gekkonidae; <i>Pseudogekko</i> ) from the Bicol Peninsula of Luzon Island. <i>Herpetologica</i> , 2020, 76, 315.	0.2	7
22	Integrative taxonomy of the rock-dwelling gecko <i>Cnemaspis siamensis</i> complex (Squamata, Gekkonidae) reveals a new species from Nakhon Si Thammarat Province, southern Thailand. <i>ZooKeys</i> , 2020, 932, 129-159.	0.5	4
23	Geographic structure of genetic variation in the Parachute Gecko <i>Ptychozoon lionotum</i> Annandale, 1905 across Indochina and Sundaland with descriptions of three new species. <i>Zootaxa</i> , 2019, 4638, 151-198.	0.2	9
24	A new species of Bent-toed Gecko (Squamata, Gekkonidae, <i>Cyrtodactylus</i> ) from the Shan Plateau in eastern Myanmar (Burma). <i>Zootaxa</i> , 2019, 4624, zootaxa.4624.3.1.	0.2	11
25	Integrative taxonomy of the Asian skinks <i>Sphenomorphus stellatus</i> (Boulenger, 1900) and <i>S. praesignis</i> (Boulenger, 1900) with the resurrection of <i>S. annamiticus</i> (Boettger, 1901) and the description of a new species from Cambodia. <i>Zootaxa</i> , 2019, 4683, zootaxa.4683.3.4.	0.2	14
26	Discovery of the westernmost population of the genus <i>Ansonia</i> (Anura, Bufonidae) with the description of a new species from the Shan Plateau of eastern Myanmar. <i>Zootaxa</i> , 2019, 4656, 545-571.	0.2	4
27	A new species of forest-dwelling <i>Cyrtodactylus</i> Gray (Squamata: Gekkonidae) from the Indawgyi Wildlife Sanctuary, Kachin State, Myanmar. <i>Zootaxa</i> , 2019, 4623, 1-25.	0.2	10
28	The discovery and description of a new species of Bent-toed Gecko of the <i>Cyrtodactylus pulchellus</i> complex (Squamata: Gekkonidae) from northern Peninsular Malaysia. <i>Zootaxa</i> , 2019, 4668, 51-75.	0.2	12
29	A new species of crocodile newt <i>Tylotriton</i> (Caudata: Salamandridae) from northern Myanmar (Burma). <i>Journal of Natural History</i> , 2019, 53, 475-495.	0.2	10
30	Taxonomic and conservation implications of population genetic admixture, mito-nuclear discordance, and male-biased dispersal of a large endangered snake, <i>Drymarchon couperi</i> . <i>PLoS ONE</i> , 2019, 14, e0214439.	1.1	21
31	Two more new species of the <i>Cyrtodactylus peguensis</i> group (Squamata: Gekkonidae) from the fringes of the Ayeyarwady Basin, Myanmar. <i>Zootaxa</i> , 2019, 4577, 274.	0.2	11
32	Six new species of the <i>Cyrtodactylus intermedius</i> complex (Squamata: Gekkonidae) from the Cardamom Mountains and associated highlands of Southeast Asia. <i>Zootaxa</i> , 2019, 4554, 1-62.	0.2	31
33	Morphological and molecular analyses reveal two new insular species of <i>Cnemaspis</i> Strauch, 1887 (Squamata, Gekkonidae) from Satun Province, southern Thailand. <i>ZooKeys</i> , 2019, 858, 127-161.	0.5	6
34	Three more new species of <i>Cyrtodactylus</i> (Squamata: Gekkonidae) from the Salween Basin of eastern Myanmar underscore the urgent need for the conservation of karst habitats. <i>Journal of Natural History</i> , 2018, 52, 1243-1294.	0.2	35
35	Twelve new species of <i>Cyrtodactylus</i> Gray (Squamata: Gekkonidae) from isolated limestone habitats in east-central and southern Myanmar demonstrate high localized diversity and unprecedented microendemism. <i>Zoological Journal of the Linnean Society</i> , 2018, 182, 862-959.	1.0	71
36	Systematics, ecomorphology, cryptic speciation and biogeography of the lizard genus <i>Tytthoscincus</i> Linkem, Diesmos & Brown (Squamata: Scincidae) from the sky-island archipelago of Peninsular Malaysia. <i>Zoological Journal of the Linnean Society</i> , 2018, 183, 635-671.	1.0	18

#	ARTICLE	IF	CITATIONS
37	Phylogenetic taxonomy of <i>Hemiphyllodactylus</i> Bleeker, 1860 (Squamata: Gekkonidae) with descriptions of three new species from Myanmar. <i>Journal of Natural History</i> , 2018, 52, 881-915.	0.2	14
38	On the taxonomy and phylogeny of the rare Selangor Mud Snake ( <i>Raclitia indica</i> ) Gray (Serpentes). <i>Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50</i>	0.2	4
39	Geographically structured genetic variation in <i>Ptychozoon lionotum</i> (Squamata: Gekkonidae) and a new species from an isolated volcano in Myanmar. <i>Zootaxa</i> , 2018, 4514, 202.	0.2	5
40	The shadow of the past: Convergence of young and old South American desert lizards as measured by head shape traits. <i>Ecology and Evolution</i> , 2018, 8, 11399-11409.	0.8	13
41	A re-description of <i>Cyrtodactylus chrysopylos</i> Bauer (Squamata: Gekkonidae) with comments on the adaptive significance of orange coloration in hatchlings and descriptions of two new species from eastern Myanmar (Burma). <i>Zootaxa</i> , 2018, 4527, 151.	0.2	16
42	Two more new species of <i>Hemiphyllodactylus</i> Bleeker (Squamata: Gekkonidae) from the Shan Hills of eastern Myanmar (Burma). <i>Zootaxa</i> , 2018, 4483, 295-316.	0.2	16
43	Two new <i>Tytthoscincus</i> Linkem, Diesmos, & Brown (Squamata; Scincidae) from Peninsular Malaysia and another case of microsyntopy between ecologically specialized, unrelated, leaf-litter species. <i>Zootaxa</i> , 2018, 4425, 87.	0.2	5
44	The rediscovery of Schaefer's Spine-jawed Snake ( <i>Xenophidion schaeferi</i> Günther & Manthey) record of the genus from Sumatra, Indonesia. <i>Zootaxa</i> , 2018, 4441, 366-378.	0.2	5
45	A new <i>Cyrtodactylus</i> Gray, 1827 (Squamata, Gekkonidae) from the Shan Hills and the biogeography of Bent-toed Geckos from eastern Myanmar. <i>Zootaxa</i> , 2018, 4446, 477-500.	0.2	27
46	A phylogenetic taxonomy of the <i>Cyrtodactylus peguensis</i> group (Reptilia: Squamata: Gekkonidae) with descriptions of two new species from Myanmar. <i>PeerJ</i> , 2018, 6, e5575.	0.9	16
47	A new species of Mud Snake (Serpentes, Homalopsidae, <i>Gyiophis</i> Murphy & Voris, 2014) from Myanmar with a first molecular phylogenetic assessment of the genus. <i>Zootaxa</i> , 2017, 4238, 571.	0.2	6
48	A new species of frog of the genus <i>Abavorana</i> Oliver, Prendini, Kraus & Raxworthy 2015 (Anura: Ranidae) from Gunung Jerai, Kedah, northwestern Peninsular Malaysia. <i>Zootaxa</i> , 2017, 4320, .	0.2	8
49	Three new karst-dwelling <i>Cnemaspis</i> Strauch, 1887 (Squamata; Gekkonidae) from Peninsular Thailand and the phylogenetic placement of <i>C. punctatouchalis</i> and <i>C. vandeventeri</i> . <i>PeerJ</i> , 2017, 5, e2884.	0.9	15
50	A new species of karst forest Bent-toed Gecko (genus <i>Tytthoscincus</i> ) from Peninsular Malaysia's endemic karst forest herpetofauna and the need for its conservation. <i>Zootaxa</i> , 2016, 4061, 1.	0.2	42
51	The transfer of two clades of Malaysian <i>Sphenomorphus</i> Fitzinger (Squamata: Scincidae) into the genus <i>Tytthoscincus</i> Linkem, Diesmos, & Brown and the description of a new Malaysian swamp-dwelling species. <i>Zootaxa</i> , 2016, 4092, 231-42.	0.2	6
52	Taxonomy, phylogeny, and distribution of <i>Bronchocela rayaensis</i> (Squamata: Agamidae) on the Thai-Malay Peninsula. <i>Zootaxa</i> , 2016, 4092, 414-20.	0.2	4
53	The phylogenetic relationships of a new Stream Toad of the genus <i>Ansonia</i> Stoliczka, 1870 (Anura). <i>Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf 50 232 Td</i>	0.2	13
54	Phylogenetic relationships of geckos of the <i>Hemiphyllodactylus harterti</i> group, a new species from Penang Island, Peninsular Malaysia, and a likely case of true cryptic speciation. <i>Zootaxa</i> , 2016, 4107, 367-80.	0.2	19

#	ARTICLE	IF	CITATIONS
55	Two new Bent-toed Geckos of the <i>Cyrtodactylus pulchellus</i> complex from Peninsular Malaysia and multiple instances of convergent adaptation to limestone forest ecosystems. <i>Zootaxa</i> , 2016, 4105, 401-29.	0.2	26
56	Different roads lead to Rome: Integrative taxonomic approaches lead to the discovery of two new lizard lineages in the <i>Liolaemus montanus</i> group (Squamata: Liolaemidae). <i>Biological Journal of the Linnean Society</i> , 2016, , .	0.7	8
57	The first riparian skink (Genus: <i>Sphenomorphus</i> Strauch, 1887) from Peninsular Malaysia and its relationship to other Indochinese and Sundaic species. <i>Zootaxa</i> , 2016, 4173, 29.	0.2	15
58	On the taxonomy and phylogeny of the skinks <i>Lipinia sekayuensis</i> Grismer, Ismail, Awang, Rizal, & Ahmad and <i>Lipinia surda</i> Boulenger from Peninsular Malaysia. <i>Zootaxa</i> , 2016, 4147, 59.	0.2	11
59	Out of Borneo, again and again: biogeography of the Stream Toad genus <i>Ansonia</i> Stoliczka (Anura: Bufonidae) and the discovery of the first limestone cave-dwelling species. <i>Biological Journal of the Linnean Society</i> , 2016, , .	0.7	19
60	Dragons in the mist: three new species of <i>Pseudocalotes</i> Fitzinger (Squamata: Agamidae) from the sky island archipelago of Peninsular Malaysia. <i>Zootaxa</i> , 2016, 4136, 461-90.	0.2	19
61	Evolution of virulence in <i>Photorhabdus</i> spp., entomopathogenic nematode symbionts. <i>Systematic and Applied Microbiology</i> , 2016, 39, 173-179.	1.2	17
62	Repeated evolution of sympatric, palaeoendemic species in closely related, co-distributed lineages of <i>Hemiphyllodactylus</i> Bleeker, 1860 (Squamata: Gekkonidae) across a sky-island archipelago in Peninsular Malaysia. <i>Zoological Journal of the Linnean Society</i> , 2015, 174, 859-876.	1.0	22
63	<i>Acanthosaura phuketensis</i> (Squamata: Agamidae), a new long-horned tree agamid from southwestern Thailand. <i>Zootaxa</i> , 2015, 4020, 473-94.	0.2	5
64	A new insular species of Rock Gecko ( <i>Cnemaspis</i> Boulenger) from Pulau Langkawi, Kedah, Peninsular Malaysia. <i>Zootaxa</i> , 2015, 3985, 203-18.	0.2	17
65	The systematics and independent evolution of cave ecomorphology in distantly related clades of Bent-toed Geckos (Genus <i>Cyrtodactylus</i> Gray, 1827) from the Mekong Delta and islands in the Gulf of Thailand. <i>Zootaxa</i> , 2015, 3980, 106.	0.2	37
66	<i>Cyrtodactylus rosichonariefi</i> sp. nov. (Squamata: Gekkonidae), a new swamp-dwelling bent-toed gecko from Bunguran Island (Great Natuna), Indonesia. <i>Zootaxa</i> , 2015, 3964, 114-24.	0.2	11
67	The fourth Bent-toed Gecko of the genus <i>Cyrtodactylus</i> (Squamata: Gekkonidae) from the island of Sumatra. <i>Zootaxa</i> , 2015, 3964, 114-24.	0.2	17
68	Integrated Analyses Resolve Conflicts over Squamate Reptile Phylogeny and Reveal Unexpected Placements for Fossil Taxa. <i>PLoS ONE</i> , 2015, 10, e0118199.	1.1	229
69	An integrative taxonomic review of the agamid genus <i>Bronchocela</i> (Kuhl, 1820) from Peninsular Malaysia with descriptions of new montane and insular endemics. <i>Zootaxa</i> , 2015, 3948, 1.	0.2	28
70	A new species of karst forest-adapted Bent-toed Gecko (genus <i>Cyrtodactylus</i> Gray, 1827) belonging to the <i>C. sworderi</i> complex from a threatened karst forest in Perak, Peninsular Malaysia. <i>Zootaxa</i> , 2014, 3755, 434-46.	0.2	28
71	A new species of insular Rock Gecko (Genus <i>Cnemaspis</i> Strauch, 1887) from the Bidong Archipelago, Terengganu, Peninsular Malaysia. <i>Zootaxa</i> , 2014, 3755, 447-56.	0.2	15
72	The phylogenetic relationships of three new species of the <i>Cyrtodactylus pulchellus</i> complex (Squamata: Gekkonidae) from poorly explored regions in northeastern Peninsular Malaysia. <i>Zootaxa</i> , 2014, 3786, 359.	0.2	37

#	ARTICLE	IF	CITATIONS
73	<strong>A new species of upland Stream Toad of the genus <i>Ansonia</i> Stoliczka, 1870 (Anura: Bufonidae) from northeastern Peninsular Malaysia</strong> . <i>Zootaxa</i> , 2014, 3764, 427.	0.2	26
74	A diminutive new species of cave-dwelling Wolf Snake (Colubridae: <i>Lycodon</i> Boie, 1826) from Peninsular Malaysia. <i>Zootaxa</i> , 2014, 3815, 51-67.	0.2	16
75	Systematics and natural history of Southeast Asian Rock Geckos (genus <i>Cnemaspis</i> Strauch, 1887) with descriptions of eight new species from Malaysia, Thailand, and Indonesia. <i>Zootaxa</i> , 2014, 3880, 1-147.	0.2	80
76	<strong>A new species of lizard in the genus <i>Caledoniscincus</i> (Reptilia: Scincidae) from far northwest New Caledonia</strong> . <i>Zootaxa</i> , 2014, 3795, 45.	0.2	2
77	Multiple origins of parthenogenesis, and a revised species phylogeny for the Southeast Asian butterfly lizards, <i>Leiolepis</i> . <i>Biological Journal of the Linnean Society</i> , 2014, 113, 1080-1093.	0.7	30
78	Fossil-calibrated phylogeny and historical biogeography of Southeast Asian water monitors ( <i>Varanus</i> )	1.2	18
79	<strong>A new species of <i>Hemiphyllodactylus</i> Bleeker, 1860 (Squamata: Gekkonidae)</strong> . <i>Zootaxa</i> , 2014, 3774, 381.	0.2	16
80	<strong>A new species of <i>Hemiphyllodactylus</i> Bleeker, 1860 (Squamata: Gekkonidae)</strong> . <i>Zootaxa</i> , 2014, 3774, 381.	0.2	21
81	<strong>Cyrtos in the city: A new Bent-toed Gecko (Genus <i>Cyrtodactylus</i>) is the only endemic species of vertebrate from Batu Caves, Selangor, Peninsular Malaysia</strong> . <i>Zootaxa</i> , 2014, 3774, 381.	0.2	28
82	Integrative taxonomy uncovers high levels of cryptic species diversity in <i>Hemiphyllodactylus</i> Bleeker, 1860 (Squamata: Gekkonidae) and the description of a new species from Peninsular Malaysia. <i>Zoological Journal of the Linnean Society</i> , 2013, 169, 849-880.	1.0	54
83	Integrative taxonomy and preliminary assessment of species limits in the <i>Liolaemus walkeri</i> complex (Squamata, Liolaemidae) with descriptions of three new species from Peru. <i>ZooKeys</i> , 2013, 364, 47-91.	0.5	40
84	<strong>A new species of karst-adapted <i>Cnemaspis</i> Strauch, 1887 (Squamata: Gekkonidae) from a threatened karst region in Pahang, Peninsular Malaysia</strong> . <i>Zootaxa</i> , 2013, 3746, 463.	0.2	14
85	<strong>A new species of lowland karst dwelling <i>Cnemaspis</i> Strauch 1887 (Squamata: Gekkonidae) from northwestern Peninsular Malaysia</strong> . <i>Zootaxa</i> , 2013, 3691, 538.	0.2	18
86	Cryptic speciation within <i>Asthenodipsas vertebralis</i> (Boulenger, 1900) (Squamata: Gekkonidae)	0.2	24
87	<strong>A. tropidonotus</strong> (Lidth de Jude, 1923) from Sumatra: an integrative taxonomic analysis. <i>Zootaxa</i> , 2013, 3664, 505-24.	0.2	23
88	Phylogenetic relationships and description of a new upland species of Bent-toed Gecko ( <i>Cyrtodactylus</i> )	0.2	23
89	<strong>A new species of lizard in the genus <i>Caledoniscincus</i> (Reptilia: Scincidae) from southern New Caledonia and a review of <i>Caledoniscincus atropunctatus</i> (Roux)</strong> . <i>Zootaxa</i> , 2013, 3694, 501.	0.2	5
90	A new species of lowland karst dwelling <i>Cnemaspis</i> Strauch 1887 (Squamata: Gekkonidae) from northwestern Peninsular Malaysia. <i>Zootaxa</i> , 2013, 3691, 538-58.	0.2	4
90	Phylogeny of bent-toed geckos ( <i>Cyrtodactylus</i> ) reveals a west to east pattern of diversification. <i>Molecular Phylogenetics and Evolution</i> , 2012, 65, 992-1003.	1.2	109

#	ARTICLE	IF	CITATIONS
91	Phylogeography, geographic variation, and taxonomy of the Bent-toed Gecko <i>Cyrtodactylus quadrivirgatus</i> Taylor, 1962 from Peninsular Malaysia with the description of a new swamp dwelling species. <i>Zootaxa</i> , 2012, 3406, 39.	0.2	45
92	A new species of scincid lizard in the genus <i>Caledoniscincus</i> (Reptilia: Scincidae) from northwest New Caledonia. <i>Zootaxa</i> , 2012, 3229, 47.	0.2	2
93	A phylogeny and taxonomy of the Thai-Malay Peninsula Bent-toed Geckos of the <i>Cyrtodactylus pulchellus</i> complex (Squamata: Gekkonidae): combined morphological and molecular analyses with descriptions of seven new species. <i>Zootaxa</i> , 2012, 3520, 1.	0.2	85
94	Identification of 104 rapidly-evolving nuclear protein-coding markers for amplification across scaled reptiles using genomic resources. <i>Conservation Genetics Resources</i> , 2012, 4, 1-10.	0.4	70
95	Peninsular Malaysia's first limbless lizard: a new species of skink of the genus <i>Larutia</i> (Blyth) from Pulau Pinang with a phylogeny of the genus. <i>Zootaxa</i> , 2011, 2799, .	0.2	16
96	A new cryptic species of <i>Acanthosaura</i> Gray, 1831 (Squamata: Agamidae) from Thailand and Cambodia. <i>Zootaxa</i> , 2010, 2488, 22.	0.2	19
97	A revision and redescription of the rock gecko <i>Cnemaspis siamensis</i> (Taylor 1925) (Squamata: Tj ETQq1 1 0.784314,rgBT /Oyerlock 10 0,2 20	0.2	20
98	Herpetofauna of gunung panti forest reserve, johor, peninsular malaysia. <i>Tropical Life Sciences Research</i> , 2010, 21, 71-82.	0.5	0
99	Two new diminutive species of <i>Cnemaspis</i> Strauch 1887 (Squamata: Gekkonidae) from Peninsular Malaysia. <i>Zootaxa</i> , 2009, 2019, 40-56.	0.2	27
100	A new species of <i>Gekko Laurenti</i> , 1768 (Squamata: Gekkonidae) from Dong Nai Province, Southeastern Vietnam. <i>Zootaxa</i> , 2009, 2238, 33-42.	0.2	13
101	Two new montane species of <i>Acanthosaura</i> Gray, 1831 (Squamata: Agamidae) from Peninsular Malaysia. <i>Zootaxa</i> , 2009, 2012, 28-46.	0.2	18
102	Two New Species of Torrent-dwelling Toads <i>Ansonia Stoliczka</i> , 1870 (Anura: Bufonidae) from Peninsular Malaysia. <i>Herpetologica</i> , 2008, 64, 321-340.	0.2	24
103	The distribution, taxonomy, and redescription of the geckos <i>Cnemaspis affinis</i> (Stoliczka 1887) and <i>C. flavolineata</i> (Nicholls 1949) with descriptions of a new montane species and two new lowland, karst-dwelling species from Peninsular Malaysia. <i>Zootaxa</i> , 2008, 1931, 1-24.	0.2	24
104	Three new species of <i>Cyrtodactylus</i> (Squamata: Gekkonidae) from Peninsular Malaysia. <i>Zootaxa</i> , 2008, 1921, 1-23.	0.2	14
105	Phylogenetic partitioning of the third-largest vertebrate genus in the world, <i>Cyrtodactylus</i> Gray, 1827 (Reptilia; Squamata; Gekkonidae) and its relevance to taxonomy and conservation. <i>Vertebrate Zoology</i> , 0, 71, 101-154.	2.0	25
106	Phylogenetic partitioning of the third-largest vertebrate genus in the world, <i>Cyrtodactylus</i> Gray, 1827 (Reptilia; Squamata; Gekkonidae) and its relevance to taxonomy and conservation. <i>Vertebrate Zoology</i> , 0, 71, 101-154.	2.0	16
107	Phylogenetic and multivariate analyses of <i>Gekko smithii</i> Gray, 1842 recover a new species from Peninsular Malaysia and support the resurrection of <i>G. albomaculatus</i> (Giebel, 1861) from Sumatra. <i>Vertebrate Zoology</i> , 0, 72, 47-80.	2.0	6
108	The taxonomy and phylogeny of the <i>Cyrtodactylus brevipalmatus</i> group (Squamata: Gekkonidae) with emphasis on <i>C. interdigitalis</i> and <i>C. ngati</i> . <i>Vertebrate Zoology</i> , 0, 72, 245-269.	2.0	5

#	ARTICLE	IF	CITATIONS
109	Two new syntopic species of wolf snakes (genus <i>Lycodon</i> H. Boie in Fitzinger, 1826) from an imperiled ecosystem in the Song Giang River Valley of southern Vietnam (Squamata: Colubridae). <i>Vertebrate Zoology</i> , 0, 72, 371-384.	2.0	1