
## Faxi Wang

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/8987262/publications.pdf Version: 2024-02-01



FAXI MANC

#	Article	IF	CITATIONS
1	Loss of ubiquitin-conjugating enzyme E2 (Ubc9) in macrophages exacerbates multiple low-dose streptozotocin-induced diabetes by attenuating M2 macrophage polarization. Cell Death and Disease, 2019, 10, 892.	2.7	44
2	Aloperine Protects Mice against DSS-Induced Colitis by PP2A-Mediated PI3K/Akt/mTOR Signaling Suppression. Mediators of Inflammation, 2017, 2017, 1-14.	1.4	40
3	Kdm2a deficiency in macrophages enhances thermogenesis to protect mice against HFD-induced obesity by enhancing H3K36me2 at the Pparg locus. Cell Death and Differentiation, 2021, 28, 1880-1899.	5.0	33
4	Soluble FGL2, a novel effector molecule of activated hepatic stellate cells, regulates T-cell function in cirrhotic patients with hepatocellular carcinoma. Hepatology International, 2014, 8, 567-575.	1.9	26
5	Targeted Inhibition of FTO Demethylase Protects Mice Against LPS-Induced Septic Shock by Suppressing NLRP3 Inflammasome. Frontiers in Immunology, 2021, 12, 663295.	2.2	26
6	A Review on Recent Advances in Aloperine Research: Pharmacological Activities and Underlying Biological Mechanisms. Frontiers in Pharmacology, 2020, 11, 538137.	1.6	23
7	Extracellular HMGB1 exacerbates autoimmune progression and recurrence of type 1 diabetes by impairing regulatory T cell stability. Diabetologia, 2020, 63, 987-1001.	2.9	23
8	The AHR Signaling Attenuates Autoimmune Responses During the Development of Type 1 Diabetes. Frontiers in Immunology, 2020, 11, 1510.	2.2	21
9	MBD2 acts as a repressor to maintain the homeostasis of the Th1 program in type 1 diabetes by regulating the STAT1-IFN-Î <sup>3</sup> axis. Cell Death and Differentiation, 2022, 29, 218-229.	5.0	18
10	Expression of Interferon Effector Gene SART1 Correlates with Interferon Treatment Response against Hepatitis B Infection. Mediators of Inflammation, 2016, 2016, 1-11.	1.4	13
11	Ubc9 deficiency selectively impairs the functionality of common lymphoid progenitors (CLPs) during bone marrow hematopoiesis. Molecular Immunology, 2019, 114, 314-322.	1.0	3