

Felice Senatore

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8995654/publications.pdf>

Version: 2024-02-01

158
papers

5,156
citations

94269

37
h-index

114278

63
g-index

160
all docs

160
docs citations

160
times ranked

5677
citing authors

#	ARTICLE	IF	CITATIONS
1	Antibacterial Activity of <i>Cuminum cyminum</i> L. and <i>Carum carvi</i> L. Essential Oils. Journal of Agricultural and Food Chemistry, 2005, 53, 57-61.	2.4	369
2	Antibacterial Activity of <i>Coriandrum sativum</i> L. and <i>Foeniculum vulgare</i> Miller Var. <i>vulgare</i> (Miller) Essential Oils. Journal of Agricultural and Food Chemistry, 2004, 52, 7862-7866.	2.4	226
3	Chemical composition and anticancer activity of essential oils of Mediterranean sage (<i>Salvia</i>) Tj ETQq1 1 0.784314 rgBT /Overlock 10 42-47.	1.8	172
4	Influence of Harvesting Time on Yield and Composition of the Essential Oil of a Thyme (<i>Thymus pulegioides</i> L.) Growing Wild in Campania (Southern Italy). Journal of Agricultural and Food Chemistry, 1996, 44, 1327-1332.	2.4	160
5	Chemical Composition and Antimicrobial Activity of the Essential Oils from Three Chemotypes of <i>Origanum vulgare</i> L. ssp. <i>hirtum</i> (Link) Letswaart Growing Wild in Campania (Southern Italy). Molecules, 2009, 14, 2735-2746.	1.7	145
6	Potential allelochemicals from the essential oil of <i>Ruta graveolens</i> . Phytochemistry, 2002, 61, 573-578.	1.4	136
7	Phytochemical composition, anti-inflammatory and antitumour activities of four <i>Teucrium</i> essential oils from Greece. Food Chemistry, 2009, 115, 679-686.	4.2	126
8	Essential oils of <i>Salvia bracteata</i> and <i>Salvia rubifolia</i> from Lebanon: Chemical composition, antimicrobial activity and inhibitory effect on human melanoma cells. Journal of Ethnopharmacology, 2009, 126, 265-272.	2.0	121
9	Supercritical carbon dioxide extraction of chamomile essential oil and its analysis by gas chromatography-mass spectrometry. Journal of Agricultural and Food Chemistry, 1994, 42, 154-158.	2.4	111
10	Chemical Constituents and Biological Activities of <i>Nepeta</i> Species. Chemistry and Biodiversity, 2011, 8, 1783-1818.	1.0	110
11	Medicinal plants and phytotherapy in the Amalfitan Coast, Salerno Province, Campania, Southern Italy. Journal of Ethnopharmacology, 1993, 39, 39-51.	2.0	106
12	Traditional phytotherapy in the Peninsula Sorrentina, Campania, Southern Italy. Journal of Ethnopharmacology, 1992, 36, 113-125.	2.0	98
13	Comparative chemical composition, free radical-scavenging and cytotoxic properties of essential oils of six <i>Stachys</i> species from different regions of the Mediterranean Area. Food Chemistry, 2009, 116, 898-905.	4.2	96
14	Correlation among environmental factors, chemical composition and antioxidative properties of essential oil and extracts of chamomile (<i>Matricaria chamomilla</i> L.) collected in Molise (South-central) Tj ETQq0 0 0 rgBT /Overlock 10 Tf	2.0	96
15	Chemical composition, antimicrobial and antioxidant activities of anethole-rich oil from leaves of selected varieties of fennel [<i>Foeniculum vulgare</i> Mill. ssp. <i>vulgare</i> var. <i>azoricum</i> (Mill.) Thell]. FÅ-toterapÅ-Åç, 2013, 90, 214-219.	1.1	93
16	Chemical Composition and Biological Activity of Essential Oils of <i>Origanum vulgare</i> L. subsp. <i>vulgare</i> L. under Different Growth Conditions. Molecules, 2013, 18, 14948-14960.	1.7	88
17	Antimicrobial and antioxidant properties of the essential oil of <i>Salvia lanigera</i> from Cyprus. Food and Chemical Toxicology, 2011, 49, 238-243.	1.8	82
18	Antibacterial and antioxidant activities in <i>Sideritis italica</i> (Miller) Greuter et Burdet essential oils. Journal of Ethnopharmacology, 2006, 107, 240-248.	2.0	76

#	ARTICLE	IF	CITATIONS
19	Chemistry and functional properties in prevention of neurodegenerative disorders of five <i>Cistus</i> species essential oils. <i>Food and Chemical Toxicology</i> , 2013, 59, 586-594.	1.8	73
20	Chemical Composition and Antimicrobial Activity of the Essential Oils from Two Species of <i>Thymus</i> Growing Wild in Southern Italy. <i>Molecules</i> , 2009, 14, 4614-4624.	1.7	58
21	Antibacterial activity of <i>Tagetes minuta</i> L. (Asteraceae) essential oil with different chemical composition. <i>Flavour and Fragrance Journal</i> , 2004, 19, 574-578.	1.2	56
22	Composition and antibacterial activity of the essential oil from <i>Crithmum maritimum</i> L. (Apiaceae) growing wild in Turkey. <i>Flavour and Fragrance Journal</i> , 2000, 15, 186-189.	1.2	53
23	Influence of harvesting time on composition of the essential oil of <i>Thymus capitatus</i> (L.) Hoffmanns. & Link. growing wild in northern Sicily and its activity on microorganisms affecting historical art crafts. <i>Arabian Journal of Chemistry</i> , 2019, 12, 2704-2712.	2.3	51
24	Volatile components of <i>Centaurea cineraria</i> L. subsp. <i>umbrosa</i> (Lacaita) Pign. and <i>Centaurea napifolia</i> L. (Asteraceae), two species growing wild in Sicily. <i>Flavour and Fragrance Journal</i> , 2003, 18, 248-251.	1.2	50
25	Protection against neurodegenerative diseases of <i>Iris pseudopumila</i> extracts and their constituents. <i>FÄ-toterapÄ-Ä¢</i> , 2009, 80, 62-67.	1.1	50
26	Volatile components of <i>Centaurea eryngioides</i> Lam. and <i>Centaurea iberica</i> Trev. var. <i>hermonis</i> Boiss. Lam., two Asteraceae growing wild in Lebanon. <i>Natural Product Research</i> , 2005, 19, 749-754.	1.0	47
27	Antibacterial and Antifungal Properties of Acetonic Extract of <i>Feijoa sellowiana</i> Fruits and Its Effect on <i>Helicobacter pylori</i> Growth. <i>Journal of Medicinal Food</i> , 2010, 13, 189-195.	0.8	46
28	Flavonoid Glycosides of <i>Barbarea vulgaris</i> L. (Brassicaceae). <i>Journal of Agricultural and Food Chemistry</i> , 2000, 48, 2659-2662.	2.4	44
29	Chemical Composition and Phytotoxic Effects of Essential Oils of <i>Salvia hierosolymitana</i> Boiss. and <i>Salvia multicaulis</i> Vahl. var. <i>simplicifolia</i> Boiss. Growing Wild in Lebanon. <i>Molecules</i> , 2009, 14, 4725-4736.	1.7	44
30	Chemical Composition and Antibacterial Activity of Essential Oils from <i>Thymus spinulosus</i> Ten. (Lamiaceae). <i>Journal of Agricultural and Food Chemistry</i> , 2003, 51, 3849-3853.	2.4	43
31	Effects of solvent-free microwave extraction on the chemical composition of essential oil of <i>Calamintha nepeta</i> (L.) Savi compared with the conventional production method. <i>Journal of Separation Science</i> , 2008, 31, 1110-1117.	1.3	43
32	Flavonoids in Subtribe Centaureinae (Cass.) Dumort. (Tribe Cardueae). <i>Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 2096-2158.</i>	1.0	43
33	Composition, antibacterial, antioxidant and antiproliferative activities of essential oils from three <i>Origanum</i> species growing wild in Lebanon and Greece. <i>Natural Product Research</i> , 2016, 30, 735-739.	1.0	42
34	Composition and antimicrobial activity of the essential oil of <i>Achillea falcata</i> L. (Asteraceae). <i>Flavour and Fragrance Journal</i> , 2005, 20, 291-294.	1.2	41
35	Chemical composition of essential oils and in vitro antioxidant properties of extracts and essential oils of <i>Calamintha origanifolia</i> and <i>Micromeria myrtifolia</i> , two Lamiaceae from the Lebanon flora. <i>Industrial Crops and Products</i> , 2014, 62, 405-411.	2.5	41
36	Chemical composition of the essential oil of <i>Salvia multicaulis</i> Vahl. var. <i>simplicifolia</i> Boiss. growing wild in Lebanon. <i>Journal of Chromatography A</i> , 2004, 1052, 237-240.	1.8	40

#	ARTICLE	IF	CITATIONS
37	Essential oils from two Peruvian <i>Satureja</i> species. <i>Flavour and Fragrance Journal</i> , 1998, 13, 1-4.	1.2	38
38	Essential oil of two <i>Lippia</i> spp. (Verbenaceae) growing wild in Guatemala. <i>Flavour and Fragrance Journal</i> , 2001, 16, 169-171.	1.2	37
39	Phytogrowth-inhibitory and antibacterial activity of <i>Verbascum sinuatum</i> . <i>Fitoterapia</i> , 2007, 78, 244-247.	1.1	37
40	Essential Oils from <i>Salvia</i> spp. (Lamiaceae). I. Chemical Composition of the Essential Oils from <i>Salvia glutinosa</i> L. Growing Wild in Southern Italy. <i>Journal of Essential Oil Research</i> , 1997, 9, 151-157.	1.3	36
41	Comparative phytochemical profile and antiproliferative activity on human melanoma cells of essential oils of three Lebanese <i>Salvia</i> species. <i>Industrial Crops and Products</i> , 2016, 83, 492-499.	2.5	35
42	Inhibition of Inducible Nitric Oxide Synthase Expression by an Acetonic Extract from <i>Feijoa sellowiana</i> Berg. Fruits. <i>Journal of Agricultural and Food Chemistry</i> , 2007, 55, 5053-5061.	2.4	34
43	<i>Genista sessilifolia</i> DC. and <i>Genista tinctoria</i> L. inhibit UV light and nitric oxide-induced DNA damage and human melanoma cell growth. <i>Chemico-Biological Interactions</i> , 2009, 180, 211-219.	1.7	34
44	Volatile constituents of aerial parts of three endemic <i>Centaurea</i> species from Turkey: <i>Centaurea amnicola</i> Hub.-Mor., <i>Centaurea consanguinea</i> DC. and <i>Centaurea ptosimopappa</i> Hayek and their antibacterial activities. <i>Natural Product Research</i> , 2008, 22, 833-839.	1.0	33
45	Characterisation of the essential oil of <i>Nepeta glomerata</i> Montbret et Aucher ex Benth from Lebanon and its biological activities. <i>Natural Product Research</i> , 2011, 25, 614-626.	1.0	32
46	Chemical Composition and Antimicrobial Screening of the Essential Oil of <i>Minthostachys verticillata</i> (Griseb.) Epl. (Lamiaceae). <i>Journal of Essential Oil Research</i> , 1998, 10, 61-65.	1.3	31
47	Volatile components from flower-heads of <i>Centaurea nicaeensis</i> All., <i>C. parlatoris</i> Helder and <i>C. solstitialis</i> L. ssp. <i>schouwii</i> (DC.) Dostal growing wild in southern Italy and their biological activity. <i>Natural Product Research</i> , 2008, 22, 825-832.	1.0	31
48	Antispasmodic Effects and Structure-Activity Relationships of Labdane Diterpenoids from <i>Marrubium globosum</i> ssp. <i>libanoticum</i> . <i>Journal of Natural Products</i> , 2009, 72, 1477-1481.	1.5	31
49	Chemical Composition of Essential Oil from Italian Populations of <i>Artemisia alba</i> Turra (Asteraceae). <i>Molecules</i> , 2012, 17, 10232-10241.	1.7	31
50	Chemical constituents of some basidiomycetes. <i>Journal of the Science of Food and Agriculture</i> , 1988, 45, 337-345.	1.7	30
51	Constituents of <i>Vitex agnus-castus</i> L. Essential Oil. <i>Flavour and Fragrance Journal</i> , 1996, 11, 179-182.	1.2	30
52	Essential Oil Composition of Stems and Fruits of <i>Caralluma europaea</i> N.E.Br. (Apocynaceae). <i>Molecules</i> , 2010, 15, 627-638.	1.7	30
53	Composition of the essential oil from flowerheads of <i>Chrysanthemum coronarium</i> L. (Asteraceae) growing wild in Southern Italy. <i>Flavour and Fragrance Journal</i> , 2004, 19, 149-152.	1.2	29
54	Phytotoxic effects of essential oils of <i>Nepeta curviflora</i> Boiss. and <i>Nepeta nuda</i> L. subsp. <i>albiflora</i> growing wild in Lebanon. <i>Journal of Plant Interactions</i> , 2009, 4, 253-259.	1.0	28

#	ARTICLE	IF	CITATIONS
55	Anthemis wiedemanniana essential oil prevents LPS-induced production of NO in RAW 264.7 macrophages and exerts antiproliferative and antibacterial activities in vitro. Natural Product Research, 2012, 26, 1594-1601.	1.0	28
56	Chemical composition of the essential oil of Salvia microstegia Boiss. et Balansa growing wild in Lebanon. Journal of Chromatography A, 2006, 1108, 276-278.	1.8	27
57	Volatile components of Centaurea calcitrapa L. and Centaurea sphaerocephala L. ssp. sphaerocephala, two Asteraceae growing wild in Sicily. Flavour and Fragrance Journal, 2006, 21, 282-285.	1.2	27
58	Antioxidant Flavonoids and Isoflavonoids from Rhizomes of Iris pseudopumila. Planta Medica, 2007, 73, 93-96.	0.7	27
59	Essential oil of a possible new chemotype of crithmum maritimum L. growing in campania (Southern) Tj ETQq1 1 0.784314 rgBT /Over	1.2	26
60	Vetiver oil production correlates with early root growth. Biochemical Systematics and Ecology, 2006, 34, 376-382.	0.6	26
61	Headspace Volatile Composition of the Flowers of Caralluma europaea N.E.Br. (Apocynaceae). Molecules, 2009, 14, 4597-4613.	1.7	26
62	Growth, essential oil characterization, and antimicrobial activity of three wild biotypes of oregano under cultivation condition in Southern Italy. Industrial Crops and Products, 2014, 62, 242-249.	2.5	26
63	Variation of Malva sylvestris essential oil yield, chemical composition and biological activity in response to different environments across Southern Italy. Industrial Crops and Products, 2017, 98, 29-37.	2.5	26
64	Chemical composition and antimicrobial activity of the essential oil of Phlomis ferruginea Ten. (Lamiaceae) growing wild in Southern Italy. Flavour and Fragrance Journal, 2006, 21, 848-851.	1.2	25
65	Metabolite profile and in vitro activities of Phagnalon saxatile (L.) Cass. relevant to treatment of Alzheimer's disease. Journal of Enzyme Inhibition and Medicinal Chemistry, 2010, 25, 97-104.	2.5	25
66	Intestinal antispasmodic effects of Helichrysum italicum (Roth) Don ssp. italicum and chemical identification of the active ingredients. Journal of Ethnopharmacology, 2013, 150, 901-906.	2.0	25
67	Cytotoxic Activity and Composition of Petroleum Ether Extract from Magydaris tomentosa (Desf.) W. D. J. Koch (Apiaceae). Molecules, 2015, 20, 1571-1578.	1.7	25
68	Chemical composition of essential oils of Senecio nutans Sch.-Bip. (Asteraceae). Flavour and Fragrance Journal, 2003, 18, 234-236.	1.2	24
69	Essential oils from the aerial parts of Centaurea cuneifolia Sibth. & Sm. and C. euxina Velen., two species growing wild in Bulgaria. Biochemical Systematics and Ecology, 2009, 37, 426-431.	0.6	24
70	Labdane Diterpenoids from Marrubium globosum ssp. libanoticum. Journal of Natural Products, 2006, 69, 836-838.	1.5	23
71	Phytochemical and Pharmacological Studies on the Acetonic Extract of Marrubium globosum ssp. libanoticum. Planta Medica, 2006, 72, 575-578.	0.7	22
72	Chemical composition and free radical scavenging activity of the essential oil of Achillea ligustica growing wild in Lipari (Aeolian Islands, Sicily). Natural Product Communications, 2013, 8, 1629-32.	0.2	22

#	ARTICLE	IF	CITATIONS
73	Antibacterial Evaluation of Cnicin and Some Natural and Semisynthetic Analogues. <i>Planta Medica</i> , 2003, 69, 277-281.	0.7	21
74	Composition of the Essential Oil of <i>Nepeta curviflora</i> Boiss. (Lamiaceae) from Lebanon. <i>Journal of Essential Oil Research</i> , 2005, 17, 268-270.	1.3	19
75	Antiproliferative Activity on Human Cancer Cell Lines after Treatment with Polyphenolic Compounds Isolated from <i>Iris pseudopumila</i> Flowers and Rhizomes. <i>Zeitschrift Fur Naturforschung - Section C Journal of Biosciences</i> , 2009, 64, 490-494.	0.6	19
76	Contribution to a Taxonomic Revision of the Sicilian <i>Helichrysum</i> Taxa by PCA Analysis of Their Essential-Oil Compositions. <i>Chemistry and Biodiversity</i> , 2016, 13, 151-159.	1.0	19
77	Chemical composition and antimicrobial activity of the essential oil from aerial parts of <i>Micromeria fruticulosa</i> (Bertol.) Grande (Lamiaceae) growing wild in Southern Italy. <i>Flavour and Fragrance Journal</i> , 2007, 22, 289-292.	1.2	18
78	Volatile constituents of the aerial parts of white salsify (<i>Tragopogon porrifolius</i> L., Asteraceae). <i>Natural Product Research</i> , 2010, 24, 663-668.	1.0	18
79	A study on the essential oil of <i>Ferulago campestris</i> : How much does extraction method influence the oil composition?. <i>Journal of Separation Science</i> , 2011, 34, 483-492.	1.3	18
80	Phytochemical Profile and Apoptotic Activity of <i>Onopordum cynarocephalum</i> . <i>Planta Medica</i> , 2012, 78, 1651-1660.	0.7	18
81	A new acetophenone derivative from flowers of <i>Helichrysum italicum</i> (Roth) Don ssp. <i>italicum</i> . <i>FÄ-toterapÄ-Äç</i> , 2014, 99, 198-203.	1.1	18
82	Essential Oil Variability in a Collection of <i>Ocimum basilicum</i> L. (Basil) Cultivars. <i>Chemistry and Biodiversity</i> , 2016, 13, 1357-1368.	1.0	18
83	Phytochemical profile of three <i>Ballota</i> species essential oils and evaluation of the effects on human cancer cells. <i>Natural Product Research</i> , 2017, 31, 436-444.	1.0	18
84	Fatty acid and free amino acid content of some mushrooms. <i>Journal of the Science of Food and Agriculture</i> , 1990, 51, 91-96.	1.7	17
85	Chemical constituents of some mushrooms. <i>Journal of the Science of Food and Agriculture</i> , 1992, 58, 499-503.	1.7	17
86	Composition of the essential oil of <i>Tagetes filifolia</i> Lag.. <i>Flavour and Fragrance Journal</i> , 1998, 13, 145-147.	1.2	17
87	<i>Eupatorium cannabinum</i> L. ssp. <i>cannabinum</i> (Asteraceae) Essential Oil: Chemical Composition and Antibacterial Activity. <i>Journal of Essential Oil Research</i> , 2001, 13, 463-466.	1.3	17
88	Essential Oils from <i>Salvia</i> sp. (Lamiaceae). III. Composition and Antimicrobial Activity of the Essential Oil of <i>Salvia palaestina</i> Benth. Growing Wild in Lebanon. <i>Journal of Essential Oil Research</i> , 2005, 17, 419-421.	1.3	17
89	Phenolic compounds of <i>Marrubium globosum</i> ssp. <i>libanoticum</i> from Lebanon. <i>Biochemical Systematics and Ecology</i> , 2006, 34, 256-258.	0.6	16
90	Volatile constituents of aerial parts of <i>Centaurea sibthorpii</i> (Sect. <i>Carduiformes</i> , Asteraceae) from Greece and their biological activity. <i>Natural Product Research</i> , 2008, 22, 840-845.	1.0	16

#	ARTICLE	IF	CITATIONS
91	Chemical composition of the essential oil from <i>Tagetes mandonii</i> Sch. Bip. (Asteraceae). <i>Flavour and Fragrance Journal</i> , 1999, 14, 32-34.	1.2	15
92	Composition and allelopathic effect of essential oils of two thistles: <i>Cirsium creticum</i> (Lam.) D. Urv. ssp. <i>triumfetti</i> (Lacaita) Werner and <i>Carduus nutans</i> L.. <i>Journal of Plant Interactions</i> , 2007, 2, 115-120.	1.0	15
93	Composition of the essential oil of <i>Petagnaea gussonei</i> (Sprengel) Rauschert, a relict species from Sicily (Southern Italy). <i>Flavour and Fragrance Journal</i> , 2008, 23, 172-177.	1.2	15
94	Chemical constituents of some species of Agaricaceae. <i>Biochemical Systematics and Ecology</i> , 1988, 16, 601-604.	0.6	14
95	Composition of the Essential Oil of <i>Nepeta betonicifolia</i> C.A. Meyer (Lamiaceae) from Turkey. <i>Journal of Essential Oil Research</i> , 2003, 15, 200-201.	1.3	14
96	Chemical Composition and Antibacterial Activity of Essential Oil of <i>Myrtus communis</i> L. Growing Wild in Italy and Turkey. <i>Journal of Essential Oil-bearing Plants: JEOP</i> , 2006, 9, 162-169.	0.7	14
97	Constituents of Leaves and Flowers Essential Oils of <i>Helichrysum pallasii</i> (Spreng.) Ledeb. Growing Wild in Lebanon. <i>Journal of Medicinal Food</i> , 2009, 12, 203-207.	0.8	14
98	Volatile compounds of flowers and leaves of <i>Sideritis italica</i> (Miller) Greuter et Burdet (Lamiaceae), a plant used as mountain tea. <i>Natural Product Research</i> , 2010, 24, 640-646.	1.0	14
99	Essential oils of three species of <i>Scutellaria</i> and their influence on <i>Spodoptera littoralis</i> . <i>Biochemical Systematics and Ecology</i> , 2013, 48, 206-210.	0.6	13
100	Chemical Composition and Antibacterial Activity of the Essential Oil of a 1,8-Cineole Chemotype of <i>Mentha aquatica</i> L. Growing Wild in Turkey. <i>Journal of Essential Oil-bearing Plants: JEOP</i> , 2005, 8, 148-153.	0.7	12
101	Phytomorphological and Essential Oil Characterization <i>in situ</i> and <i>ex situ</i> of Wild Biotypes of <i>Oregano</i> Collected in the Campania Region (Southern Italy). <i>Chemistry and Biodiversity</i> , 2013, 10, 2078-2090.	1.0	12
102	Chemical Composition of the Essential Oils of Three Endemic Species of <i>Anthemis</i> Sect. <i>Hiorthia</i> (DC.) R.Fern. Growing Wild in Sicily and Chemotaxonomic Volatile Markers of the Genus <i>Anthemis</i> L.: An Update. <i>Chemistry and Biodiversity</i> , 2014, 11, 652-672.	1.0	12
103	Antibacterial and allelopathic activity of methanolic extract from <i>Iris pseudopumila</i> rhizomes. <i>FÄ-toterapÄ-Äç</i> , 2006, 77, 460-462.	1.1	11
104	Chemical composition of essential oils of <i>Anthemis secundiramea</i> Biv. subsp. <i>secundiramea</i> (Asteraceae) collected wild in Sicily and their activity on micro-organisms affecting historical art craft. <i>Natural Product Research</i> , 2019, 33, 970-979.	1.0	11
105	Composition and antibacterial activity of the essential oil of <i>Anisochilus carnosus</i> (Linn. ?!) Benth., a Tamil plant acclimatized in Sicily. <i>Flavour and Fragrance Journal</i> , 2003, 18, 202-204.	1.2	10
106	Chemical Composition of the Essential Oils of <i>Centaurea Sicana</i> and <i>C. Giardiniae</i> Growing Wild in Sicily. <i>Natural Product Communications</i> , 2008, 3, 1934578X0800300.	0.2	10
107	Activity against Microorganisms Affecting Cellulosic Objects of the Volatile Constituents of <i>Leonotis nepetaefolia</i> from Nicaragua. <i>Natural Product Communications</i> , 2014, 9, 1934578X1400901.	0.2	10
108	Chemical composition of the essential oil from <i>Thapsia garganica</i> L. (Apiaceae) grown wild in Sicily and its antimicrobial activity. <i>Natural Product Research</i> , 2016, 30, 1042-1052.	1.0	10

#	ARTICLE	IF	CITATIONS
109	Chemical composition of the essential oil from <i>Pulicaria vulgaris</i> var. <i>graeca</i> (Sch.-Bip.) Fiori (Asteraceae) growing wild in Sicily and its antimicrobial activity. <i>Natural Product Research</i> , 2016, 30, 259-267.	1.0	10
110	Oligosaccharides in Five Different <i>Vicia faba</i> L. Cultivars. <i>Biochemical Systematics and Ecology</i> , 1989, 17, 559-561.	0.6	9
111	Volatile constituents of <i>Minthostachys setosa</i> (Briq.) Epl. (Lamiaceae) from Peru. <i>Flavour and Fragrance Journal</i> , 1998, 13, 263-265.	1.2	9
112	Essential Oils from <i>Salvia</i> spp. (Lamiaceae). II. Chemical Composition of the Essential Oil from <i>Salvia pratensis</i> L. subsp. <i>haematodes</i> (L.) Briq. Inflorescences. <i>Journal of Essential Oil Research</i> , 1998, 10, 135-137.	1.3	9
113	Composition of the essential oil of <i>Pallenis spinosa</i> (L.) Cass. (Asteraceae). <i>Flavour and Fragrance Journal</i> , 2003, 18, 195-197.	1.2	9
114	Chemical Composition of the Essential Oil of <i>Phagnalon saxatile</i> (L.) Cass. (Asteraceae) Growing Wild in Southern Italy. <i>Journal of Essential Oil-bearing Plants: JEOP</i> , 2005, 8, 258-263.	0.7	9
115	Analysis of Essential Oils from <i>Scutellaria orientalis</i> ssp. <i>alpina</i> and <i>S. utriculata</i> by GC and GC-MS. <i>Natural Product Communications</i> , 2011, 6, 1934578X1100600.	0.2	9
116	Chemical composition, antimicrobial and antioxidant activity of the essential oils from <i>Pimpinella</i> <i>tragicum</i> Vill. subsp. <i>glauca</i> (C. Presl.) C. Brullo & Brullo (Apiaceae) growing wild in Sicily. <i>Natural Product Research</i> , 2013, 27, 2338-2346.	1.0	9
117	GC and GC-MS analysis of the essential oil of <i>Nepeta cilicica</i> Boiss. ex Benth. from Lebanon. <i>Natural Product Research</i> , 2013, 27, 1975-1981.	1.0	9
118	Volatile Constituents of the Aerial Parts of <i>Pulicaria sicula</i> (L.) <i>Moris</i> Growing Wild in Sicily: Chemotaxonomic Volatile Markers of the Genus <i>Pulicaria</i> Gaertn. <i>Chemistry and Biodiversity</i> , 2015, 12, 781-799.	1.0	9
119	Composition of the Essential Oil of <i>Allium neapolitanum</i> Cirillo Growing Wild in Sicily and its Activity on Microorganisms Affecting Historical Art Crafts. <i>Journal of Oleo Science</i> , 2015, 64, 1315-1320.	0.6	9
120	Sterols from three <i>Lactarius</i> species. <i>Biochemical Systematics and Ecology</i> , 1981, 9, 247-248.	0.6	8
121	Free amino acids from different cultivars of <i>Vicia faba</i> . <i>Journal of Agricultural and Food Chemistry</i> , 1983, 31, 836-838.	2.4	8
122	Fatty acids, free amino acids and sterols from some species of <i>Stropharia</i> and <i>Stereum</i> . <i>Biochemical Systematics and Ecology</i> , 1990, 18, 103-106.	0.6	8
123	Composition of the essential oil of <i>Santolina neapolitana</i> Jordan et Fourr. <i>Flavour and Fragrance Journal</i> , 1994, 9, 77-79.	1.2	8
124	Composition of the essential oil of <i>Minthostachys spicata</i> (Benth.) Epl. <i>Flavour and Fragrance Journal</i> , 1995, 10, 43-45.	1.2	8
125	Quercetagenin 6-O-β-D-glucopyranoside from <i>Tagetes mandonii</i> . <i>Phytochemistry</i> , 1997, 45, 201-202.	1.4	8
126	Essential Oil Composition of <i>Teucrium divaricatum</i> Sieb. ssp. <i>villosum</i> (Celak.) Rech. fil. Growing Wild in Lebanon. <i>Journal of Medicinal Food</i> , 2010, 13, 1281-1285.	0.8	8

#	ARTICLE	IF	CITATIONS
127	Characterisation and antimicrobial activity of the volatile components of the flowers of <i>Magyaris tomentosa</i> (Desf.) DC. collected in Sicily and Algeria. <i>Natural Product Research</i> , 2014, 28, 1152-1158.	1.0	8
128	Chemical Composition of the Essential Oil of <i>Bupleurum Fontanesii</i> (Apiaceae) Growing Wild in Sicily and its Activity on Microorganisms Affecting Historical Art Crafts. <i>Natural Product Communications</i> , 2016, 11, 1934578X1601100.	0.2	8
129	GC and GC/MS Analysis of the Essential Oil of <i>Salvia hierosolymitana</i> Boiss. Growing Wild in Lebanon. <i>Natural Product Communications</i> , 2007, 2, 1934578X0700200.	0.2	7
130	Essential Oil Composition and Antibacterial Activity of <i>Anthemis mixta</i> and <i>A. TomENTOSA</i> (Asteraceae). <i>Natural Product Communications</i> , 2012, 7, 1934578X1200701.	0.2	7
131	Two new quercetagenin O-glucosides from <i>Tagetes mandonii</i> . <i>Biochemical Systematics and Ecology</i> , 1999, 27, 309-311.	0.6	6
132	4-hydroxybenzyl glucosinolate from <i>Cardaria draba</i> (Cruciferae). <i>Biochemical Systematics and Ecology</i> , 2003, 31, 1205-1207.	0.6	6
133	Chemical composition and antibacterial activity of essential oils from five culinary herbs of the Lamiaceae family growing in Campania, Southern Italy. <i>Journal of Essential Oil-bearing Plants: JEOP</i> , 2003, 6, 166-173.	0.7	6
134	Essential oil composition and antifeedant properties of <i>Bellardia trixago</i> (L.) All. (sin. <i>Bartsia trixago</i>) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 5	0.6	6
135	<i>Daphne oleoides</i> : An alternative source of important sesquiterpenes. <i>International Journal of Food Properties</i> , 2017, 20, 549-559.	1.3	6
136	Sterols, fatty acids and free amino acids from two <i>Helvella</i> species. <i>Biochemical Systematics and Ecology</i> , 1982, 10, 285-287.	0.6	5
137	Essential oil of <i>Eremocharis triradiata</i> (Wolff.) Johnston (Apiaceae) growing wild in PerÃ©. <i>Flavour and Fragrance Journal</i> , 1997, 12, 257-259.	1.2	5
138	Volatile Constituents of <i>Calamintha organifolia</i> Boiss. Growing Wild in Lebanon. <i>Natural Product Communications</i> , 2007, 2, 1934578X0700201.	0.2	5
139	Essential Oil Composition of <i>Tanacetum vulgare</i> Subsp. <i>Siculum</i> (Guss.) Raimondo et Spadaro (Asteraceae) from Sicily. <i>Natural Product Communications</i> , 2009, 4, 1934578X0900400.	0.2	5
140	Chemical Composition of the Essential Oils of <i>Centaurea formanekii</i> and <i>C. orphanidea</i> ssp. <i>thessala</i> , Growing Wild in Greece. <i>Natural Product Communications</i> , 2012, 7, 1934578X1200700.	0.2	5
141	Chemical Composition of the Essential Oils of Three Species of Apiaceae Growing Wild in Sicily: <i>Bonannia graeca</i> , <i>Eryngium maritimum</i> and <i>Opopanax chironium</i> . <i>Natural Product Communications</i> , 2013, 8, 1934578X1300800.	0.2	5
142	Chemical Composition and Antimicrobial Activity of the Essential Oil from Flowers of <i>Eryngium triquetrum</i> (Apiaceae) Collected Wild in Sicily. <i>Natural Product Communications</i> , 2016, 11, 1934578X1601100.	0.2	5
143	<i>Carica candicans</i> Gray (Mito), an Alimentary Resource from Peruvian Flora. <i>Journal of Agricultural and Food Chemistry</i> , 1999, 47, 3682-3684.	2.4	4
144	Chemical Composition and Antibacterial Activity of Extracts of <i>Helleborus bocconeii</i> Ten. subsp. <i>intermedius</i> . <i>Natural Product Communications</i> , 2007, 2, 1934578X0700200.	0.2	4

#	ARTICLE	IF	CITATIONS
145	Antibacterial Activity and Composition of the Essential Oil of <i>Peperomia Galioides</i> HBK (Piperaceae) from Peru. <i>Natural Product Communications</i> , 2008, 3, 1934578X0800300.	0.2	4
146	Chemical Composition and Free Radical Scavenging Activity of the Essential Oil of <i>Achillea ligustica</i> Growing Wild in Lipari (Aeolian Islands, Sicily). <i>Natural Product Communications</i> , 2013, 8, 1934578X1300801.	0.2	4
147	Chemical composition of the essential oil from the aerial parts of <i>Ononis reclinata</i> L. (Fabaceae) grown wild in Sicily. <i>Natural Product Research</i> , 2017, 31, 7-15.	1.0	4
148	Volatile Components of Aerial Parts of <i>Centaurea nigrescens</i> and <i>C. stenolepis</i> Growing Wild in the Balkans. <i>Natural Product Communications</i> , 2010, 5, 1934578X1000500.	0.2	3
149	Antiproliferative and Cytotoxic Effects on Malignant Melanoma Cells of Essential Oils from the Aerial Parts of <i>Genista sessilifolia</i> and <i>G. tinctoria</i> . <i>Natural Product Communications</i> , 2010, 5, 1934578X1000500.	0.2	3
150	Comparative Chemical Composition and Antioxidant Properties of the Essential Oils of three <i>Sideritis libanotica</i> Subspecies. <i>Natural Product Communications</i> , 2015, 10, 1934578X1501000.	0.2	3
151	Essential oil from aerial parts of <i>Vernonia colorata</i> Drake and <i>Vernonia nigritiana</i> Oliver et Hiern. (Asteraceae) growing wild in Mali. <i>Journal of Essential Oil-bearing Plants: JEOP</i> , 2004, 7, 267-274.	0.7	2
152	Volatile Components of <i>Centaurea Bracteata</i> and <i>C. Pannonica</i> subsp. <i>Pannonica</i> growing wild in Croatia. <i>Natural Product Communications</i> , 2010, 5, 1934578X1000501.	0.2	2
153	Cytotoxic Properties of <i>Marrubium globosum</i> ssp. <i>libanoticum</i> and its Bioactive Components. <i>Natural Product Communications</i> , 2013, 8, 1934578X1300800.	0.2	2
154	Chemical Composition of the Essential Oil of the Local Endemics <i>Centaurea davidovii</i> and <i>C. parilica</i> (Asteraceae, sect. <i>Lepteranthus</i>) from Bulgaria. <i>Natural Product Communications</i> , 2014, 9, 1934578X1400900.	0.2	2
155	Volatile constituents of <i>Stachys palaestina</i> L. (Palestine woundwort) growing in Lebanon. <i>Natural Product Research</i> , 2014, 28, 1674-1679.	1.0	2
156	Effect of Three <i>Centaurea</i> Species Collected from Central Anatolia Region of Turkey on Human Melanoma Cells. <i>Natural Product Communications</i> , 2016, 11, 1934578X1601100.	0.2	1
157	Chemical Composition of the Essential Oil of <i>Mentha pulegium</i> Growing Wild in Sicily and its Activity on Microorganisms Affecting Historical Art Crafts. <i>Natural Product Communications</i> , 2017, 12, 1934578X1701200.	0.2	1
158	Volatile Components from Aerial parts of <i>Centaurea gracilentia</i> and <i>C. ovina</i> ssp. <i>besserana</i> Growing Wild in Bulgaria. <i>Natural Product Communications</i> , 2011, 6, 1934578X1100600.	0.2	0