

Felice Senatore

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8995654/publications.pdf>

Version: 2024-02-01

158
papers

5,156
citations

94269

37
h-index

114278

63
g-index

160
all docs

160
docs citations

160
times ranked

5677
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 1 | Influence of harvesting time on composition of the essential oil of <i>Thymus capitatus</i> (L.) Hoffmanns. & Link. growing wild in northern Sicily and its activity on microorganisms affecting historical art crafts. <i>Arabian Journal of Chemistry</i> , 2019, 12, 2704-2712. | 2.3 | 51 |
| 2 | Chemical composition of essential oils of <i>Anthemis secundiramea</i> Biv. subsp. <i>secundiramea</i> (Asteraceae) collected wild in Sicily and their activity on micro-organisms affecting historical art craft. <i>Natural Product Research</i> , 2019, 33, 970-979. | 1.0 | 11 |
| 3 | <i>Daphne oleoides</i> : An alternative source of important sesquiterpenes. <i>International Journal of Food Properties</i> , 2017, 20, 549-559. | 1.3 | 6 |
| 4 | Phytochemical profile of three <i>Ballota</i> species essential oils and evaluation of the effects on human cancer cells. <i>Natural Product Research</i> , 2017, 31, 436-444. | 1.0 | 18 |
| 5 | Chemical composition of the essential oil from the aerial parts of <i>Ononis reclinata</i> L. (Fabaceae) grown wild in Sicily. <i>Natural Product Research</i> , 2017, 31, 7-15. | 1.0 | 4 |
| 6 | Variation of <i>Malva sylvestris</i> essential oil yield, chemical composition and biological activity in response to different environments across Southern Italy. <i>Industrial Crops and Products</i> , 2017, 98, 29-37. | 2.5 | 26 |
| 7 | Chemical Composition of the Essential Oil of <i>Mentha pulegium</i> Growing Wild in Sicily and its Activity on Microorganisms Affecting Historical Art Crafts. <i>Natural Product Communications</i> , 2017, 12, 1934578X1701200. | 0.2 | 1 |
| 8 | Chemical Composition and Antimicrobial Activity of the Essential Oil from Flowers of <i>Eryngium triquetrum</i> (Apiaceae) Collected Wild in Sicily. <i>Natural Product Communications</i> , 2016, 11, 1934578X1601100. | 0.2 | 5 |
| 9 | Effect of Three <i>Centaurea</i> Species Collected from Central Anatolia Region of Turkey on Human Melanoma Cells. <i>Natural Product Communications</i> , 2016, 11, 1934578X1601100. | 0.2 | 1 |
| 10 | Chemical Composition of the Essential Oil of <i>Bupleurum Fontanesii</i> (Apiaceae) Growing Wild in Sicily and its Activity on Microorganisms Affecting Historical Art Crafts. <i>Natural Product Communications</i> , 2016, 11, 1934578X1601100. | 0.2 | 8 |
| 11 | Contribution to a Taxonomic Revision of the Sicilian <i>Helichrysum</i> Taxa by PCA Analysis of Their Essential-Oil Compositions. <i>Chemistry and Biodiversity</i> , 2016, 13, 151-159. | 1.0 | 19 |
| 12 | Essential Oil Variability in a Collection of <i>Ocimum basilicum</i> L. (Basil) Cultivars. <i>Chemistry and Biodiversity</i> , 2016, 13, 1357-1368. | 1.0 | 18 |
| 13 | Comparative phytochemical profile and antiproliferative activity on human melanoma cells of essential oils of three lebanese <i>Salvia</i> species. <i>Industrial Crops and Products</i> , 2016, 83, 492-499. | 2.5 | 35 |
| 14 | Chemical composition of the essential oil from <i>Thapsia garganica</i> L. (Apiaceae) grown wild in Sicily and its antimicrobial activity. <i>Natural Product Research</i> , 2016, 30, 1042-1052. | 1.0 | 10 |
| 15 | Chemical composition of the essential oil from <i>Pulicaria vulgaris</i> var. <i>graeca</i> (Sch.-Bip.) Fiori (Asteraceae) growing wild in Sicily and its antimicrobial activity. <i>Natural Product Research</i> , 2016, 30, 259-267. | 1.0 | 10 |
| 16 | Composition, antibacterial, antioxidant and antiproliferative activities of essential oils from three <i>Origanum</i> species growing wild in Lebanon and Greece. <i>Natural Product Research</i> , 2016, 30, 735-739. | 1.0 | 42 |
| 17 | Cytotoxic Activity and Composition of Petroleum Ether Extract from <i>Magydaris tomentosa</i> (Desf.) W. D. J. Koch (Apiaceae). <i>Molecules</i> , 2015, 20, 1571-1578. | 1.7 | 25 |
| 18 | Volatile Constituents of the Aerial Parts of <i>Pulicaria sicula</i> (L.) <i>Moris</i> Growing Wild in Sicily: Chemotaxonomic Volatile Markers of the Genus <i>Pulicaria</i> <i>Gaertn</i> .. <i>Chemistry and Biodiversity</i> , 2015, 12, 781-799. | 1.0 | 9 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 19 | Comparative Chemical Composition and Antioxidant Properties of the Essential Oils of three <i>Sideritis libanotica</i> Subspecies. <i>Natural Product Communications</i> , 2015, 10, 1934578X1501000. | 0.2 | 3 |
| 20 | Composition of the Essential Oil of <i>Allium neapolitanum</i> ; Cirillo Growing Wild in Sicily and its Activity on Microorganisms Affecting Historical Art Crafts. <i>Journal of Oleo Science</i> , 2015, 64, 1315-1320. | 0.6 | 9 |
| 21 | Correlation among environmental factors, chemical composition and antioxidative properties of essential oil and extracts of chamomile (<i>Matricaria chamomilla</i> L.) collected in Molise (South-central Italy). <i>Journal of Essential Oil Research</i> , 2015, 27, 113-119. | 0.2 | 1 |
| 22 | Activity against Microorganisms Affecting Cellulosic Objects of the Volatile Constituents of <i>Leonotis nepetaefolia</i> from Nicaragua. <i>Natural Product Communications</i> , 2014, 9, 1934578X1400901. | 0.2 | 10 |
| 23 | Chemical Composition of the Essential Oil of the Local Endemics <i>Centaurea davidovii</i> and <i>C. parilica</i> (Asteraceae, sect. <i>Lepteranthus</i>) from Bulgaria. <i>Natural Product Communications</i> , 2014, 9, 1934578X1400900. | 0.2 | 2 |
| 24 | Chemical Composition of the Essential Oils of Three Endemic Species of <i>Anthemis</i> Sect. <i>Hiorthia</i> (DC.) R.Fern. Growing Wild in Sicily and Chemotaxonomic Volatile Markers of the Genus <i>Anthemis</i> L.: An Update. <i>Chemistry and Biodiversity</i> , 2014, 11, 652-672. | 1.0 | 12 |
| 25 | Characterisation and antimicrobial activity of the volatile components of the flowers of <i>Magyaris tomentosa</i> (Desf.) DC. collected in Sicily and Algeria. <i>Natural Product Research</i> , 2014, 28, 1152-1158. | 1.0 | 8 |
| 26 | Volatile constituents of <i>Stachys palaestina</i> L. (Palestine woundwort) growing in Lebanon. <i>Natural Product Research</i> , 2014, 28, 1674-1679. | 1.0 | 2 |
| 27 | Growth, essential oil characterization, and antimicrobial activity of three wild biotypes of oregano under cultivation condition in Southern Italy. <i>Industrial Crops and Products</i> , 2014, 62, 242-249. | 2.5 | 26 |
| 28 | A new acetophenone derivative from flowers of <i>Helichrysum italicum</i> (Roth) Don ssp. <i>italicum</i> . <i>Fitoterapia</i> , 2014, 99, 198-203. | 1.1 | 18 |
| 29 | Chemical composition of essential oils and in vitro antioxidant properties of extracts and essential oils of <i>Calamintha origanifolia</i> and <i>Micromeria myrtifolia</i> , two Lamiaceae from the Lebanon flora. <i>Industrial Crops and Products</i> , 2014, 62, 405-411. | 2.5 | 41 |
| 30 | Chemistry and functional properties in prevention of neurodegenerative disorders of five <i>Cistus</i> species essential oils. <i>Food and Chemical Toxicology</i> , 2013, 59, 586-594. | 1.8 | 73 |
| 31 | Intestinal antispasmodic effects of <i>Helichrysum italicum</i> (Roth) Don ssp. <i>italicum</i> and chemical identification of the active ingredients. <i>Journal of Ethnopharmacology</i> , 2013, 150, 901-906. | 2.0 | 25 |
| 32 | Chemical composition, antimicrobial and antioxidant activity of the essential oils from <i>Pimpinella tragioides</i> Vill. subsp. <i>glauca</i> (C. Presl.) C. Brullo & Brullo (Apiaceae) growing wild in Sicily. <i>Natural Product Research</i> , 2013, 27, 2338-2346. | 1.0 | 9 |
| 33 | GC and GC-MS analysis of the essential oil of <i>Nepeta cilicica</i> Boiss. ex Benth. from Lebanon. <i>Natural Product Research</i> , 2013, 27, 1975-1981. | 1.0 | 9 |
| 34 | Chemical composition and anticancer activity of essential oils of Mediterranean sage (<i>Salvia officinalis</i> L.) growing wild in Sicily. <i>Journal of Essential Oil Research</i> , 2013, 25, 42-47. | 1.8 | 172 |
| 35 | Chemical composition, antimicrobial and antioxidant activities of anethole-rich oil from leaves of selected varieties of fennel [<i>Foeniculum vulgare</i> Mill. ssp. <i>vulgare</i> var. <i>azoricum</i> (Mill.) Thell.]. <i>Fitoterapia</i> , 2013, 90, 214-219. | 1.1 | 93 |
| 36 | Essential oils of three species of <i>Scutellaria</i> and their influence on <i>Spodoptera littoralis</i> . <i>Biochemical Systematics and Ecology</i> , 2013, 48, 206-210. | 0.6 | 13 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 37 | Phytomorphological and Essential Oil Characterization <i>in situ</i> and <i>ex situ</i> of Wild Biotypes of Oregano Collected in the Campania Region (Southern Italy). <i>Chemistry and Biodiversity</i> , 2013, 10, 2078-2090. | 1.0 | 12 |
| 38 | Chemical Composition and Biological Activity of Essential Oils of <i>Origanum vulgare</i> L. subsp. <i>vulgare</i> L. under Different Growth Conditions. <i>Molecules</i> , 2013, 18, 14948-14960. | 1.7 | 88 |
| 39 | Chemical Composition of the Essential Oils of Three Species of Apiaceae Growing Wild in Sicily: <i>Bonannia graeca</i> , <i>Eryngium maritimum</i> and <i>Opopanax chironium</i> . <i>Natural Product Communications</i> , 2013, 8, 1934578X1300800. | 0.2 | 5 |
| 40 | Cytotoxic Properties of <i>Marrubium globosum</i> ssp. <i>libanoticum</i> and its Bioactive Components. <i>Natural Product Communications</i> , 2013, 8, 1934578X1300800. | 0.2 | 2 |
| 41 | Chemical Composition and Free Radical Scavenging Activity of the Essential Oil of <i>Achillea ligustica</i> Growing Wild in Lipari (Aeolian Islands, Sicily). <i>Natural Product Communications</i> , 2013, 8, 1934578X1300801. | 0.2 | 4 |
| 42 | Chemical composition and free radical scavenging activity of the essential oil of <i>Achillea ligustica</i> growing wild in Lipari (Aeolian Islands, Sicily). <i>Natural Product Communications</i> , 2013, 8, 1629-32. | 0.2 | 22 |
| 43 | <i>Anthemis wiedemanniana</i> essential oil prevents LPS-induced production of NO in RAW 264.7 macrophages and exerts antiproliferative and antibacterial activities <i>in vitro</i> . <i>Natural Product Research</i> , 2012, 26, 1594-1601. | 1.0 | 28 |
| 44 | Phytochemical Profile and Apoptotic Activity of <i>Onopordum cynarocephalum</i> . <i>Planta Medica</i> , 2012, 78, 1651-1660. | 0.7 | 18 |
| 45 | Chemical Composition of Essential Oil from Italian Populations of <i>Artemisia alba</i> Turra (Asteraceae). <i>Molecules</i> , 2012, 17, 10232-10241. | 1.7 | 31 |
| 46 | Flavonoids in Subtribe Centaureinae (Cass.) Dumort. (Tribe Cardueae). <i>Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 3</i> 2096-2158. | 1.0 | 43 |
| 47 | Chemical Composition of the Essential Oils of <i>Centaurea formanekii</i> and <i>C. orphanidea</i> ssp. <i>thessala</i> , Growing Wild in Greece. <i>Natural Product Communications</i> , 2012, 7, 1934578X1200700. | 0.2 | 5 |
| 48 | Essential Oil Composition and Antibacterial Activity of <i>Anthemis mixta</i> and <i>A. Tomentosa</i> (Asteraceae). <i>Natural Product Communications</i> , 2012, 7, 1934578X1200701. | 0.2 | 7 |
| 49 | Characterisation of the essential oil of <i>Nepeta glomerata</i> Montbret et Aucher ex Benth from Lebanon and its biological activities. <i>Natural Product Research</i> , 2011, 25, 614-626. | 1.0 | 32 |
| 50 | Antimicrobial and antioxidant properties of the essential oil of <i>Salvia lanigera</i> from Cyprus. <i>Food and Chemical Toxicology</i> , 2011, 49, 238-243. | 1.8 | 82 |
| 51 | Volatile Components from Aerial parts of <i>Centaurea gracilentia</i> and <i>C. ovina</i> ssp. <i>besserana</i> Growing Wild in Bulgaria. <i>Natural Product Communications</i> , 2011, 6, 1934578X1100600. | 0.2 | 0 |
| 52 | Analysis of Essential Oils from <i>Scutellaria orientalis</i> ssp. <i>alpina</i> and <i>S. utriculata</i> by GC and GC-MS. <i>Natural Product Communications</i> , 2011, 6, 1934578X1100600. | 0.2 | 9 |
| 53 | A study on the essential oil of <i>Ferulago campestris</i> : How much does extraction method influence the oil composition?. <i>Journal of Separation Science</i> , 2011, 34, 483-492. | 1.3 | 18 |
| 54 | Chemical Constituents and Biological Activities of <i>Nepeta</i> Species. <i>Chemistry and Biodiversity</i> , 2011, 8, 1783-1818. | 1.0 | 110 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 55 | Volatile Components of Aerial Parts of <i>Centaurea nigrescens</i> and <i>C. stenolepis</i> Growing Wild in the Balkans. <i>Natural Product Communications</i> , 2010, 5, 1934578X1000500. | 0.2 | 3 |
| 56 | Antiproliferative and Cytotoxic Effects on Malignant Melanoma Cells of Essential Oils from the Aerial Parts of <i>Genista sessilifolia</i> and <i>G. tinctoria</i> . <i>Natural Product Communications</i> , 2010, 5, 1934578X1000500. | 0.2 | 3 |
| 57 | Volatile Components of <i>Centaurea Bracteata</i> and <i>C. Pannonica</i> subsp. <i>Pannonica</i> growing wild in Croatia. <i>Natural Product Communications</i> , 2010, 5, 1934578X1000501. | 0.2 | 2 |
| 58 | Volatile constituents of the aerial parts of white salsify (<i>Tragopogon porrifolius</i> L., Asteraceae). <i>Natural Product Research</i> , 2010, 24, 663-668. | 1.0 | 18 |
| 59 | Essential Oil Composition of Stems and Fruits of <i>Caralluma europaea</i> N.E.Br. (Apocynaceae). <i>Molecules</i> , 2010, 15, 627-638. | 1.7 | 30 |
| 60 | Essential Oil Composition of <i>Teucrium divaricatum</i> Sieb. ssp. <i>villosum</i> (Celak.) Rech. fil. Growing Wild in Lebanon. <i>Journal of Medicinal Food</i> , 2010, 13, 1281-1285. | 0.8 | 8 |
| 61 | Antibacterial and Antifungal Properties of Acetonic Extract of <i>Feijoa sellowiana</i> Fruits and Its Effect on <i>Helicobacter pylori</i> Growth. <i>Journal of Medicinal Food</i> , 2010, 13, 189-195. | 0.8 | 46 |
| 62 | Volatile compounds of flowers and leaves of <i>Sideritis italica</i> (Miller) Greuter et Burdet (Lamiaceae), a plant used as mountain tea. <i>Natural Product Research</i> , 2010, 24, 640-646. | 1.0 | 14 |
| 63 | Metabolite profile and <i>in vitro</i> activities of <i>Phagnalon saxatile</i> (L.) Cass. relevant to treatment of Alzheimer's disease. <i>Journal of Enzyme Inhibition and Medicinal Chemistry</i> , 2010, 25, 97-104. | 2.5 | 25 |
| 64 | Essential Oil Composition of <i>Tanacetum vulgare</i> Subsp. <i>Siculum</i> (Guss.) Raimondo et Spadaro (Asteraceae) from Sicily. <i>Natural Product Communications</i> , 2009, 4, 1934578X0900400. | 0.2 | 5 |
| 65 | Chemical Composition and Antimicrobial Activity of the Essential Oils from Two Species of <i>Thymus</i> Growing Wild in Southern Italy. <i>Molecules</i> , 2009, 14, 4614-4624. | 1.7 | 58 |
| 66 | Chemical Composition and Phytotoxic Effects of Essential Oils of <i>Salvia hierosolymitana</i> Boiss. and <i>Salvia multicaulis</i> Vahl. var. <i>simplicifolia</i> Boiss. Growing Wild in Lebanon. <i>Molecules</i> , 2009, 14, 4725-4736. | 1.7 | 44 |
| 67 | Antiproliferative Activity on Human Cancer Cell Lines after Treatment with Polyphenolic Compounds Isolated from <i>Iris pseudopumila</i> Flowers and Rhizomes. <i>Zeitschrift Fur Naturforschung - Section C Journal of Biosciences</i> , 2009, 64, 490-494. | 0.6 | 19 |
| 68 | Phytotoxic effects of essential oils of <i>Nepeta curviflora</i> Boiss. and <i>Nepeta nuda</i> L. subsp. <i>albiflora</i> growing wild in Lebanon. <i>Journal of Plant Interactions</i> , 2009, 4, 253-259. | 1.0 | 28 |
| 69 | Protection against neurodegenerative diseases of <i>Iris pseudopumila</i> extracts and their constituents. <i>FAA-toterapA-Å</i> , 2009, 80, 62-67. | 1.1 | 50 |
| 70 | Essential oils from the aerial parts of <i>Centaurea cuneifolia</i> Sibth. & Sm. and <i>C. euxina</i> Velen., two species growing wild in Bulgaria. <i>Biochemical Systematics and Ecology</i> , 2009, 37, 426-431. | 0.6 | 24 |
| 71 | <i>Genista sessilifolia</i> DC. and <i>Genista tinctoria</i> L. inhibit UV light and nitric oxide-induced DNA damage and human melanoma cell growth. <i>Chemico-Biological Interactions</i> , 2009, 180, 211-219. | 1.7 | 34 |
| 72 | Phytochemical composition, anti-inflammatory and antitumour activities of four <i>Teucrium</i> essential oils from Greece. <i>Food Chemistry</i> , 2009, 115, 679-686. | 4.2 | 126 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 73 | Comparative chemical composition, free radical-scavenging and cytotoxic properties of essential oils of six <i>Stachys</i> species from different regions of the Mediterranean Area. <i>Food Chemistry</i> , 2009, 116, 898-905. | 4.2 | 96 |
| 74 | Antispasmodic Effects and Structure-Activity Relationships of Labdane Diterpenoids from <i>Marrubium globosum</i> ssp. <i>libanoticum</i> . <i>Journal of Natural Products</i> , 2009, 72, 1477-1481. | 1.5 | 31 |
| 75 | Essential oils of <i>Salvia bracteata</i> and <i>Salvia rubifolia</i> from Lebanon: Chemical composition, antimicrobial activity and inhibitory effect on human melanoma cells. <i>Journal of Ethnopharmacology</i> , 2009, 126, 265-272. | 2.0 | 121 |
| 76 | Chemical Composition and Antimicrobial Activity of the Essential Oils from Three Chemotypes of <i>Origanum vulgare</i> L. ssp. <i>hirtum</i> (Link) Letswaart Growing Wild in Campania (Southern Italy). <i>Molecules</i> , 2009, 14, 2735-2746. | 1.7 | 145 |
| 77 | Constituents of Leaves and Flowers Essential Oils of <i>Helichrysum pallasii</i> (Spreng.) Ledeb. Growing Wild in Lebanon. <i>Journal of Medicinal Food</i> , 2009, 12, 203-207. | 0.8 | 14 |
| 78 | Headspace Volatile Composition of the Flowers of <i>Caralluma europaea</i> N.E.Br. (Apocynaceae). <i>Molecules</i> , 2009, 14, 4597-4613. | 1.7 | 26 |
| 79 | Effects of solvent-free microwave extraction on the chemical composition of essential oil of <i>Calamintha nepeta</i> (L.) Savi compared with the conventional production method. <i>Journal of Separation Science</i> , 2008, 31, 1110-1117. | 1.3 | 43 |
| 80 | Composition of the essential oil of <i>Petagnaea gussonei</i> (Sprengel) Rauschert, a relict species from Sicily (Southern Italy). <i>Flavour and Fragrance Journal</i> , 2008, 23, 172-177. | 1.2 | 15 |
| 81 | Essential oil composition and antifeedant properties of <i>Bellardia trixago</i> (L.) All. (sin. <i>Bartsia trixago</i>) Tj ETQq1 1 0.784314 rgBT /Overl 0.6 | 0.6 | 6 |
| 82 | Volatile constituents of aerial parts of three endemic <i>Centaurea</i> species from Turkey: <i>Centaurea amanicola</i> Hub.-Mor., <i>Centaurea consanguinea</i> DC. and <i>Centaurea ptosimopappa</i> Hayek and their antibacterial activities. <i>Natural Product Research</i> , 2008, 22, 833-839. | 1.0 | 33 |
| 83 | Volatile components from flower-heads of <i>Centaurea nicaeensis</i> All., <i>C. parlatoris</i> Helder and <i>C. solstitialis</i> L. ssp. <i>schouwii</i> (DC.) Dostal growing wild in southern Italy and their biological activity. <i>Natural Product Research</i> , 2008, 22, 825-832. | 1.0 | 31 |
| 84 | Volatile constituents of aerial parts of <i>Centaurea sibthorpii</i> (Sect. <i>Carduiformes</i> , Asteraceae) from Greece and their biological activity. <i>Natural Product Research</i> , 2008, 22, 840-845. | 1.0 | 16 |
| 85 | Chemical Composition of the Essential Oils of <i>Centaurea Sicana</i> and <i>C. Giardinae</i> Growing Wild in Sicily. <i>Natural Product Communications</i> , 2008, 3, 1934578X0800300. | 0.2 | 10 |
| 86 | Antibacterial Activity and Composition of the Essential Oil of <i>Peperomia galioides</i> HBK (Piperaceae) from Peru. <i>Natural Product Communications</i> , 2008, 3, 1934578X0800300. | 0.2 | 4 |
| 87 | Antioxidant Flavonoids and Isoflavonoids from Rhizomes of <i>Iris pseudopumila</i> . <i>Planta Medica</i> , 2007, 73, 93-96. | 0.7 | 27 |
| 88 | Composition and allelopathic effect of essential oils of two thistles: <i>Cirsium creticum</i> (Lam.) D.'Urv. ssp. <i>triumfetti</i> (Lacaita) Werner and <i>Carduus nutans</i> L.. <i>Journal of Plant Interactions</i> , 2007, 2, 115-120. | 1.0 | 15 |
| 89 | Inhibition of Inducible Nitric Oxide Synthase Expression by an Acetonic Extract from <i>Feijoa sellowiana</i> Berg. Fruits. <i>Journal of Agricultural and Food Chemistry</i> , 2007, 55, 5053-5061. | 2.4 | 34 |
| 90 | Volatile Constituents of <i>Calamintha organifolia</i> Boiss. Growing Wild in Lebanon. <i>Natural Product Communications</i> , 2007, 2, 1934578X0700201. | 0.2 | 5 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 91 | Chemical Composition and Antibacterial Activity of Extracts of <i>Helleborus bocconeii</i> Ten. subsp. <i>intermedius</i> . <i>Natural Product Communications</i> , 2007, 2, 1934578X0700200. | 0.2 | 4 |
| 92 | GC and GC/MS Analysis of the Essential Oil of <i>Salvia hierosolymitana</i> Boiss. Growing Wild in Lebanon. <i>Natural Product Communications</i> , 2007, 2, 1934578X0700200. | 0.2 | 7 |
| 93 | Chemical composition and antimicrobial activity of the essential oil from aerial parts of <i>Micromeria fruticulosa</i> (Bertol.) Grande (Lamiaceae) growing wild in Southern Italy. <i>Flavour and Fragrance Journal</i> , 2007, 22, 289-292. | 1.2 | 18 |
| 94 | Phytogrowth-inhibitory and antibacterial activity of <i>Verbascum sinuatum</i> . <i>FÄ-toterapÄ-Ät</i> , 2007, 78, 244-247. | 1.1 | 37 |
| 95 | Chemical Composition and Antibacterial Activity of Essential Oil of <i>Myrtus communis</i> L. Growing Wild in Italy and Turkey. <i>Journal of Essential Oil-bearing Plants: JEOP</i> , 2006, 9, 162-169. | 0.7 | 14 |
| 96 | Labdane Diterpenoids from <i>Marrubium globosum</i> ssp. <i>libanoticum</i> . <i>Journal of Natural Products</i> , 2006, 69, 836-838. | 1.5 | 23 |
| 97 | Antibacterial and antioxidant activities in <i>Sideritis italica</i> (Miller) Greuter et Burdet essential oils. <i>Journal of Ethnopharmacology</i> , 2006, 107, 240-248. | 2.0 | 76 |
| 98 | Chemical composition of the essential oil of <i>Salvia microstegia</i> Boiss. et Balansa growing wild in Lebanon. <i>Journal of Chromatography A</i> , 2006, 1108, 276-278. | 1.8 | 27 |
| 99 | Antibacterial and allelopathic activity of methanolic extract from <i>Iris pseudopumila</i> rhizomes. <i>FÄ-toterapÄ-Ät</i> , 2006, 77, 460-462. | 1.1 | 11 |
| 100 | Phenolic compounds of <i>Marrubium globosum</i> ssp. <i>libanoticum</i> from Lebanon. <i>Biochemical Systematics and Ecology</i> , 2006, 34, 256-258. | 0.6 | 16 |
| 101 | Vetiver oil production correlates with early root growth. <i>Biochemical Systematics and Ecology</i> , 2006, 34, 376-382. | 0.6 | 26 |
| 102 | Volatile components of <i>Centaurea calcitrapa</i> L. and <i>Centaurea sphaerocephala</i> L. ssp. <i>sphaerocephala</i> , two Asteraceae growing wild in Sicily. <i>Flavour and Fragrance Journal</i> , 2006, 21, 282-285. | 1.2 | 27 |
| 103 | Chemical composition and antimicrobial activity of the essential oil of <i>Phlomis ferruginea</i> Ten. (Lamiaceae) growing wild in Southern Italy. <i>Flavour and Fragrance Journal</i> , 2006, 21, 848-851. | 1.2 | 25 |
| 104 | Phytochemical and Pharmacological Studies on the Acetonic Extract of <i>Marrubium globosum</i> ssp. <i>libanoticum</i> . <i>Planta Medica</i> , 2006, 72, 575-578. | 0.7 | 22 |
| 105 | Composition and antimicrobial activity of the essential oil of <i>Achillea falcata</i> L. (Asteraceae). <i>Flavour and Fragrance Journal</i> , 2005, 20, 291-294. | 1.2 | 41 |
| 106 | Volatile components of <i>Centaurea eryngioides</i> Lam. and <i>Centaurea iberica</i> Trev. var. <i>hermonis</i> Boiss. Lam., two Asteraceae growing wild in Lebanon. <i>Natural Product Research</i> , 2005, 19, 749-754. | 1.0 | 47 |
| 107 | Essential Oils from <i>Salvia</i> sp. (Lamiaceae). III. Composition and Antimicrobial Activity of the Essential Oil of <i>Salvia palaestina</i> Benth. Growing Wild in Lebanon. <i>Journal of Essential Oil Research</i> , 2005, 17, 419-421. | 1.3 | 17 |
| 108 | Composition of the Essential Oil of <i>Nepeta curviflora</i> Boiss. (Lamiaceae) from Lebanon. <i>Journal of Essential Oil Research</i> , 2005, 17, 268-270. | 1.3 | 19 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 109 | Chemical Composition and Antibacterial Activity of the Essential Oil of a 1,8-Cineole Chemotype of <i>Mentha aquatica</i> L. Growing Wild in Turkey. <i>Journal of Essential Oil-bearing Plants: JEOP</i> , 2005, 8, 148-153. | 0.7 | 12 |
| 110 | Antibacterial Activity of <i>Cuminum cyminum</i> L. and <i>Carum carvi</i> L. Essential Oils. <i>Journal of Agricultural and Food Chemistry</i> , 2005, 53, 57-61. | 2.4 | 369 |
| 111 | Chemical Composition of the Essential Oil of <i>Phagnalon Saxatile</i> (L.) Cass. (Asteraceae) Growing Wild in Southern Italy. <i>Journal of Essential Oil-bearing Plants: JEOP</i> , 2005, 8, 258-263. | 0.7 | 9 |
| 112 | Essential oil from aerial parts of <i>Vernonia colorata</i> Drake and <i>Vernonia nigritiana</i> Oliver et Hiern. (Asteraceae) growing wild in Mali. <i>Journal of Essential Oil-bearing Plants: JEOP</i> , 2004, 7, 267-274. | 0.7 | 2 |
| 113 | Composition of the essential oil from flowerheads of <i>Chrysanthemum coronarium</i> L. (Asteraceae) growing wild in Southern Italy. <i>Flavour and Fragrance Journal</i> , 2004, 19, 149-152. | 1.2 | 29 |
| 114 | Antibacterial activity of <i>Tagetes minuta</i> L. (Asteraceae) essential oil with different chemical composition. <i>Flavour and Fragrance Journal</i> , 2004, 19, 574-578. | 1.2 | 56 |
| 115 | Chemical composition of the essential oil of <i>Salvia multicaulis</i> Vahl. var. <i>simplicifolia</i> Boiss. growing wild in Lebanon. <i>Journal of Chromatography A</i> , 2004, 1052, 237-240. | 1.8 | 40 |
| 116 | Antibacterial Activity of <i>Coriandrum sativum</i> L. and <i>Foeniculum vulgare</i> Miller Var. <i>vulgare</i> (Miller) Essential Oils. <i>Journal of Agricultural and Food Chemistry</i> , 2004, 52, 7862-7866. | 2.4 | 226 |
| 117 | 4-hydroxybenzyl glucosinolate from <i>Cardaria draba</i> (Cruciferae). <i>Biochemical Systematics and Ecology</i> , 2003, 31, 1205-1207. | 0.6 | 6 |
| 118 | Volatile components of <i>Centaurea cineraria</i> L. subsp. <i>umbrosa</i> (Lacaita) Pign. and <i>Centaurea napifolia</i> L. (Asteraceae), two species growing wild in Sicily. <i>Flavour and Fragrance Journal</i> , 2003, 18, 248-251. | 1.2 | 50 |
| 119 | Composition of the essential oil of <i>Pallenis spinosa</i> (L.) Cass. (Asteraceae). <i>Flavour and Fragrance Journal</i> , 2003, 18, 195-197. | 1.2 | 9 |
| 120 | Composition and antibacterial activity of the essential oil of <i>Anisochilus carnosus</i> (Linn. ?!) Benth., a Tamil plant acclimatized in Sicily. <i>Flavour and Fragrance Journal</i> , 2003, 18, 202-204. | 1.2 | 10 |
| 121 | Chemical composition of essential oils of <i>Senecio nutans</i> Sch.-Bip. (Asteraceae). <i>Flavour and Fragrance Journal</i> , 2003, 18, 234-236. | 1.2 | 24 |
| 122 | Chemical composition and antibacterial activity of essential oils from five culinary herbs of the Lamiaceae family growing in Campania, Southern Italy. <i>Journal of Essential Oil-bearing Plants: JEOP</i> , 2003, 6, 166-173. | 0.7 | 6 |
| 123 | Chemical Composition and Antibacterial Activity of Essential Oils from <i>Thymus spinulosus</i> Ten. (Lamiaceae). <i>Journal of Agricultural and Food Chemistry</i> , 2003, 51, 3849-3853. | 2.4 | 43 |
| 124 | Composition of the Essential Oil of <i>Nepeta betonicifolia</i> C.A. Meyer (Lamiaceae) from Turkey. <i>Journal of Essential Oil Research</i> , 2003, 15, 200-201. | 1.3 | 14 |
| 125 | Antibacterial Evaluation of Cnicin and Some Natural and Semisynthetic Analogues. <i>Planta Medica</i> , 2003, 69, 277-281. | 0.7 | 21 |
| 126 | Potential allelochemicals from the essential oil of <i>Ruta graveolens</i> . <i>Phytochemistry</i> , 2002, 61, 573-578. | 1.4 | 136 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 127 | Essential oil of two <i>Lippia</i> spp. (Verbenaceae) growing wild in Guatemala. <i>Flavour and Fragrance Journal</i> , 2001, 16, 169-171. | 1.2 | 37 |
| 128 | <i>Eupatorium cannabinum</i> L. ssp. <i>cannabinum</i> (Asteraceae) Essential Oil: Chemical Composition and Antibacterial Activity. <i>Journal of Essential Oil Research</i> , 2001, 13, 463-466. | 1.3 | 17 |
| 129 | Composition and antibacterial activity of the essential oil from <i>Crithmum maritimum</i> L. (Apiaceae) growing wild in Turkey. <i>Flavour and Fragrance Journal</i> , 2000, 15, 186-189. | 1.2 | 53 |
| 130 | Flavonoid Glycosides of <i>Barbarea vulgaris</i> L. (Brassicaceae). <i>Journal of Agricultural and Food Chemistry</i> , 2000, 48, 2659-2662. | 2.4 | 44 |
| 131 | Two new quercetagenin O-glucosides from <i>Tagetes mandonii</i> . <i>Biochemical Systematics and Ecology</i> , 1999, 27, 309-311. | 0.6 | 6 |
| 132 | Chemical composition of the essential oil from <i>Tagetes mandonii</i> Sch. Bip. (Asteraceae). <i>Flavour and Fragrance Journal</i> , 1999, 14, 32-34. | 1.2 | 15 |
| 133 | <i>Carica candicans</i> Gray (Mito), an Alimentary Resource from Peruvian Flora. <i>Journal of Agricultural and Food Chemistry</i> , 1999, 47, 3682-3684. | 2.4 | 4 |
| 134 | Essential oils from two Peruvian <i>Satureja</i> species. <i>Flavour and Fragrance Journal</i> , 1998, 13, 1-4. | 1.2 | 38 |
| 135 | Composition of the essential oil of <i>Tagetes filifolia</i> Lag.. <i>Flavour and Fragrance Journal</i> , 1998, 13, 145-147. | 1.2 | 17 |
| 136 | Volatile constituents of <i>Minthostachys setosa</i> (Briq.) Epl. (Lamiaceae) from Peru. <i>Flavour and Fragrance Journal</i> , 1998, 13, 263-265. | 1.2 | 9 |
| 137 | Essential Oils from <i>Salvia</i> spp. (Lamiaceae). II. Chemical Composition of the Essential Oil from <i>Salvia pratensis</i> L. subsp. <i>haematodes</i> (L.) Briq. Inflorescences. <i>Journal of Essential Oil Research</i> , 1998, 10, 135-137. | 1.3 | 9 |
| 138 | Chemical Composition and Antimicrobial Screening of the Essential Oil of <i>Minthostachys verticillata</i> (Griseb.) Epl. (Lamiaceae). <i>Journal of Essential Oil Research</i> , 1998, 10, 61-65. | 1.3 | 31 |
| 139 | Essential Oils from <i>Salvia</i> spp. (Lamiaceae). I. Chemical Composition of the Essential Oils from <i>Salvia glutinosa</i> L. Growing Wild in Southern Italy. <i>Journal of Essential Oil Research</i> , 1997, 9, 151-157. | 1.3 | 36 |
| 140 | Quercetagenin 6-O- β -D-glucopyranoside from <i>Tagetes mandonii</i> . <i>Phytochemistry</i> , 1997, 45, 201-202. | 1.4 | 8 |
| 141 | Essential oil of <i>Eremocharis triradiata</i> (Wolff.) Johnston (Apiaceae) growing wild in Perù. <i>Flavour and Fragrance Journal</i> , 1997, 12, 257-259. | 1.2 | 5 |
| 142 | Influence of Harvesting Time on Yield and Composition of the Essential Oil of a Thyme (<i>Thymus pulegioides</i> L.) Growing Wild in Campania (Southern Italy). <i>Journal of Agricultural and Food Chemistry</i> , 1996, 44, 1327-1332. | 2.4 | 160 |
| 143 | Constituents of <i>Vitex agnus-castus</i> L. Essential Oil. <i>Flavour and Fragrance Journal</i> , 1996, 11, 179-182. | 1.2 | 30 |
| 144 | Composition of the essential oil of <i>Minthostachys spicata</i> (Benth.) Epl. <i>Flavour and Fragrance Journal</i> , 1995, 10, 43-45. | 1.2 | 8 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 145 | Composition of the essential oil of <i>Santolina neapolitana</i> Jordan et Fourc. Flavour and Fragrance Journal, 1994, 9, 77-79. | 1.2 | 8 |
| 146 | Essential oil of a possible new chemotype of <i>Crithmum maritimum</i> L. growing in Campania (Southern Italy). Journal of Agricultural and Food Chemistry, 1994, 42, 154-158. | 2.4 | 111 |
| 147 | Supercritical carbon dioxide extraction of chamomile essential oil and its analysis by gas chromatography-mass spectrometry. Journal of Agricultural and Food Chemistry, 1994, 42, 154-158. | 2.4 | 111 |
| 148 | Medicinal plants and phytotherapy in the Amalfitan Coast, Salerno Province, Campania, Southern Italy. Journal of Ethnopharmacology, 1993, 39, 39-51. | 2.0 | 106 |
| 149 | Traditional phytotherapy in the Peninsula Sorrentina, Campania, Southern Italy. Journal of Ethnopharmacology, 1992, 36, 113-125. | 2.0 | 98 |
| 150 | Chemical constituents of some mushrooms. Journal of the Science of Food and Agriculture, 1992, 58, 499-503. | 1.7 | 17 |
| 151 | Fatty acids, free amino acids and sterols from some species of <i>Stropharia</i> and <i>Stereum</i> . Biochemical Systematics and Ecology, 1990, 18, 103-106. | 0.6 | 8 |
| 152 | Fatty acid and free amino acid content of some mushrooms. Journal of the Science of Food and Agriculture, 1990, 51, 91-96. | 1.7 | 17 |
| 153 | Oligosaccharides in Five Different <i>Vicia faba</i> L. Cultivars. Biochemical Systematics and Ecology, 1989, 17, 559-561. | 0.6 | 9 |
| 154 | Chemical constituents of some basidiomycetes. Journal of the Science of Food and Agriculture, 1988, 45, 337-345. | 1.7 | 30 |
| 155 | Chemical constituents of some species of Agaricaceae. Biochemical Systematics and Ecology, 1988, 16, 601-604. | 0.6 | 14 |
| 156 | Free amino acids from different cultivars of <i>Vicia faba</i> . Journal of Agricultural and Food Chemistry, 1983, 31, 836-838. | 2.4 | 8 |
| 157 | Sterols, fatty acids and free amino acids from two <i>Helvella</i> species. Biochemical Systematics and Ecology, 1982, 10, 285-287. | 0.6 | 5 |
| 158 | Sterols from three <i>Lactarius</i> species. Biochemical Systematics and Ecology, 1981, 9, 247-248. | 0.6 | 8 |