

# Håkan Bengtsson

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9004847/publications.pdf>

Version: 2024-02-01

15  
papers

1,188  
citations

933447

10  
h-index

1058476

14  
g-index

15  
all docs

15  
docs citations

15  
times ranked

1123  
citing authors

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 1  | Injuries affect team performance negatively in professional football: an 11-year follow-up of the UEFA Champions League injury study. <i>British Journal of Sports Medicine</i> , 2013, 47, 738-742.   | 6.7 | 540       |
| 2  | Muscle injury rates in professional football increase with fixture congestion: an 11-year follow-up of the UEFA Champions League injury study. <i>British Journal of Sports Medicine</i> , 2013, 47, 743-747.  | 6.7 | 191       |
| 3  | Time before return to play for the most common injuries in professional football: a 16-year follow-up of the UEFA Elite Club Injury Study. <i>British Journal of Sports Medicine</i> , 2020, 54, 421-426.  | 6.7 | 138       |
| 4  | Injury rates decreased in men's professional football: an 18-year prospective cohort study of almost 12 000 injuries sustained during 1.8 million hours of play. <i>British Journal of Sports Medicine</i> , 2021, 55, 1084-1092.  | 6.7 | 88        |
| 5  | Muscle injury rate in professional football is higher in matches played within 5 days since the previous match: a 14-year prospective study with more than 130 000 match observations. <i>British Journal of Sports Medicine</i> , 2018, 52, 1116-1122.  | 6.7 | 65        |
| 6  | Upper extremity injuries in male elite football players. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , 2013, 21, 1626-1632.  | 4.2 | 37        |
| 7  | Epidemiology and return to play following isolated syndesmotic injuries of the ankle: a prospective cohort study of 3677 male professional footballers in the UEFA Elite Club Injury Study. <i>British Journal of Sports Medicine</i> , 2019, 53, 959-964.   | 6.7 | 35        |
| 8  | Match Injury Rates in Professional Soccer Vary With Match Result, Match Venue, and Type of Competition. <i>American Journal of Sports Medicine</i> , 2013, 41, 1505-1510.  | 4.2 | 23        |
| 9  | Role of illness in male professional football: not a major contributor to time loss. <i>British Journal of Sports Medicine</i> , 2016, 50, 699-702.  | 6.7 | 22        |
| 10 | Influence of the COVID-19 Lockdown and Restart on the Injury Incidence and Injury Burden in Men's Professional Football Leagues in 2020: The UEFA Elite Club Injury Study. <i>Sports Medicine - Open</i> , 2022, 8, 67.  | 3.1 | 15        |
| 11 | Few training sessions between return to play and first match appearance are associated with an increased propensity for injury: a prospective cohort study of male professional football players during 16 consecutive seasons. <i>British Journal of Sports Medicine</i> , 2020, 54, 427-432.         | 6.7 | 13        |
| 12 | Hand, Wrist, and Forearm Injuries in Male Professional Soccer Players: A Prospective Cohort Study of 558 Team-Seasons From 2001-2002 to 2018-2019. <i>Orthopaedic Journal of Sports Medicine</i> , 2021, 9, 232596712097709.   | 1.7 | 8         |
| 13 | Injury epidemiology in professional football in South America compared with Europe. <i>BMJ Open Sport and Exercise Medicine</i> , 2021, 7, e001172.  | 2.9 | 5         |
| 14 | Anterior ankle impingement syndrome is less frequent, but associated with a longer absence and higher re-injury rate compared to posterior syndrome: a prospective cohort study of 6754 male professional soccer players. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , 2022, 30, 4262-4269. | 4.2 | 5         |
| 15 | Re-injuries in Professional Football: The UEFA Elite Club Injury Study. , 2018, , 953-962.   |     | 3         |