

Maria A Baturova

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9005893/publications.pdf>

Version: 2024-02-01

11
papers

152
citations

1307594

7
h-index

1281871

11
g-index

11
all docs

11
docs citations

11
times ranked

298
citing authors

#	ARTICLE	IF	CITATIONS
1	Electrocardiographic and Echocardiographic predictors of paroxysmal atrial fibrillation detected after ischemic stroke. <i>BMC Cardiovascular Disorders</i> , 2016, 16, 209.	1.7	39
2	Predictors of new onset atrial fibrillation during 10-year follow-up after first-ever ischemic stroke. <i>International Journal of Cardiology</i> , 2015, 199, 248-252.	1.7	25
3	Atrial fibrillation in patients with ischaemic stroke in the Swedish national patient registers: how much do we miss?. <i>Europace</i> , 2014, 16, 1714-1719.	1.7	17
4	Atrial fibrillation as a clinical characteristic of arrhythmogenic right ventricular cardiomyopathy: Experience from the Nordic ARVC Registry. <i>International Journal of Cardiology</i> , 2020, 298, 39-43.	1.7	14
5	Atrial time and voltage dispersion are both needed to predict new-onset atrial fibrillation in ischemic stroke patients. <i>BMC Cardiovascular Disorders</i> , 2017, 17, 200.	1.7	13
6	Interatrial block in prediction of all-cause mortality after first-ever ischemic stroke. <i>BMC Cardiovascular Disorders</i> , 2019, 19, 37.	1.7	12
7	Secondary prevention and lifestyle indices after stroke in a long-term perspective. <i>Acta Neurologica Scandinavica</i> , 2018, 138, 227-234.	2.1	9
8	Documentation of atrial fibrillation prior to first-ever ischemic stroke. <i>Acta Neurologica Scandinavica</i> , 2014, 129, 412-419.	2.1	7
9	Usefulness of Electrocardiographic Left Atrial Abnormality to Predict Response to Cardiac Resynchronization Therapy in Patients With Mild Heart Failure and Left Bundle Branch Block (a) Tj ETQq1 1 0.784314 rgBT /Qverlock 1.6	1.6	7
10	Evolution of P-wave indices during long-term follow-up as markers of atrial substrate progression in arrhythmogenic right ventricular cardiomyopathy. <i>Europace</i> , 2021, 23, i29-i37.	1.7	5
11	Non-permanent atrial fibrillation and oral anticoagulant therapy are related to survival during 10 years after first-ever ischemic stroke. <i>International Journal of Cardiology</i> , 2017, 232, 134-139.	1.7	4