

List of Publications by Citations

**Source:** <https://exaly.com/author-pdf/9015538/tao-li-publications-by-citations.pdf>  
**Version:** 2024-04-04

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.  
The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

141 papers	7,304 citations	38 h-index	83 g-index
147 ext. papers	8,566 ext. citations	6.8 avg, IF	4.92 L-index

#	Paper	IF	Citations
141	Large recurrent microdeletions associated with schizophrenia. <i>Nature</i> , <b>2008</b> , 455, 232-6	50.4	1427
140	Prevalence of mental disorders in China: a cross-sectional epidemiological study. <i>Lancet Psychiatry</i> , <b>2019</b> , 6, 211-224	23.3	594
139	MDR1 pharmacogenetics: frequency of the C3435T mutation in exon 26 is significantly influenced by ethnicity. <i>Pharmacogenetics and Genomics</i> , <b>2001</b> , 11, 217-21		335
138	Short-term effects of antipsychotic treatment on cerebral function in drug-naïve first-episode schizophrenia revealed by "resting state" functional magnetic resonance imaging. <i>Archives of General Psychiatry</i> , <b>2010</b> , 67, 783-92		294
137	Genome-wide association identifies a common variant in the reelin gene that increases the risk of schizophrenia only in women. <i>PLoS Genetics</i> , <b>2008</b> , 4, e28	6	270
136	Genome-wide association study identifies a susceptibility locus for schizophrenia in Han Chinese at 11p11.2. <i>Nature Genetics</i> , <b>2011</b> , 43, 1228-31	36.3	231
135	Association of cerebral deficits with clinical symptoms in antipsychotic-naïve first-episode schizophrenia: an optimized voxel-based morphometry and resting state functional connectivity study. <i>American Journal of Psychiatry</i> , <b>2009</b> , 166, 196-205	11.9	217
134	Ethnic variation in the thymidylate synthase enhancer region polymorphism among Caucasian and Asian populations. <i>Genomics</i> , <b>1999</b> , 58, 310-2	4.3	177
133	Reduced default mode network functional connectivity in patients with recurrent major depressive disorder. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2019</b> , 116, 9078-9083	11.5	168
132	Association between 5-HT2A gene promoter polymorphism and anorexia nervosa. <i>Lancet, The</i> , <b>1997</b> , 350, 412	40	146
131	The prefrontal dysfunction in individuals with Internet gaming disorder: a meta-analysis of functional magnetic resonance imaging studies. <i>Addiction Biology</i> , <b>2015</b> , 20, 799-808	4.6	130
130	Localization of cerebral functional deficits in treatment-naïve, first-episode schizophrenia using resting-state fMRI. <i>NeuroImage</i> , <b>2010</b> , 49, 2901-6	7.9	120
129	Changes of brain morphometry in first-episode, drug-naïve, non-late-life adult patients with major depression: an optimized voxel-based morphometry study. <i>Biological Psychiatry</i> , <b>2010</b> , 67, 186-8	7.9	118
128	Hair cortisol level as a biomarker for altered hypothalamic-pituitary-adrenal activity in female adolescents with posttraumatic stress disorder after the 2008 Wenchuan earthquake. <i>Biological Psychiatry</i> , <b>2012</b> , 72, 65-9	7.9	116
127	Risk indicators for post-traumatic stress disorder in adolescents exposed to the 5.12 Wenchuan earthquake in China. <i>Psychiatry Research</i> , <b>2011</b> , 189, 385-91	9.9	108
126	Anatomical and functional brain abnormalities in drug-naïve first-episode schizophrenia. <i>American Journal of Psychiatry</i> , <b>2013</b> , 170, 1308-16	11.9	106
125	Brain-Wide Analysis of Functional Connectivity in First-Episode and Chronic Stages of Schizophrenia. <i>Schizophrenia Bulletin</i> , <b>2017</b> , 43, 436-448	1.3	103

124	Allelic functional variation of serotonin transporter expression is a susceptibility factor for late onset Alzheimer's disease. <i>NeuroReport</i> , <b>1997</b> , 8, 683-6	1.7	98
123	Voxel-based, brain-wide association study of aberrant functional connectivity in schizophrenia implicates thalamocortical circuitry. <i>NPJ Schizophrenia</i> , <b>2015</b> , 1, 15016	5.5	94
122	Association analysis of the DRD4 and COMT genes in methamphetamine abuse. <i>American Journal of Medical Genetics Part A</i> , <b>2004</b> , 129B, 120-4		85
121	Evaluation of a susceptibility gene for schizophrenia: genotype based meta-analysis of RGS4 polymorphisms from thirteen independent samples. <i>Biological Psychiatry</i> , <b>2006</b> , 60, 152-62	7.9	80
120	Altered cortical thickness related to clinical severity but not the untreated disease duration in schizophrenia. <i>Schizophrenia Bulletin</i> , <b>2015</b> , 41, 201-10	1.3	75
119	Evidence for association between novel polymorphisms in the PRODH gene and schizophrenia in a Chinese population. <i>American Journal of Medical Genetics Part A</i> , <b>2004</b> , 129B, 13-5		68
118	Allelic association analysis of the dopamine D2, D3, 5-HT2A, and GABA(A)gamma2 receptors and serotonin transporter genes with heroin abuse in Chinese subjects. <i>American Journal of Medical Genetics Part A</i> , <b>2002</b> , 114, 329-35		68
117	No association between schizophrenia and homozygosity at the D3 dopamine receptor gene. <i>American Journal of Medical Genetics Part A</i> , <b>1993</b> , 48, 83-6		66
116	Potential linkage disequilibrium between schizophrenia and locus D22S278 on the long arm of chromosome 22. <i>American Journal of Medical Genetics Part A</i> , <b>1995</b> , 60, 465-7		55
115	Contrasting and convergent patterns of amygdala connectivity in mania and depression: a resting-state study. <i>Journal of Affective Disorders</i> , <b>2015</b> , 173, 53-8	6.6	52
114	Reaction time of the Continuous Performance Test is an endophenotypic marker for schizophrenia: a study of first-episode neuroleptic-naïve schizophrenia, their non-psychotic first-degree relatives and healthy population controls. <i>Schizophrenia Research</i> , <b>2007</b> , 89, 293-8	3.6	51
113	Neurocognitive deficits in first-episode schizophrenic patients and their first-degree relatives. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , <b>2007</b> , 144B, 407-16	3.5	50
112	-141 C del/ins polymorphism of the dopamine receptor 2 gene is associated with schizophrenia in a British population. <i>American Journal of Medical Genetics Part A</i> , <b>1999</b> , 88, 407-10		50
111	Abnormal Resting-State Connectivity in a Substantia Nigra-Related Striato-Thalamo-Cortical Network in a Large Sample of First-Episode Drug-Naïve Patients With Schizophrenia. <i>Schizophrenia Bulletin</i> , <b>2018</b> , 44, 419-431	1.3	46
110	Assessment of white matter abnormalities in paranoid schizophrenia and bipolar mania patients. <i>Psychiatry Research - Neuroimaging</i> , <b>2011</b> , 194, 347-353	2.9	46
109	Association study of 45 candidate genes in nicotine dependence in Han Chinese. <i>Addictive Behaviors</i> , <b>2012</b> , 37, 622-6	4.2	45
108	Association analysis between dopamine receptor genes and bipolar affective disorder. <i>Psychiatry Research</i> , <b>1999</b> , 86, 193-201	9.9	45
107	Voxel based morphometric and diffusion tensor imaging analysis in male bipolar patients with first-episode mania. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , <b>2012</b> , 36, 231-8	5.5	44

106	Association of the dysbindin gene with bipolar affective disorder. <i>American Journal of Psychiatry</i> , <b>2006</b> , 163, 1636-8	11.9	43
105	Common variants on Xq28 conferring risk of schizophrenia in Han Chinese. <i>Schizophrenia Bulletin</i> , <b>2014</b> , 40, 777-86	1.3	40
104	A quantitative association study between schizotypal traits and COMT, PRODH and BDNF genes in a healthy Chinese population. <i>Psychiatry Research</i> , <b>2007</b> , 153, 7-15	9.9	39
103	Association analysis of polymorphisms in the DRD4 gene and heroin abuse in Chinese subjects. <i>American Journal of Medical Genetics Part A</i> , <b>2000</b> , 96, 616-21		34
102	White matter alterations in first episode treatment-naïve patients with deficit schizophrenia: a combined VBM and DTI study. <i>Scientific Reports</i> , <b>2015</b> , 5, 12994	4.9	33
101	Lack of association between schizophrenia and the phospholipase-A2 genes cPLA2 and sPLA2. <i>American Journal of Medical Genetics Part A</i> , <b>2001</b> , 105, 246-249		33
100	Overlapping clusters of gray matter deficits in paranoid schizophrenia and psychotic bipolar mania with family history. <i>Neuroscience Letters</i> , <b>2011</b> , 489, 94-8	3.3	32
99	Analysis of CAG/CTG repeat size in Chinese subjects with schizophrenia and bipolar affective disorder using the repeat expansion detection method. <i>Biological Psychiatry</i> , <b>1998</b> , 44, 1160-5	7.9	32
98	Possible association between Cys311Ser polymorphism of paraoxonase 2 gene and late-onset Alzheimer's disease in Chinese. <i>Molecular Brain Research</i> , <b>2004</b> , 120, 201-4		32
97	Reward pathway dysfunction in gambling disorder: A meta-analysis of functional magnetic resonance imaging studies. <i>Behavioural Brain Research</i> , <b>2014</b> , 275, 243-51	3.4	31
96	Advanced paternal age increases the risk of schizophrenia and obsessive-compulsive disorder in a Chinese Han population. <i>Psychiatry Research</i> , <b>2012</b> , 198, 353-9	9.9	31
95	Psychiatric epidemiological surveys in China 1960-2010: how real is the increase of mental disorders?. <i>Current Opinion in Psychiatry</i> , <b>2011</b> , 24, 324-30	4.9	31
94	Positional pathway screen of wnt signaling genes in schizophrenia: association with DKK4. <i>Biological Psychiatry</i> , <b>2008</b> , 63, 13-6	7.9	31
93	Association analysis of polymorphisms in the 1bpioid gene and heroin abuse in Chinese subjects. <i>Addiction Biology</i> , <b>2000</b> , 5, 181-6	4.6	30
92	Transmission disequilibrium analysis of a triplet repeat within the hKCa3 gene using family trios with schizophrenia. <i>Biochemical and Biophysical Research Communications</i> , <b>1998</b> , 251, 662-5	3.4	30
91	Voxel-based morphometric analysis on the volume of gray matter in bipolar I disorder. <i>Psychiatry Research - Neuroimaging</i> , <b>2011</b> , 191, 92-7	2.9	29
90	An association study of RGS4 polymorphisms with clinical phenotypes of schizophrenia in a Chinese population. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , <b>2008</b> , 147B, 77-85	3.5	29
89	Increased co-expression of genes harboring the damaging de novo mutations in Chinese schizophrenic patients during prenatal development. <i>Scientific Reports</i> , <b>2015</b> , 5, 18209	4.9	27

88	Common and distinct patterns of grey matter alterations in borderline personality disorder and bipolar disorder: voxel-based meta-analysis. <i>British Journal of Psychiatry</i> , <b>2019</b> , 215, 395-403	5.4	26
87	An epidemiological survey of alcohol use disorders in a Tibetan population. <i>Psychiatry Research</i> , <b>2008</b> , 159, 56-66	9.9	25
86	Sex differences in the clinical characteristics and brain gray matter volume alterations in unmedicated patients with major depressive disorder. <i>Scientific Reports</i> , <b>2017</b> , 7, 2515	4.9	24
85	No association between alleles or genotypes at the dopamine transporter gene and schizophrenia. <i>Psychiatry Research</i> , <b>1994</b> , 52, 17-23	9.9	24
84	The China Mental Health Survey (CMHS): I. background, aims and measures. <i>Social Psychiatry and Psychiatric Epidemiology</i> , <b>2016</b> , 51, 1559-1569	4.5	23
83	Classification of First-Episode Schizophrenia Using Multimodal Brain Features: A Combined Structural and Diffusion Imaging Study. <i>Schizophrenia Bulletin</i> , <b>2019</b> , 45, 591-599	1.3	23
82	Premorbid tobacco smoking is associated with later age at onset in schizophrenia. <i>Psychiatry Research</i> , <b>2010</b> , 178, 461-6	9.9	23
81	102T/C polymorphism of serotonin receptor type 2A gene is not associated with schizophrenia in either Chinese or British populations <b>1999</b> , 88, 95-98		23
80	Association study of dopamine receptor genes polymorphism with cognitive functions in bipolar I disorder patients. <i>Journal of Affective Disorders</i> , <b>2015</b> , 170, 85-90	6.6	22
79	Correlation and familial aggregation of dimensions of psychosis in affected sibling pairs from China. <i>British Journal of Psychiatry</i> , <b>2008</b> , 193, 305-10	5.4	22
78	Mutation screening and association analysis of the parkin gene in Parkinson's disease patients from South-West China. <i>European Neurology</i> , <b>2003</b> , 49, 85-9	2.1	22
77	Genome-Wide Association of Heroin Dependence in Han Chinese. <i>PLoS ONE</i> , <b>2016</b> , 11, e0167388	3.7	22
76	The role of single nucleotide polymorphism of D2 dopamine receptor gene on major depressive disorder and response to antidepressant treatment. <i>Psychiatry Research</i> , <b>2012</b> , 200, 1047-50	9.9	21
75	A COMT gene haplotype associated with methamphetamine abuse. <i>Pharmacogenetics and Genomics</i> , <b>2011</b> , 21, 731-40	1.9	20
74	Factor structures of the neurocognitive assessments and familial analysis in first-episode schizophrenia patients, their relatives and controls. <i>Australian and New Zealand Journal of Psychiatry</i> , <b>2010</b> , 44, 109-19	2.6	19
73	Effect of Damaging Rare Mutations in Synapse-Related Gene Sets on Response to Short-term Antipsychotic Medication in Chinese Patients With Schizophrenia: A Randomized Clinical Trial. <i>JAMA Psychiatry</i> , <b>2018</b> , 75, 1261-1269	14.5	19
72	Normative data on a battery of neuropsychological tests in the Han Chinese population. <i>Journal of Neuropsychology</i> , <b>2011</b> , 5, 126-42	2.6	18
71	Prognostic Utility of Multivariate Morphometry in Schizophrenia. <i>Frontiers in Psychiatry</i> , <b>2019</b> , 10, 245	5	17

70	A splitting brain: Imbalanced neural networks in schizophrenia. <i>Psychiatry Research - Neuroimaging</i> , <b>2015</b> , 232, 145-53	2.9	17
69	Analysis of voxel-mirrored homotopic connectivity in medication-free, current major depressive disorder. <i>Journal of Affective Disorders</i> , <b>2018</b> , 240, 171-176	6.6	17
68	Identification of neuroglycan C and interacting partners as potential susceptibility genes for schizophrenia in a Southern Chinese population. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , <b>2010</b> , 153B, 103-13	3.5	16
67	The differential clinical and neurocognitive profiles of COMT SNP rs165599 genotypes in schizophrenia. <i>Journal of the International Neuropsychological Society</i> , <b>2005</b> , 11, 202-4	3.1	16
66	Genetic association of the cytochrome c oxidase-related genes with Alzheimer's disease in Han Chinese. <i>Neuropsychopharmacology</i> , <b>2018</b> , 43, 2264-2276	8.7	15
65	No association of the LRRK2 genetic variants with Alzheimer's disease in Han Chinese individuals. <i>Neurobiology of Aging</i> , <b>2014</b> , 35, 444.e5-9	5.6	15
64	Sex-Specific Patterns of Aberrant Brain Function in First-Episode Treatment-Naïve Patients with Schizophrenia. <i>International Journal of Molecular Sciences</i> , <b>2015</b> , 16, 16125-43	6.3	15
63	Genome-wide association analysis with gray matter volume as a quantitative phenotype in first-episode treatment-naïve patients with schizophrenia. <i>PLoS ONE</i> , <b>2013</b> , 8, e75083	3.7	15
62	White Matter Abnormalities in Major Depression Biotypes Identified by Diffusion Tensor Imaging. <i>Neuroscience Bulletin</i> , <b>2019</b> , 35, 867-876	4.3	14
61	Contingent negative variation in patients with deficit schizophrenia or bipolar I disorder with psychotic features: measurement and correlation with clinical characteristics. <i>Nordic Journal of Psychiatry</i> , <b>2015</b> , 69, 196-203	2.3	14
60	"Brain Connectivity Deviates by Sex and Hemisphere in the First Episode of Schizophrenia"-A Route to the Genetic Basis of Language and Psychosis?. <i>Schizophrenia Bulletin</i> , <b>2019</b> , 45, 484-494	1.3	14
59	Bipolar 1 disorder is not associated with the RGS4, PRODH, COMT and GRK3 genes. <i>Psychiatric Genetics</i> , <b>2006</b> , 16, 229-30	2.9	14
58	Age-Related Reduction in Cortical Thickness in First-Episode Treatment-Naïve Patients with Schizophrenia. <i>Neuroscience Bulletin</i> , <b>2019</b> , 35, 688-696	4.3	13
57	Hair thyroid hormones concentration in patients with depression changes with disease episodes in female Chinese. <i>Psychiatry Research</i> , <b>2014</b> , 220, 251-3	9.9	13
56	Whole-exome sequencing for the identification of susceptibility genes of Kashin-Beck disease. <i>PLoS ONE</i> , <b>2014</b> , 9, e92298	3.7	13
55	Multi-components of evoked-brain potentials in deficit and nondeficit schizophrenia. <i>Asia-Pacific Psychiatry</i> , <b>2013</b> , 5, 69-79	3.2	13
54	The 1239G/C polymorphism in exon 5 of BACE1 gene may be associated with sporadic Alzheimer's disease in Chinese Hans. <i>American Journal of Medical Genetics Part A</i> , <b>2004</b> , 124B, 54-7		13
53	Associations of Internet Addiction Severity With Psychopathology, Serious Mental Illness, and Suicidality: Large-Sample Cross-Sectional Study. <i>Journal of Medical Internet Research</i> , <b>2020</b> , 22, e17560	7.6	13



52	Association of white matter deficits with clinical symptoms in antipsychotic-naïve first-episode schizophrenia: an optimized VBM study using 3T. <i>Magnetic Resonance Materials in Physics, Biology, and Medicine</i> , <b>2014</b> , 27, 283-90	2.8	12
51	A joint study of whole exome sequencing and structural MRI analysis in major depressive disorder. <i>Psychological Medicine</i> , <b>2020</b> , 50, 384-395	6.9	12
50	Functional brain networks in never-treated and treated long-term ill schizophrenia patients. <i>Neuropsychopharmacology</i> , <b>2019</b> , 44, 1940-1947	8.7	11
49	Low-rank network signatures in the triple network separate schizophrenia and major depressive disorder. <i>NeuroImage: Clinical</i> , <b>2019</b> , 22, 101725	5.3	11
48	Association of APC and REEP5 gene polymorphisms with major depression disorder and treatment response to antidepressants in a Han Chinese population. <i>General Hospital Psychiatry</i> , <b>2012</b> , 34, 571-7	5.6	11
47	PRODH gene is associated with executive function in schizophrenic families. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , <b>2008</b> , 147B, 654-7	3.5	11
46	Co-treatment of buspirone with atypical antipsychotic drugs (AAPDs) improved neurocognitive function in chronic schizophrenia. <i>Schizophrenia Research</i> , <b>2019</b> , 209, 135-140	3.6	10
45	Convergence and divergence of neurocognitive patterns in schizophrenia and depression. <i>Schizophrenia Research</i> , <b>2018</b> , 192, 327-334	3.6	10
44	Neurocognitive Graphs of First-Episode Schizophrenia and Major Depression Based on Cognitive Features. <i>Neuroscience Bulletin</i> , <b>2018</b> , 34, 312-320	4.3	10
43	Detection of metabolites in the white matter of frontal lobes and hippocampus with proton in first-episode treatment-naïve schizophrenia patients. <i>Microbial Biotechnology</i> , <b>2012</b> , 6, 166-75	3.3	10
42	CNTF and psychiatric disorders. <i>Nature Genetics</i> , <b>1996</b> , 13, 143-4	36.3	10
41	Rural birth/upbringing and childhood adversities are associated with psychotic experiences in university students in China. <i>Schizophrenia Research</i> , <b>2019</b> , 209, 105-112	3.6	9
40	A powerful conditional gene-based association approach implicated functionally important genes for schizophrenia. <i>Bioinformatics</i> , <b>2019</b> , 35, 628-635	7.2	9
39	The androgen receptor gene polyglycine repeat polymorphism is associated with memory performance in healthy Chinese individuals. <i>Psychoneuroendocrinology</i> , <b>2009</b> , 34, 947-52	5	9
38	Affected sibling pair linkage analysis of qualitative and quantitative traits for schizophrenia on chromosome 22 in a Chinese population. <i>American Journal of Medical Genetics Part A</i> , <b>2001</b> , 105, 321-7		9
37	The development and validation of Huaxi emotional-distress index (HEI): A Chinese questionnaire for screening depression and anxiety in non-psychiatric clinical settings. <i>Comprehensive Psychiatry</i> , <b>2017</b> , 76, 87-97	7.3	8
36	Depth-dependent abnormal cortical myelination in first-episode treatment-naïve schizophrenia. <i>Human Brain Mapping</i> , <b>2020</b> , 41, 2782-2793	5.9	8
35	No association between a promoter polymorphism in the noradrenaline transporter gene and anorexia nervosa. <i>Psychiatric Genetics</i> , <b>2007</b> , 17, 247-8	2.9	8

34	Increased expression of a novel miRNA in peripheral blood is negatively correlated with hippocampal volume in patients with major depressive disorder. <i>Journal of Affective Disorders</i> , <b>2019</b> , 245, 205-212	6.6	8
33	Genes in immune pathways associated with abnormal white matter integrity in first-episode and treatment-naïve patients with schizophrenia. <i>British Journal of Psychiatry</i> , <b>2019</b> , 214, 281-287	5.4	7
32	Minor physical anomalies in patients with schizophrenia in a Chinese population. <i>Psychiatry Research</i> , <b>2012</b> , 200, 223-7	9.9	7
31	Testing the role of genetic variation of the MC4R gene in Chinese population in antipsychotic-induced metabolic disturbance. <i>Science China Life Sciences</i> , <b>2019</b> , 62, 535-543	8.5	6
30	Association of the 3' region of the neuregulin 1 gene with bipolar I disorder in the Chinese Han population. <i>Journal of Affective Disorders</i> , <b>2014</b> , 162, 81-8	6.6	6
29	The functional MMP-9 microsatellite marker is not associated with episodic memory in humans. <i>Psychiatric Genetics</i> , <b>2008</b> , 18, 252	2.9	6
28	Linkage disequilibrium analysis of polymorphisms in the gene for myelin oligodendrocyte glycoprotein in Tourette's syndrome patients from a Chinese sample. <i>American Journal of Medical Genetics Part A</i> , <b>2004</b> , 124B, 76-80		6
27	Neural substrate of unrelenting negative symptoms in schizophrenia: a longitudinal resting-state fMRI study. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , <b>2018</b> , 268, 641-651	5.1	6
26	Distinguishable brain networks relate disease susceptibility to symptom expression in schizophrenia. <i>Human Brain Mapping</i> , <b>2018</b> , 39, 3503-3515	5.9	5
25	Genes in the serotonin pathway are associated with bipolar affective disorder in a Han Chinese population. <i>Neuroscience Bulletin</i> , <b>2014</b> , 30, 33-42	4.3	5
24	Cortical surface area correlates with STON2 gene Ser307Pro polymorphism in first-episode treatment-naïve patients with schizophrenia. <i>PLoS ONE</i> , <b>2013</b> , 8, e64090	3.7	5
23	Interaction among genes influencing ethanol metabolism and sex is association with alcohol use disorders in a Tibet population. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , <b>2010</b> , 153B, 561-569	3.5	5
22	Lower regional grey matter in alcohol use disorders: evidence from a voxel-based meta-analysis. <i>BMC Psychiatry</i> , <b>2021</b> , 21, 247	4.2	5
21	The tree shrew cerebellum atlas: Systematic nomenclature, neurochemical characterization, and afferent projections. <i>Journal of Comparative Neurology</i> , <b>2018</b> , 526, 2744-2775	3.4	5
20	Gender-specific interactions between alcohol metabolism genes and severity of quantitative alcohol-related-traits in a Tibetan population. <i>Neuroscience Letters</i> , <b>2011</b> , 495, 22-5	3.3	4
19	Structural variants in the Chinese population and their impact on phenotypes, diseases and population adaptation. <i>Nature Communications</i> , <b>2021</b> , 12, 6501	17.4	3
18	Revisiting tandem repeats in psychiatric disorders from perspectives of genetics, physiology, and brain evolution. <i>Molecular Psychiatry</i> , <b>2021</b> ,	15.1	3
17	No NRG1 V266L in Chinese patients with schizophrenia. <i>Psychiatric Genetics</i> , <b>2011</b> , 21, 47-9	2.9	2



16	Selective serotonin reuptake inhibitors in anorexia nervosa. <i>Lancet, The</i> , <b>1997</b> , 350, 1033-1034	4.0	2
15	A plasma metabolomics study suggests alteration of multiple metabolic pathways in patients with bipolar disorder. <i>Psychiatry Research</i> , <b>2021</b> , 299, 113880	9.9	2
14	Contribution of genes in the GABAergic pathway to bipolar disorder and its executive function deficit in the Chinese Han population. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , <b>2018</b> , 177, 50-67	3.5	2
13	Interaction Between Variations in Dopamine D2 and Serotonin 2A Receptor is Associated with Short-Term Response to Antipsychotics in Schizophrenia. <i>Neuroscience Bulletin</i> , <b>2019</b> , 35, 1102-1105	4.3	1
12	A Comprehensive Analysis of the CaMK2A Gene and Susceptibility to Alzheimer's Disease in the Han Chinese Population. <i>Frontiers in Aging Neuroscience</i> , <b>2019</b> , 11, 84	5.3	1
11	A Prospective Twin Registry in Southwestern China (TRiSC): exploring the effects of genetic and environmental factors on cognitive and behavioral development and mental health wellbeing in children and adolescents. <i>Twin Research and Human Genetics</i> , <b>2009</b> , 12, 312-9	2.2	1
10	The interactions between childhood adversities and recent stress were associated with early-adulthood depression among Chinese undergraduate students. <i>Depression and Anxiety</i> , <b>2021</b> , 38, 961-971	8.4	1
9	A brief web-based screening plus emotional-disorder health education was associated with improvement of mental health awareness and service-seeking attitudes among patients seeking nonpsychiatric clinical services in China. <i>Depression and Anxiety</i> , <b>2021</b> , 38, 571-587	8.4	1
8	Urbanicity and Psychosis. <i>JAMA Psychiatry</i> , <b>2018</b> , 75, 1301	14.5	1
7	Auditory event-related potentials, neurocognition, and global functioning in drug naïve first-episode schizophrenia and bipolar disorder. <i>Psychological Medicine</i> , <b>2021</b> , 1-10	6.9	1
6	Alterations in CRY2 and PER3 gene expression associated with thalamic-limbic connectivity structural abnormalities in patients with bipolar depression or unipolar depression. <i>Journal of Affective Disorders</i> , <b>2022</b> , 298, 472-480	6.6	0
5	The Gender-Sensitive Social Risk Factors for Internet Addiction in College Undergraduate Students. <i>Psychiatry Investigation</i> , <b>2021</b> , 18, 636-644	3.1	0
4	Comprehensive and integrative analyses identify TYW5 as a schizophrenia risk gene.. <i>BMC Medicine</i> , <b>2022</b> , 20, 169	11.4	0
3	Functional Striatal Abnormalities: A Distinct Brain Signature of Schizophrenia. <i>Neuroscience Bulletin</i> , <b>2021</b> , 37, 284-286	4.3	
2	Generation and characterization of human-derived iPSC lines from one pair of dizygotic twins discordant for schizophrenia.. <i>Stem Cell Research</i> , <b>2022</b> , 60, 102710	1.6	
1	Correlation between reduced telomere length and behavioural and emotional problems in left-behind children in a rural area in China.. <i>Psychoneuroendocrinology</i> , <b>2022</b> , 140, 105732	5	