

Frank de Vries

List of Publications by Citations

Source: <https://exaly.com/author-pdf/9018462/frank-de-vries-publications-by-citations.pdf>

Version: 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

232
papers

5,897
citations

46
h-index

66
g-index

273
ext. papers

6,846
ext. citations

5
avg, IF

5.68
L-index

#	Paper	IF	Citations
232	Fracture risk with intermittent high-dose oral glucocorticoid therapy. <i>Arthritis and Rheumatism</i> , 2007 , 56, 208-14		209
231	Epidemiology of fractures in the United Kingdom 1988-2012: Variation with age, sex, geography, ethnicity and socioeconomic status. <i>Bone</i> , 2016 , 87, 19-26	4.7	196
230	Use of statins and risk of fractures. <i>JAMA - Journal of the American Medical Association</i> , 2001 , 285, 1850-57.4	5.4	190
229	Risk of fracture after bariatric surgery in the United Kingdom: population based, retrospective cohort study. <i>BMJ, The</i> , 2012 , 345, e5085	5.9	132
228	Antidepressant prescribing in five European countries: application of common definitions to assess the prevalence, clinical observations, and methodological implications. <i>European Journal of Clinical Pharmacology</i> , 2014 , 70, 849-57	2.8	110
227	Severity of obstructive airway disease and risk of osteoporotic fracture. <i>European Respiratory Journal</i> , 2005 , 25, 879-84	13.6	110
226	Risk of hip/femur fracture after stroke: a population-based case-control study. <i>Stroke</i> , 2009 , 40, 3281-5	6.7	107
225	Risk of lactic acidosis or elevated lactate concentrations in metformin users with renal impairment: a population-based cohort study. <i>Diabetes Care</i> , 2014 , 37, 2218-24	14.6	102
224	High risk of drug-induced microscopic colitis with concomitant use of NSAIDs and proton pump inhibitors. <i>Alimentary Pharmacology and Therapeutics</i> , 2016 , 43, 1004-13	6.1	101
223	The risk of colorectal cancer in patients with type 2 diabetes: associations with treatment stage and obesity. <i>Diabetes Care</i> , 2015 , 38, 495-502	14.6	94
222	The epidemiology of extra-articular manifestations in ankylosing spondylitis: a population-based matched cohort study. <i>Annals of the Rheumatic Diseases</i> , 2015 , 74, 1373-8	2.4	89
221	The risk of fracture in patients with multiple sclerosis: the UK general practice research database. <i>Journal of Bone and Mineral Research</i> , 2011 , 26, 2271-9	6.3	84
220	Use of proton pump inhibitors and risk of hip/femur fracture: a population-based case-control study. <i>Osteoporosis International</i> , 2011 , 22, 903-10	5.3	79
219	Reporting to Improve Reproducibility and Facilitate Validity Assessment for Healthcare Database Studies V1.0. <i>Pharmacoepidemiology and Drug Safety</i> , 2017 , 26, 1018-1032	2.6	78
218	Fracture risk in patients receiving acid-suppressant medication alone and in combination with bisphosphonates. <i>Osteoporosis International</i> , 2009 , 20, 1989-98	5.3	78
217	Stability of Blood Eosinophils in Patients with Chronic Obstructive Pulmonary Disease and in Control Subjects, and the Impact of Sex, Age, Smoking, and Baseline Counts. <i>American Journal of Respiratory and Critical Care Medicine</i> , 2017 , 195, 1402-1404	10.2	77
216	Use of insulin and insulin analogs and risk of cancer - systematic review and meta-analysis of observational studies. <i>Current Drug Safety</i> , 2013 , 8, 333-48	1.4	77

215	Remission of Type 2 Diabetes Mellitus in Patients After Different Types of Bariatric Surgery: A Population-Based Cohort Study in the United Kingdom. <i>JAMA Surgery</i> , 2015 , 150, 1126-33	5.4	76
214	Use of antidepressant drugs and risk of osteoporotic and non-osteoporotic fractures. <i>Bone</i> , 2010 , 47, 604-9	4.7	76
213	Mortality in British hip fracture patients, 2000-2010: a population-based retrospective cohort study. <i>Bone</i> , 2014 , 66, 171-7	4.7	75
212	Exposure to benzodiazepines (anxiolytics, hypnotics and related drugs) in seven European electronic healthcare databases: a cross-national descriptive study from the PROTECT-EU Project. <i>Pharmacoepidemiology and Drug Safety</i> , 2016 , 25 Suppl 1, 56-65	2.6	73
211	The effect of body mass index on the risk of post-operative complications during the 6 months following total hip replacement or total knee replacement surgery. <i>Osteoarthritis and Cartilage</i> , 2014 , 22, 918-27	6.2	72
210	Use of anti-depressants and the risk of fracture of the hip or femur. <i>Osteoporosis International</i> , 2009 , 20, 1705-13	5.3	72
209	HIV infection and its association with an excess risk of clinical fractures: a nationwide case-control study. <i>Journal of Acquired Immune Deficiency Syndromes (1999)</i> , 2014 , 66, 90-5	3.1	69
208	Causes of death in patients with multiple sclerosis and matched referent subjects: a population-based cohort study. <i>European Journal of Neurology</i> , 2012 , 19, 1007-14	6	68
207	Antipsychotic use and the risk of hip/femur fracture: a population-based case-control study. <i>Osteoporosis International</i> , 2009 , 20, 1499-506	5.3	65
206	Use of beta-blockers and the risk of hip/femur fracture in the United Kingdom and The Netherlands. <i>Calcified Tissue International</i> , 2007 , 80, 69-75	3.9	61
205	Use of inhaled and oral glucocorticoids, severity of inflammatory disease and risk of hip/femur fracture: a population-based case-control study. <i>Journal of Internal Medicine</i> , 2007 , 261, 170-7	10.8	59
204	Statin use and reduced cancer-related mortality. <i>New England Journal of Medicine</i> , 2013 , 368, 576-7	59.2	58
203	Timing of acute myocardial infarction in patients undergoing total hip or knee replacement: a nationwide cohort study. <i>Archives of Internal Medicine</i> , 2012 , 172, 1229-35		56
202	Use of dipeptidyl peptidase-4 inhibitors for type 2 diabetes mellitus and risk of fracture. <i>Bone</i> , 2014 , 68, 124-30	4.7	55
201	Ankylosing spondylitis and risk of ischaemic heart disease: a population-based cohort study. <i>Annals of the Rheumatic Diseases</i> , 2016 , 75, 203-9	2.4	54
200	Potential impact of benzodiazepine use on the rate of hip fractures in five large European countries and the United States. <i>Calcified Tissue International</i> , 2012 , 91, 24-31	3.9	53
199	Lithium use and the risk of fractures. <i>Bone</i> , 2007 , 40, 1252-8	4.7	53
198	Ethnic and geographic variations in the epidemiology of childhood fractures in the United Kingdom. <i>Bone</i> , 2016 , 85, 9-14	4.7	53

197	Reanalysis of two studies with contrasting results on the association between statin use and fracture risk: the General Practice Research Database. <i>International Journal of Epidemiology</i> , 2006 , 35, 1301-8	7.8	52
196	Anti-osteoporosis drug prescribing after hip fracture in the UK: 2000-2010. <i>Osteoporosis International</i> , 2015 , 26, 1919-28	5.3	51
195	Risk of fractures in patients with multiple sclerosis: a population-based cohort study. <i>Neurology</i> , 2012 , 78, 1967-73	6.5	49
194	Reporting to Improve Reproducibility and Facilitate Validity Assessment for Healthcare Database Studies V1.0. <i>Value in Health</i> , 2017 , 20, 1009-1022	3.3	47
193	Ankylosing spondylitis confers substantially increased risk of clinical spine fractures: a nationwide case-control study. <i>Osteoporosis International</i> , 2015 , 26, 85-91	5.3	47
192	Trends in oral anti-osteoporosis drug prescription in the United Kingdom between 1990 and 2012: Variation by age, sex, geographic location and ethnicity. <i>Bone</i> , 2017 , 94, 50-55	4.7	47
191	Bone fracture risk is not associated with the use of glucagon-like peptide-1 receptor agonists: a population-based cohort analysis. <i>Calcified Tissue International</i> , 2015 , 97, 104-12	3.9	46
190	Elevated risk of clinical fractures and associated risk factors in patients with systemic lupus erythematosus versus matched controls: a population-based study in the United Kingdom. <i>Osteoporosis International</i> , 2014 , 25, 1275-83	5.3	46
189	Adjustment for continuous confounders: an example of how to prevent residual confounding. <i>Cmaj</i> , 2013 , 185, 401-6	3.5	46
188	Use of beta-2 agonists and risk of hip/femur fracture: a population-based case-control study. <i>Pharmacoepidemiology and Drug Safety</i> , 2007 , 16, 612-9	2.6	46
187	Use of Glucagon-Like-Peptide 1 Receptor Agonists and Risk of Fracture as Compared to Use of Other Anti-hyperglycemic Drugs. <i>Calcified Tissue International</i> , 2015 , 97, 506-15	3.9	45
186	Risk of hypoglycaemia in users of sulphonylureas compared with metformin in relation to renal function and sulphonylurea metabolite group: population based cohort study. <i>BMJ, The</i> , 2016 , 354, i3625	5.9	45
185	Risk of fracture in patients with Parkinson's disease. <i>Osteoporosis International</i> , 2013 , 24, 2283-90	5.3	43
184	Bridging differences in outcomes of pharmacoepidemiological studies: design and first results of the PROTECT project. <i>Current Clinical Pharmacology</i> , 2014 , 9, 130-8	2.5	43
183	Cancer risk among insulin users: comparing analogues with human insulin in the CARING five-country cohort study. <i>Diabetologia</i> , 2017 , 60, 1691-1703	10.3	41
182	Secular trends in fracture incidence in the UK between 1990 and 2012. <i>Osteoporosis International</i> , 2016 , 27, 3197-3206	5.3	40
181	Changes in mortality patterns following total hip or knee arthroplasty over the past two decades: a nationwide cohort study. <i>Arthritis and Rheumatology</i> , 2014 , 66, 311-8	9.5	40
180	CARING (CAnCer Risk and INsulin analogues): the association of diabetes mellitus and cancer risk with focus on possible determinants - a systematic review and a meta-analysis. <i>Current Drug Safety</i> , 2013 , 8, 296-332	1.4	40

179	Prevalence of antibiotic use: a comparison across various European health care data sources. <i>Pharmacoepidemiology and Drug Safety</i> , 2016 , 25 Suppl 1, 11-20	2.6	40
178	Calibration of FRAX \square 3.1 to the Dutch population with data on the epidemiology of hip fractures. <i>Osteoporosis International</i> , 2012 , 23, 861-9	5.3	37
177	Concomitant use of ibuprofen and paracetamol and the risk of major clinical safety outcomes. <i>British Journal of Clinical Pharmacology</i> , 2010 , 70, 429-38	3.8	37
176	Incidence rates and trends of hip/femur fractures in five European countries: comparison using e-healthcare records databases. <i>Calcified Tissue International</i> , 2014 , 94, 580-9	3.9	36
175	Use of organic nitrates and the risk of hip fracture: a population-based case-control study. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2010 , 95, 1924-31	5.6	34
174	Current and Future Incidence and Costs of Osteoporosis-Related Fractures in The Netherlands: Combining Claims Data with BMD Measurements. <i>Calcified Tissue International</i> , 2016 , 98, 235-43	3.9	33
173	Timing of stroke in patients undergoing total hip replacement and matched controls: a nationwide cohort study. <i>Stroke</i> , 2012 , 43, 3225-9	6.7	33
172	Risk of fracture with thiazolidinediones: an individual patient data meta-analysis. <i>Frontiers in Endocrinology</i> , 2013 , 4, 11	5.7	32
171	Incidence of fractures in patients with multiple sclerosis: the Danish National Health Registers. <i>Multiple Sclerosis Journal</i> , 2012 , 18, 622-7	5	32
170	Long-term use of dipeptidyl peptidase-4 inhibitors and risk of fracture: A retrospective population-based cohort study. <i>Diabetes, Obesity and Metabolism</i> , 2017 , 19, 421-428	6.7	31
169	Use of metformin and survival of diabetic women with breast cancer. <i>Current Drug Safety</i> , 2013 , 8, 357-63	3.4	31
168	Antiepileptic drug use in seven electronic health record databases in Europe: a methodologic comparison. <i>Epilepsia</i> , 2014 , 55, 666-673	6.4	30
167	The use of incretins and fractures - a meta-analysis on population-based real life data. <i>British Journal of Clinical Pharmacology</i> , 2017 , 83, 923-926	3.8	30
166	The accuracy of date of death recording in the Clinical Practice Research Datalink GOLD database in England compared with the Office for National Statistics death registrations. <i>Pharmacoepidemiology and Drug Safety</i> , 2019 , 28, 563-569	2.6	28
165	The epidemiology of fractures in Denmark in 2011. <i>Osteoporosis International</i> , 2016 , 27, 2017-25	5.3	28
164	Long-acting β 2-agonists in adult asthma and the pattern of risk of death and severe asthma outcomes: a study using the GPRD. <i>European Respiratory Journal</i> , 2010 , 36, 494-502	13.6	28
163	Excess risk of hip fractures attributable to the use of antidepressants in five European countries and the USA. <i>Osteoporosis International</i> , 2014 , 25, 847-55	5.3	27
162	Oral bisphosphonate use and total knee/hip implant survival: validation of results in an external population-based cohort. <i>Arthritis and Rheumatology</i> , 2014 , 66, 3233-40	9.5	27

161	Use of thiazolidinediones and risk of osteoporotic fracture: disease or drugs?. <i>Pharmacoepidemiology and Drug Safety</i> , 2012 , 21, 507-14	2.6	26
160	Use of beta2 agonists and risk of acute myocardial infarction in patients with hypertension. <i>British Journal of Clinical Pharmacology</i> , 2008 , 65, 580-6	3.8	26
159	Chlamydia trachomatis and the Risk of Pelvic Inflammatory Disease, Ectopic Pregnancy, and Female Infertility: A Retrospective Cohort Study Among Primary Care Patients. <i>Clinical Infectious Diseases</i> , 2019 , 69, 1517-1525	11.6	25
158	Long-term persistence with anti-osteoporosis drugs after fracture. <i>Osteoporosis International</i> , 2015 , 26, 1831-40	5.3	25
157	The epidemiology of mortality after fracture in England: variation by age, sex, time, geographic location, and ethnicity. <i>Osteoporosis International</i> , 2017 , 28, 161-168	5.3	25
156	Use of dipeptidyl peptidase 4 inhibitors and fracture risk compared to use of other anti-hyperglycemic drugs. <i>Pharmacoepidemiology and Drug Safety</i> , 2015 , 24, 1017-25	2.6	25
155	Fracture risk before and after total hip replacement in patients with osteoarthritis: potential benefits of bisphosphonate use. <i>Arthritis and Rheumatism</i> , 2011 , 63, 992-1001		25
154	Dopaminergic drugs and the risk of hip or femur fracture: a population-based case-control study. <i>Osteoporosis International</i> , 2011 , 22, 2197-204	5.3	24
153	Cancer recording in patients with and without type 2 diabetes in the Clinical Practice Research Datalink primary care data and linked hospital admission data: a cohort study. <i>BMJ Open</i> , 2018 , 8, e020827	2.7	23
152	Individuals With Type 2 Diabetes Mellitus Are at an Increased Risk of Gout But This Is Not Due to Diabetes: A Population-Based Cohort Study. <i>Medicine (United States)</i> , 2015 , 94, e1358	1.8	23
151	A simple score for estimating the long-term risk of fracture in patients with multiple sclerosis. <i>Neurology</i> , 2012 , 79, 922-8	6.5	23
150	Patterns of risk of cancer in patients with metal-on-metal hip replacements versus other bearing surface types: a record linkage study between a prospective joint registry and general practice electronic health records in England. <i>PLoS ONE</i> , 2013 , 8, e65891	3.7	23
149	Predicting the 10-year risk of hip and major osteoporotic fracture in rheumatoid arthritis and in the general population: an independent validation and update of UK FRAX without bone mineral density. <i>Annals of the Rheumatic Diseases</i> , 2016 , 75, 2095-2100	2.4	23
148	The risk of venous thromboembolism in patients with multiple sclerosis: the Clinical Practice Research Datalink. <i>Journal of Thrombosis and Haemostasis</i> , 2014 , 12, 444-51	15.4	22
147	Cardiovascular safety of vildagliptin in patients with type 2 diabetes: A European multi-database, non-interventional post-authorization safety study. <i>Diabetes, Obesity and Metabolism</i> , 2017 , 19, 1473-1478	6.7	21
146	Use of incretin agents and risk of pancreatic cancer: a population-based cohort study. <i>Diabetes, Obesity and Metabolism</i> , 2016 , 18, 258-65	6.7	21
145	Risk of fracture with thiazolidinediones: disease or drugs?. <i>Calcified Tissue International</i> , 2012 , 90, 450-7	3.9	20
144	The risk of fracture in incident multiple sclerosis patients: the Danish National Health Registers. <i>Multiple Sclerosis Journal</i> , 2012 , 18, 1609-16	5	20

143	Safety and efficacy of new oral anticoagulants and low-molecular-weight heparins compared with aspirin in patients undergoing total knee and hip replacements. <i>Pharmacoepidemiology and Drug Safety</i> , 2016 , 25, 1245-1252	2.6	20
142	Impact of cumulative exposure to high-dose oral glucocorticoids on fracture risk in Denmark: a population-based case-control study. <i>Archives of Osteoporosis</i> , 2018 , 13, 30	2.9	19
141	Medication adherence among gout patients initiated allopurinol: a retrospective cohort study in the Clinical Practice Research Datalink (CPRD). <i>Rheumatology</i> , 2018 , 57, 1641-1650	3.9	19
140	Use of proton pump inhibitors and risk of iron deficiency: a population-based case-control study. <i>Journal of Internal Medicine</i> , 2019 , 285, 205-214	10.8	19
139	The Epidemiology of Hip and Major Osteoporotic Fractures in a Dutch Population of Community-Dwelling Elderly: Implications for the Dutch FRAX [®] Algorithm. <i>PLoS ONE</i> , 2015 , 10, e0143800	3.7	19
138	Severity of Diabetes Mellitus and Total Hip or Knee Replacement: A Population-Based Case-Control Study. <i>Medicine (United States)</i> , 2016 , 95, e3739	1.8	19
137	Risk of major bleeding and stroke associated with the use of vitamin K antagonists, nonvitamin K antagonist oral anticoagulants and aspirin in patients with atrial fibrillation: a cohort study. <i>British Journal of Clinical Pharmacology</i> , 2017 , 83, 1844-1859	3.8	17
136	No Decreased Risk of Gastrointestinal Cancers in Users of Metformin in The Netherlands; A Time-Varying Analysis of Metformin Exposure. <i>Cancer Prevention Research</i> , 2017 , 10, 290-297	3.2	17
135	The risk of major and any (non-hip) fragility fracture after hip fracture in the United Kingdom: 2000-2010. <i>Osteoporosis International</i> , 2014 , 25, 2555-63	5.3	17
134	The pattern of risk of myocardial infarction in patients taking asthma medication: a study with the General Practice Research Database. <i>Journal of Hypertension</i> , 2009 , 27, 1485-92	1.9	17
133	Use of Dipeptidyl-Peptidase-4 Inhibitors and the Risk of Pneumonia: A Population-Based Cohort Study. <i>PLoS ONE</i> , 2015 , 10, e0139367	3.7	17
132	The IMI PROTECT project: purpose, organizational structure, and procedures. <i>Pharmacoepidemiology and Drug Safety</i> , 2016 , 25 Suppl 1, 5-10	2.6	17
131	Development and validation of an analytical method using UPLC-MS/MS to quantify everolimus in dried blood spots in the oncology setting. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2018 , 149, 106-113	3.5	16
130	Risk of vertebral and non-vertebral fractures in patients with sarcoidosis: a population-based cohort. <i>Osteoporosis International</i> , 2016 , 27, 1603-1610	5.3	16
129	Influence of metformin intake on the risk of bladder cancer in type 2 diabetes patients. <i>British Journal of Clinical Pharmacology</i> , 2015 , 80, 1464-72	3.8	15
128	Risk of community-acquired pneumonia in chronic obstructive pulmonary disease stratified by smoking status: a population-based cohort study in the United Kingdom. <i>International Journal of COPD</i> , 2017 , 12, 2425-2432	3	15
127	Secular trends in major osteoporotic fractures among 50+ adults in Denmark between 1995 and 2010. <i>Osteoporosis International</i> , 2019 , 30, 2217-2223	5.3	15
126	Use of incretin agents and risk of acute and chronic pancreatitis: A population-based cohort study. <i>Diabetes, Obesity and Metabolism</i> , 2017 , 19, 401-411	6.7	15

125	Data management and data analysis techniques in pharmacoepidemiological studies using a pre-planned multi-database approach: a systematic literature review. <i>Pharmacoepidemiology and Drug Safety</i> , 2015 , 24, 897-905	2.6	15
124	Increase in prophylaxis of glucocorticoid-induced osteoporosis by pharmacist feedback: a randomised controlled trial. <i>Osteoporosis International</i> , 2014 , 25, 385-92	5.3	15
123	The Impact of the Choice of Data Source in Record Linkage Studies Estimating Mortality in Venous Thromboembolism. <i>PLoS ONE</i> , 2016 , 11, e0148349	3.7	15
122	Mortality, causes of death and influence of medication use in patients with systemic lupus erythematosus vs matched controls. <i>Rheumatology</i> , 2021 , 60, 207-216	3.9	15
121	Risk of fracture in patients with muscular dystrophies. <i>Osteoporosis International</i> , 2014 , 25, 509-18	5.3	14
120	Knee arthroplasty and risk of hip fracture: a population-based, case-control study. <i>Calcified Tissue International</i> , 2012 , 90, 144-50	3.9	14
119	Risk of infections in patients with gout: a population-based cohort study. <i>Scientific Reports</i> , 2017 , 7, 14294.9	4.9	14
118	The Netherlands Chlamydia cohort study (NECCST) protocol to assess the risk of late complications following Chlamydia trachomatis infection in women. <i>BMC Infectious Diseases</i> , 2017 , 17, 264	4	14
117	Risk of bladder cancer in patients with diabetes: a retrospective cohort study. <i>BMJ Open</i> , 2015 , 5, e007430	3.0	14
116	Gastrointestinal cancer incidence in type 2 diabetes mellitus; results from a large population-based cohort study in the UK. <i>Cancer Epidemiology</i> , 2018 , 54, 104-111	2.8	14
115	Impact of detection bias on the risk of gastrointestinal cancer and its subsites in type 2 diabetes mellitus. <i>European Journal of Cancer</i> , 2017 , 79, 61-71	7.5	13
114	Statins and Risk of Lower Limb Revision Surgery: The Influence of Differences in Study Design Using Electronic Health Records From the United Kingdom and Denmark. <i>American Journal of Epidemiology</i> , 2016 , 184, 58-66	3.8	13
113	Balance measures for propensity score methods: a clinical example on beta-agonist use and the risk of myocardial infarction. <i>Pharmacoepidemiology and Drug Safety</i> , 2011 , 20, 1130-7	2.6	13
112	Use of thiazolidinediones and the risk of elective hip or knee replacement: a population based case-control study. <i>British Journal of Clinical Pharmacology</i> , 2016 , 81, 370-8	3.8	13
111	Risk of gastrointestinal bleeding in patients undergoing total hip or knee replacement compared with matched controls: a nationwide cohort study. <i>American Journal of Gastroenterology</i> , 2013 , 108, 1277-85	0.7	12
110	Fracture rate in patients with myasthenia gravis: the general practice research database. <i>Osteoporosis International</i> , 2013 , 24, 467-76	5.3	12
109	Blood Eosinophil Counts, Withdrawal of Inhaled Corticosteroids and Risk of COPD Exacerbations and Mortality in the Clinical Practice Research Datalink (CPRD). <i>COPD: Journal of Chronic Obstructive Pulmonary Disease</i> , 2019 , 16, 152-159	2	11
108	Use of high-dose intermittent systemic glucocorticoids and the risk of fracture in patients with chronic obstructive pulmonary disease. <i>Bone</i> , 2018 , 110, 238-243	4.7	11

107	Use of systemic glucocorticoids and the risk of major osteoporotic fractures in patients with sarcoidosis. <i>Osteoporosis International</i> , 2017 , 28, 2859-2866	5.3	11
106	Quality of INR control and outcomes following venous thromboembolism. <i>Clinical and Applied Thrombosis/Hemostasis</i> , 2012 , 18, 370-8	3.3	11
105	Use of thiazolidinediones and risk of bladder cancer: disease or drugs?. <i>Current Drug Safety</i> , 2013 , 8, 364-70	4.0	11
104	Concomitant use of oral glucocorticoids and proton pump inhibitors and risk of osteoporotic fractures among patients with rheumatoid arthritis: a population-based cohort study. <i>Annals of the Rheumatic Diseases</i> , 2020 ,	2.4	11
103	Proton pump inhibitors, fracture risk and selection bias: three studies, same database, two answers. <i>Osteoporosis International</i> , 2011 , 22, 1641-2	5.3	10
102	Insulin glargine use and breast cancer risk: Associations with cumulative exposure. <i>Acta Oncologica</i> , 2016 , 55, 851-8	3.2	10
101	Impact of varying outcomes and definitions of suicidality on the associations of antiepileptic drugs and suicidality: comparisons from UK Clinical Practice Research Datalink (CPRD) and Danish national registries (DNR). <i>Pharmacoepidemiology and Drug Safety</i> , 2016 , 25 Suppl 1, 142-55	2.6	10
100	Blood eosinophilia, use of inhaled corticosteroids, and risk of COPD exacerbations and mortality. <i>Pharmacoepidemiology and Drug Safety</i> , 2018 , 27, 1191-1199	2.6	10
99	The association between insulin use and volumetric bone mineral density, bone micro-architecture and bone strength of the distal radius in patients with type 2 diabetes - The Maastricht study. <i>Bone</i> , 2017 , 101, 156-161	4.7	9
98	Risk of myocardial infarction in patients with atrial fibrillation using vitamin K antagonists, aspirin or direct acting oral anticoagulants. <i>British Journal of Clinical Pharmacology</i> , 2017 , 83, 1835-1843	3.8	9
97	Psychometric Validation of the Autism Impact Measure (AIM). <i>Journal of Autism and Developmental Disorders</i> , 2019 , 49, 2559-2570	4.6	9
96	A compilation of research working groups on drug utilisation across Europe. <i>BMC Research Notes</i> , 2014 , 7, 143	2.3	9
95	Pattern of risks of systemic lupus erythematosus among statin users: a population-based cohort study. <i>Annals of the Rheumatic Diseases</i> , 2017 , 76, 1723-1730	2.4	9
94	Pattern of risks of rheumatoid arthritis among patients using statins: A cohort study with the clinical practice research datalink. <i>PLoS ONE</i> , 2018 , 13, e0193297	3.7	9
93	Smoke-free legislation and the incidence of paediatric respiratory infections and wheezing/asthma: interrupted time series analyses in the four UK nations. <i>Scientific Reports</i> , 2015 , 5, 15246	4.9	8
92	COX-2-selective NSAIDs and risk of hip or knee replacements: a population-based case-control study. <i>Calcified Tissue International</i> , 2012 , 91, 387-94	3.9	8
91	Use of biguanides and the risk of colorectal cancer: a register-based cohort study. <i>Current Drug Safety</i> , 2013 , 8, 349-56	1.4	8
90	Diabetes and fractures: new evidence of atypical femoral fractures?. <i>Osteoporosis International</i> , 2020 , 31, 447-455	5.3	8

89	Increased risk of all-cause mortality associated with domperidone use in Parkinson [®] patients: a population-based cohort study in the UK. <i>British Journal of Clinical Pharmacology</i> , 2018 , 84, 2551-2561	3.8	8
88	Association of type 2 diabetes mellitus with self-reported knee pain and clinical knee osteoarthritis: The Maastricht Study. <i>Diabetes and Metabolism</i> , 2018 , 44, 296-299	5.4	7
87	Association between use of antidepressants or benzodiazepines and the risk of subsequent fracture among those aged 65+ in the Netherlands. <i>Osteoporosis International</i> , 2018 , 29, 2477-2485	5.3	7
86	Risk of cataract and glaucoma in patients with multiple sclerosis. <i>Multiple Sclerosis Journal</i> , 2012 , 18, 628-38	5	7
85	Psychostimulants/Atomoxetine and Serious Cardiovascular Events in Children with ADHD or Autism Spectrum Disorder. <i>CNS Drugs</i> , 2020 , 34, 93-101	6.7	7
84	Inpatient drug utilization in Europe: nationwide data sources and a review of publications on a selected group of medicines (PROTECT project). <i>Basic and Clinical Pharmacology and Toxicology</i> , 2015 , 116, 201-11	3.1	6
83	Risk of a first-ever acute myocardial infarction and all-cause mortality with sulphonylurea treatment: A population-based cohort study. <i>Diabetes, Obesity and Metabolism</i> , 2018 , 20, 1056-1060	6.7	6
82	Comparability of the age and sex distribution of the UK Clinical Practice Research Datalink and the total Dutch population. <i>Pharmacoepidemiology and Drug Safety</i> , 2016 , 25, 1460-1464	2.6	6
81	Secular trends in major osteoporotic fractures among 50+ adults in Denmark between 1995 and 2010. <i>Archives of Osteoporosis</i> , 2018 , 13, 91	2.9	6
80	Gastric acid-suppressive agents and risk of Clostridium difficile-associated disease. <i>JAMA - Journal of the American Medical Association</i> , 2006 , 295, 2599; author reply 2600-1	27.4	6
79	On epidemiology of fractures and variation with age and ethnicity. <i>Bone</i> , 2016 , 93, 230-231	4.7	5
78	Economic impact of Patient [®] Own Medication use during hospitalisation: a multicentre pre-post implementation study. <i>International Journal of Clinical Pharmacy</i> , 2019 , 41, 1658-1665	2.3	5
77	Risk of fracture in patients with Charcot-Marie-Tooth disease. <i>Muscle and Nerve</i> , 2014 , 50, 919-24	3.4	5
76	Use of inhaled corticosteroids and the risk of non-fatal acute myocardial infarction. <i>Journal of Hypertension</i> , 2008 , 26, 124-9	1.9	5
75	Metformin use in type 2 diabetic patients is not associated with lower arterial stiffness: the Maastricht Study. <i>Journal of Hypertension</i> , 2019 , 37, 365-371	1.9	5
74	Disease burden of knee osteoarthritis patients with a joint replacement compared to matched controls: a population-based analysis of a Dutch medical claims database. <i>Osteoarthritis and Cartilage</i> , 2018 , 26, 202-210	6.2	5
73	Glucose-Lowering Drugs and Fracture Risk-a Systematic Review. <i>Current Osteoporosis Reports</i> , 2020 , 18, 737-758	5.4	4
72	Incidence of subsequent fractures in the UK between 1990 and 2012 among individuals 50 years or older. <i>Osteoporosis International</i> , 2018 , 29, 2469-2475	5.3	4

71	The Association Between β -Blocker Use and Cardiorespiratory Fitness: The Maastricht Study. <i>Journal of Cardiovascular Pharmacology and Therapeutics</i> , 2019 , 24, 37-45	2.6	4
70	External Validation Of The Updated ADO Score In COPD Patients From The Birmingham COPD Cohort. <i>International Journal of COPD</i> , 2019 , 14, 2395-2407	3	4
69	Five-year fracture risk estimation in patients with Parkinson's disease. <i>Bone</i> , 2013 , 56, 266-70	4.7	4
68	Pioglitazone and bladder cancer: two studies, same database, two answers. <i>British Journal of Clinical Pharmacology</i> , 2013 , 76, 484-5	3.8	4
67	The Association of Oral Bisphosphonate Use With Mortality Risk Following a Major Osteoporotic Fracture in the United Kingdom: Population-Based Cohort Study. <i>Journal of the American Medical Directors Association</i> , 2020 , 21, 811-816	5.9	4
66	Use of Sodium-Glucose Co-Transporter-2-Inhibitors (SGLT2-Is) and Risk of Lower Limb Amputation. <i>Current Drug Safety</i> , 2021 , 16, 62-72	1.4	4
65	All-cause mortality with current and past use of antidepressants or benzodiazepines after major osteoporotic and hip fracture. <i>Osteoporosis International</i> , 2019 , 30, 573-581	5.3	3
64	The Impact of Dose and Simultaneous Use of Acid-Reducing Agents on the Effectiveness of Vemurafenib in Metastatic BRAF V600 Mutated Melanoma: a Retrospective Cohort Study. <i>Targeted Oncology</i> , 2018 , 13, 363-370	5	3
63	Two-year persistence with teriparatide improved significantly after introduction of an educational and motivational support program. <i>Osteoporosis International</i> , 2019 , 30, 1837-1844	5.3	3
62	Risk of fracture in patients with Guillain-Barré syndrome. <i>Osteoporosis International</i> , 2014 , 25, 1845-51	5.3	3
61	THU0069 The Epidemiology of Extra-Articular Manifestations in Ankylosing Spondylitis: A Population-Based Matched Cohort Study. <i>Annals of the Rheumatic Diseases</i> , 2014 , 73, 200.3-200	2.4	3
60	Prescribing Errors With Low-Molecular-Weight Heparins. <i>Journal of Patient Safety</i> , 2021 , 17, e587-e592	1.9	3
59	Harmful effects of proton pump inhibitors: discrepancies between observational studies and randomized clinical trials. <i>JAMA Internal Medicine</i> , 2013 , 173, 1559	11.5	3
58	Prolonged outpatient vitamin K antagonist use and risk of venous thromboembolism in patients undergoing total hip or knee replacement. <i>Journal of Thrombosis and Haemostasis</i> , 2013 , 11, 642-50	15.4	3
57	Risk of fractures in patients with multiple sclerosis: a population-based cohort study. <i>Neurology</i> , 2012 , 79, 1934-5	6.5	3
56	Pregnancies and Time to Pregnancy in Women With and Without a Previous Chlamydia trachomatis Infection. <i>Sexually Transmitted Diseases</i> , 2020 , 47, 739-747	2.4	3
55	The stability of the ADO score among UK COPD patients from The Health Improvement Network. <i>ERJ Open Research</i> , 2020 , 6,	3.5	3
54	Increased Risk of Falls, Fall-related Injuries and Fractures in People with Type 1 and Type 2 Diabetes - A Nationwide Cohort Study. <i>Current Drug Safety</i> , 2021 , 16, 52-61	1.4	3

53	Trends in moderate and severe exacerbations among COPD patients in the UK from 2005 to 2013. <i>Respiratory Medicine</i> , 2018 , 144, 1-6	4.6	3
52	Low-dose oral glucocorticoid therapy and risk of osteoporotic fractures in patients with rheumatoid arthritis: a cohort study using the Clinical Practice Research Datalink. <i>Rheumatology</i> , 2021 ,	3.9	3
51	Impact of the definition of osteoarthritis and of the timing of its onset on the association between type 2 diabetes mellitus and osteoarthritis: Clinical Practice Research Datalink. <i>Diabetes Research and Clinical Practice</i> , 2019 , 148, 240-248	7.4	2
50	Risk of vascular disease with gout: overadjustment of the statistical analyses?. <i>Annals of the Rheumatic Diseases</i> , 2015 , 74, e9	2.4	2
49	No decrease in fracture risk despite 15 years of treatment evolution for multiple myeloma patients: A Danish nationwide case-control study. <i>Bone</i> , 2020 , 134, 115299	4.7	2
48	The risk of new fragility fractures in patients with chronic kidney disease and hip fracture-a population-based cohort study in the UK. <i>Osteoporosis International</i> , 2020 , 31, 1487-1497	5.3	2
47	Obstructive sleep apnea and the risk of gout: a population-based case-control study. <i>Arthritis Research and Therapy</i> , 2020 , 22, 92	5.7	2
46	Factors Associated with Antihypertensive Medication Non-Adherence: A Cross-Sectional Study Among Lebanese Hypertensive Adults. <i>Patient Preference and Adherence</i> , 2020 , 14, 663-673	2.4	2
45	Association between vildagliptin and risk of angioedema, foot ulcers, skin lesions, hepatic toxicity, and serious infections in patients with type 2 diabetes mellitus: A European multidatabase, noninterventional, postauthorization safety study. <i>Endocrinology, Diabetes and Metabolism</i> , 2019 , 2, e00084	2.7	2
44	The difficulty in evaluating all findings in study on use of macrolides in mother and child and risk of infantile hypertrophic pyloric stenosis. <i>BMJ, The</i> , 2014 , 349, g5201	5.9	2
43	Thiazolidinediones and cancer: duplicate publication bias?. <i>Oncologist</i> , 2013 , 18, 1147	5.7	2
42	Acid suppressants and hip fracture: duplicate publication bias?. <i>Bone</i> , 2011 , 49, 920; author reply 921	4.7	2
41	Liquid Chromatography-Tandem Mass Spectrometry to Monitor Unbound and Total Ceftriaxone in Serum of Critically Ill Patients. <i>Current Clinical Pharmacology</i> , 2021 , 16, 341-349	2.5	2
40	Incidence, hospitalization and mortality and their changes over time in people with a first ever diabetic foot ulcer. <i>Diabetic Medicine</i> , 2021 , e14725	3.5	2
39	Validation of an analytical method using HPLC-MS/MS to quantify osimertinib in human plasma and supplementary stability results. <i>Biomedical Chromatography</i> , 2020 , 34, e4771	1.7	2
38	The risk of community-acquired pneumonia in children using gastric acid suppressants. <i>European Respiratory Journal</i> , 2021 , 58,	13.6	2
37	Thiazolidinediones and Glucagon-Like Peptide-1 Receptor Agonists and the Risk of Nonalcoholic Fatty Liver Disease: A Cohort Study. <i>Hepatology</i> , 2021 , 74, 2467-2477	11.2	2
36	Adjusting for unmeasured confounding using validation data: Simplified two-stage calibration for survival and dichotomous outcomes. <i>Statistics in Medicine</i> , 2019 , 38, 2719-2734	2.3	1

35	Drug utilization in the Maastricht Study: A comparison with nationwide data. <i>Medicine (United States)</i> , 2020 , 99, e18524	1.8	1
34	Letter regarding "Cognitive outcomes in meningioma patients undergoing surgery: individual changes over time and predictors of late cognitive functioning". <i>Neuro-Oncology</i> , 2020 , 22, 580-581	1	1
33	Effects of colchicine on risk of cardiovascular events and mortality among patients with gout: the effect of index date selection?. <i>Annals of the Rheumatic Diseases</i> , 2016 , 75, e26	2.4	1
32	Mortality risk in atrial fibrillation: the role of aspirin, vitamin K and non-vitamin K antagonists. <i>International Journal of Clinical Pharmacy</i> , 2019 , 41, 1536-1544	2.3	1
31	Drug-drug interactions with tyrosine-kinase inhibitors. <i>Lancet Oncology, The</i> , 2014 , 15, e416	21.7	1
30	THU0090 Cardiovascular Morbidity in Patients with Ankylosing Spondylitis: A Population-Based Cohort Study. <i>Annals of the Rheumatic Diseases</i> , 2014 , 73, 207.3-208	2.4	1
29	Oral Abstracts 3: Adolescent and Young Adult * O13. Hypermobility is a Risk Factor for Musculoskeletal Pain in Adolescence: Findings From a Prospective Cohort Study. <i>Rheumatology</i> , 2013 , 52, i33-i43	3.9	1
28	Experiences with Lean Six Sigma as improvement strategy to reduce parenteral medication administration errors and associated potential risk of harm. <i>BMJ Quality Improvement Reports</i> , 2017 , 6,		1
27	Identification of antithrombotic drugs related to total joint replacement using anonymised free-text notes: a search strategy in the Clinical Practice Research Datalink. <i>BMJ Open</i> , 2015 , 5, e009017 ³		1
26	Effective writing and dealing with reviewers. <i>Journal of Clinical Epidemiology</i> , 2014 , 67, 830	5.7	1
25	Use of hormonal contraceptives and risk of HIV-1 transmission. <i>Lancet Infectious Diseases, The</i> , 2012 , 12, 508-9; author reply 510-1	25.5	1
24	Severity of obstructive airway disease and risk of fracture. <i>Bone</i> , 2013 , 52, 711	4.7	1
23	Inverse association between duration of use of acid-suppressive medications and fracture risk. <i>Archives of Internal Medicine</i> , 2011 , 171, 1044; author reply 1044		1
22	His452Tyr 5-HT polymorphism and intravaginal ejaculation latency time in Dutch men with lifelong premature ejaculation. <i>International Journal of Impotence Research</i> , 2021 ,	2.3	1
21	Risperidone versus aripiprazole fracture risk in children and adolescents with autism spectrum disorders. <i>Autism Research</i> , 2021 , 14, 1800-1814	5.1	1
20	Response to: Correspondence on Concomitant use of oral glucocorticoids and proton pump inhibitors and risk of osteoporotic fractures among patients with rheumatoid arthritis: a population-based cohort study by Zheng. <i>Annals of the Rheumatic Diseases</i> , 2021 ,	2.4	1
19	The use of oral glucocorticoids and the risk of major osteoporotic fracture in patients with myasthenia gravis. <i>Osteoporosis International</i> , 2021 , 1	5.3	0
18	C-reactive protein as a biomarker of response to inhaled corticosteroids among patients with COPD. <i>Pulmonary Pharmacology and Therapeutics</i> , 2020 , 60, 101870	3.5	0

17	Inconsistencies in Reporting Studies of Lactic Acidosis. <i>JAMA Internal Medicine</i> , 2019 , 179, 455-456	11.5
16	Use of parenteral glucocorticoids and the risk of new onset type 2 diabetes mellitus: A case-control study. <i>Diabetes Research and Clinical Practice</i> , 2018 , 139, 100-106	7.4
15	OP0017 Medication Taking Behaviour of Allopurinol Treatment among Patients with Gout: A Retrospective Cohort Study in The Clinical Practice Research Datalink (CPRD). <i>Annals of the Rheumatic Diseases</i> , 2016 , 75, 59.2-59	2.4
14	Comments on "Use of metformin and risk of kidney cancer in patients with type 2 diabetes", Chin-Hsiao Tseng, Eur J Cancer, 2016, No. 52, pp. 19-25. <i>European Journal of Cancer</i> , 2016 , 61, 157-8	7.5
13	THU0334 Severity of Diabetes Mellitus and Risk of Total Hip or Knee Replacement: A Population Based Case-Control Study. <i>Annals of the Rheumatic Diseases</i> , 2015 , 74, 316.2-316	2.4
12	AB1108 Identification of Antithrombotic Drugs Related to Total Joint Replacement Using Anonymised Free Text Notes. <i>Annals of the Rheumatic Diseases</i> , 2015 , 74, 1271.2-1271	2.4
11	SAT0333 Use of Glitazones and The Risk of Elective Hip Or Knee Replacement: A Population Based Case-Control Study. <i>Annals of the Rheumatic Diseases</i> , 2015 , 74, 779.3-780	2.4
10	Use of thiazolidinediones and risk of osteoporotic fracture: Disease or drugs?. <i>Bone</i> , 2010 , 47, S372-S373	4.7
9	Acid suppressants, fracture, and statistical adjustments for body mass index, smoking status, and calcium/vitamin D intake. <i>American Journal of Gastroenterology</i> , 2011 , 106, 2051	0.7
8	Interpretation of the association between beta-blockers and hip/femur fracture. <i>American Journal of Nephrology</i> , 2012 , 35, 101-2	4.6
7	P.2.a.032 Resource use in depressed patients with sleep disturbance. <i>European Neuropsychopharmacology</i> , 2009 , 19, S378-S379	1.2
6	Is there a need for review-a-thons?. <i>Lancet Rheumatology, The</i> , 2020 , 2, e205	14.2
5	Letter to the Editor. Cognitive performance of patients with low-grade glioma. <i>Journal of Neurosurgery</i> , 2020 , 1-2	3.2
4	No Decrease in Fracture Risk Despite 15 Years of Treatment Evolution in Multiple Myeloma Patients: A Danish Nationwide Case-Control Study. <i>Blood</i> , 2019 , 134, 5503-5503	2.2
3	Association of concomitant use of acid reducing agents in full-dose vemurafenib users with risk of progression in BRAF V600 mutation-positive unresectable or metastatic melanoma patients: A retrospective cohort study.. <i>Journal of Clinical Oncology</i> , 2017 , 35, 9540-9540	2.2
2	Determinants of treatment modification before and after implementation of the updated 2015 NICE guideline on type 2 diabetes: A retrospective cohort study. <i>Diabetes Research and Clinical Practice</i> , 2021 , 176, 108828	7.4
1	Characteristics of patients with lung cancer in clinical practice and their potential eligibility for clinical trials evaluating tyrosine kinase inhibitors or immune checkpoint inhibitors.. <i>Cancer Epidemiology</i> , 2022 , 78, 102149	2.8