

Jasenka Zubcevic

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/9019664/jasenka-zubcevic-publications-by-year.pdf>

Version: 2024-04-24

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

54
papers

2,159
citations

21
h-index

46
g-index

62
ext. papers

2,711
ext. citations

5.6
avg, IF

4.71
L-index

#	Paper	IF	Citations
54	Identification of a Gut Commensal That Compromises the Blood Pressure-Lowering Effect of Ester Angiotensin-Converting Enzyme Inhibitors.. <i>Hypertension</i> , 2022 , 101161HYPERTENSIONAHA12118711	8.5	4
53	Tumor Necrosis Factor Alpha and the Gastrointestinal Epithelium: Implications for the Gut-Brain Axis and Hypertension. <i>Cellular and Molecular Neurobiology</i> , 2021 , 1	4.6	0
52	Central Administration of Hydrogen Sulfide Donor NaHS Reduces Iba1-Positive Cells in the PVN and Attenuates Rodent Angiotensin II Hypertension. <i>Frontiers in Neuroscience</i> , 2021 , 15, 690919	5.1	2
51	The importance of bone marrow and the immune system in driving increases in blood pressure and sympathetic nerve activity in hypertension. <i>Experimental Physiology</i> , 2020 , 105, 1815-1826	2.4	5
50	Gastrointestinal dysbiosis following diethylhexyl phthalate exposure in zebrafish (Danio rerio): Altered microbial diversity, functionality, and network connectivity. <i>Environmental Pollution</i> , 2020 , 265, 114496	9.3	13
49	Genetic ablation of bone marrow beta-adrenergic receptors in mice modulates miRNA-transcriptome networks of neuroinflammation in the paraventricular nucleus. <i>Physiological Genomics</i> , 2020 , 52, 169-177	3.6	3
48	Reduced responsiveness of bone marrow hematopoietic cells to sympathetic activation is protective against high fat diet-induced obesity and gut dysbiosis. <i>FASEB Journal</i> , 2020 , 34, 1-1	0.9	
47	Sub-diaphragmatic vagal nerve stimulation alleviates rodent hypertension associated with gut dysbiosis and reduced serotonergic vagal afferent signaling. <i>FASEB Journal</i> , 2020 , 34, 1-1	0.9	
46	Genetic ablation of bone marrow beta-adrenergic receptors alters miRNA-transcriptome networks for microglia activation and inflammation in the paraventricular nucleus of the hypothalamus. <i>FASEB Journal</i> , 2020 , 34, 1-1	0.9	
45	Nicotine exposure during pregnancy alters the maternal gut microbiome and both cecal and plasma short chain fatty acids in Sprague Dawley rats.. <i>FASEB Journal</i> , 2020 , 34, 1-1	0.9	2
44	Ghrelin Signaling Affects Feeding Behavior, Metabolism, and Memory through the Vagus Nerve. <i>Current Biology</i> , 2020 , 30, 4510-4518.e6	6.3	15
43	Pulmonary arterial hypertension-associated changes in gut pathology and microbiota. <i>ERJ Open Research</i> , 2020 , 6,	3.5	11
42	Gut microbiota and neuroinflammation in pathogenesis of hypertension: A potential role for hydrogen sulfide. <i>Pharmacological Research</i> , 2020 , 153, 104677	10.2	15
41	Attenuated Amiloride-Sensitive Current and Augmented Calcium-Activated Chloride Current in Marsh Rice Rat (<i>Oryzomys palustris</i>) Airways. <i>IScience</i> , 2019 , 19, 737-748	6.1	3
40	Impaired Autonomic Nervous System-Microbiome Circuit in Hypertension. <i>Circulation Research</i> , 2019 , 125, 104-116	15.7	47
39	Elevated bone marrow sympathetic drive precedes systemic inflammation in angiotensin II hypertension. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2019 , 317, H279-H289	5.2	20
38	Comparison of Isoflurane, Ketamine-Dexmedetomidine, and Ketamine-Xylazine for General Anesthesia during Oral Procedures in Rice Rats (). <i>Journal of the American Association for Laboratory Animal Science</i> , 2019 , 58, 40-49	1.3	6

37	Sustained Captopril-Induced Reduction in Blood Pressure Is Associated With Alterations in Gut-Brain Axis in the Spontaneously Hypertensive Rat. <i>Journal of the American Heart Association</i> , 2019 , 8, e010721	6	37
36	Central Administration of Hydrogen Sulfide Alleviates Rodent Angiotensin II Hypertension. <i>FASEB Journal</i> , 2019 , 33, 835.18	0.9	
35	Impaired butyrate absorption in the proximal colon, low serum butyrate and diminished central effects of butyrate on blood pressure in spontaneously hypertensive rats. <i>Acta Physiologica</i> , 2019 , 226, e13256	5.6	41
34	Involvement of Neuroinflammation in the Pathogenesis of Monocrotaline-Induced Pulmonary Hypertension. <i>Hypertension</i> , 2018 , 71, 1156-1163	8.5	27
33	Cocaine differentially affects synaptic activity in memory and midbrain areas of female and male rats: an in vivo MEMRI study. <i>Brain Imaging and Behavior</i> , 2018 , 12, 201-216	4.1	7
32	Butyrate regulates inflammatory cytokine expression without affecting oxidative respiration in primary astrocytes from spontaneously hypertensive rats. <i>Physiological Reports</i> , 2018 , 6, e13732	2.6	19
31	MEMRI reveals altered activity in brain regions associated with anxiety, locomotion, and cardiovascular reactivity on the elevated plus maze in the WKY vs SHR rats. <i>Brain Imaging and Behavior</i> , 2018 , 12, 1318-1331	4.1	6
30	Increased Abundance of Lactobacillales in the Colon of Beta-Adrenergic Receptor Knock Out Mouse Is Associated With Increased Gut Bacterial Production of Short Chain Fatty Acids and Reduced IL17 Expression in Circulating CD4 Immune Cells. <i>Frontiers in Physiology</i> , 2018 , 9, 1593	4.6	20
29	Transcriptional networks in rodent models support a role for gut-brain communication in neurogenic hypertension: a review of the evidence. <i>Physiological Genomics</i> , 2017 , 49, 327-338	3.6	8
28	Hypertension-Linked Pathophysiological Alterations in the Gut. <i>Circulation Research</i> , 2017 , 120, 312-323	15.7	247
27	Shifts in the Gut Microbiota Composition Due to Depleted Bone Marrow Beta Adrenergic Signaling Are Associated with Suppressed Inflammatory Transcriptional Networks in the Mouse Colon. <i>Frontiers in Physiology</i> , 2017 , 8, 220	4.6	18
26	A Single Angiotensin II Hypertensive Stimulus Is Associated with Prolonged Neuronal and Immune System Activation in Wistar-Kyoto Rats. <i>Frontiers in Physiology</i> , 2017 , 8, 592	4.6	33
25	Gut-Brain Axis in Regulation of Blood Pressure. <i>Frontiers in Physiology</i> , 2017 , 8, 845	4.6	33
24	Loss of bone marrow adrenergic beta 1 and 2 receptors modifies transcriptional networks, reduces circulating inflammatory factors, and regulates blood pressure. <i>Physiological Genomics</i> , 2016 , 48, 526-363	3.6	16
23	Gut dysbiosis is linked to hypertension. <i>Hypertension</i> , 2015 , 65, 1331-40	8.5	716
22	Involvement of bone marrow cells and neuroinflammation in hypertension. <i>Circulation Research</i> , 2015 , 117, 178-91	15.7	116
21	Optimizing the Microfil Dye Perfusion Technique will Allow for Rat Cerebral Vascular Network Analysis through uCT Image Quantification. <i>FASEB Journal</i> , 2015 , 29, 647.1	0.9	
20	Functional neural-bone marrow pathways: implications in hypertension and cardiovascular disease. <i>Hypertension</i> , 2014 , 63, e129-39	8.5	32

19	Altered inflammatory response is associated with an impaired autonomic input to the bone marrow in the spontaneously hypertensive rat. <i>Hypertension</i> , 2014 , 63, 542-50	8.5	70
18	Nucleus of the solitary tract (pro)renin receptor-mediated antihypertensive effect involves nuclear factor- κ B-cytokine signaling in the spontaneously hypertensive rat. <i>Hypertension</i> , 2013 , 61, 622-7	8.5	31
17	Chronic knockdown of the nucleus of the solitary tract AT1 receptors increases blood inflammatory-endothelial progenitor cell ratio and exacerbates hypertension in the spontaneously hypertensive rat. <i>Hypertension</i> , 2013 , 61, 1328-33	8.5	29
16	Neuroimmune communication in hypertension and obesity: a new therapeutic angle?. <i>Pharmacology & Therapeutics</i> , 2013 , 138, 428-40	13.9	39
15	Dysfunctional brain-bone marrow communication: a paradigm shift in the pathophysiology of hypertension. <i>Current Hypertension Reports</i> , 2013 , 15, 377-89	4.7	21
14	Abstract 606: Reconstitution Of Bone Marrow With WKY Cells Lowers Central/Peripheral Inflammation And Blood Pressure In The SHR. <i>Hypertension</i> , 2013 , 62,	8.5	1
13	Increased Expression of Prorenin Receptor (PRR) in the NTS of Spontaneously Hypertensive Rats (SHR) May Be A Compensatory Mechanism of Hypertension. <i>FASEB Journal</i> , 2013 , 27, 903.8	0.9	
12	Brain-mediated dysregulation of the bone marrow activity in angiotensin II-induced hypertension. <i>Hypertension</i> , 2012 , 60, 1316-23	8.5	51
11	Dysfunctional bone marrow-derived endothelial progenitor cells in chronic Ang II infusion rat model of hypertension. <i>FASEB Journal</i> , 2012 , 26, 878.7	0.9	
10	In vivo MEMRI reveals persistent activation of the brain autonomic areas by an acute systemic angiotensin II injection. <i>FASEB Journal</i> , 2012 , 26, lb801	0.9	
9	NTS (pro)renin receptor (PRR)-mediated antihypertensive effect involves NF-KappaB-cytokine signaling in the spontaneously hypertensive rats (SHR). <i>FASEB Journal</i> , 2012 , 26, 684.26	0.9	
8	Autonomic-immune-vascular interaction: an emerging concept for neurogenic hypertension. <i>Hypertension</i> , 2011 , 57, 1026-33	8.5	144
7	Role of GABAergic neurones in the nucleus tractus solitarii in modulation of cardiovascular activity. <i>Experimental Physiology</i> , 2010 , 95, 909-18	2.4	28
6	Shift to an involvement of phosphatidylinositol 3-kinase in angiotensin II actions on nucleus tractus solitarii neurons of the spontaneously hypertensive rat. <i>Circulation Research</i> , 2009 , 105, 1248-55	15.7	26
5	Chronic blockade of phosphatidylinositol 3-kinase in the nucleus tractus solitarii is prohypertensive in the spontaneously hypertensive rat. <i>Hypertension</i> , 2009 , 53, 97-103	8.5	17
4	Role of phosphoinositide-3-kinase (PI3K) in the nucleus of the solitary tract (NTS) in the modulation of baroreceptor reflex function in the hypertensive rat. <i>FASEB Journal</i> , 2008 , 22, 737.34	0.9	
3	Chronic inhibition of phosphoinositide-3-kinase (PI3K) in the nucleus of the solitary tract (NTS) of hypertensive rats increases blood pressure. <i>FASEB Journal</i> , 2007 , 21, A899	0.9	1
2	Structure-based discovery of a novel angiotensin-converting enzyme 2 inhibitor. <i>Hypertension</i> , 2004 , 44, 903-6	8.5	142

- 1 Cloning and characterization of a secreted form of angiotensin-converting enzyme 2. *Regulatory Peptides*, **2004**, 122, 61-7