

Tingting Yang

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9021178/publications.pdf>

Version: 2024-02-01

14
papers

478
citations

1040056

9
h-index

996975

15
g-index

16
all docs

16
docs citations

16
times ranked

479
citing authors

#	ARTICLE	IF	CITATIONS
1	Quercetin improves nonalcoholic fatty liver by ameliorating inflammation, oxidative stress, and lipid metabolism in <i>db/db</i> mice. <i>Phytotherapy Research</i> , 2019, 33, 3140-3152.	5.8	127
2	Quantitative profiling of 19 bile acids in rat plasma, liver, bile and different intestinal section contents to investigate bile acid homeostasis and the application of temporal variation of endogenous bile acids. <i>Journal of Steroid Biochemistry and Molecular Biology</i> , 2017, 172, 69-78.	2.5	107
3	Bile acid homeostasis paradigm and its connotation with cholestatic liver diseases. <i>Drug Discovery Today</i> , 2019, 24, 112-128.	6.4	47
4	Targeting mammalian serine/threonine-protein kinase 4 through Yes-associated protein/TEA domain transcription factor-mediated epithelial-mesenchymal transition ameliorates diabetic nephropathy orchestrated renal fibrosis. <i>Metabolism: Clinical and Experimental</i> , 2020, 108, 154258.	3.4	35
5	Early indications of ANIT-induced cholestatic liver injury: Alteration of hepatocyte polarization and bile acid homeostasis. <i>Food and Chemical Toxicology</i> , 2017, 110, 1-12.	3.6	34
6	Amelioration of non-alcoholic fatty liver disease by sodium butyrate is linked to the modulation of intestinal tight junctions in <i>db/db</i> mice. <i>Food and Function</i> , 2020, 11, 10675-10689.	4.6	33
7	YY1: A novel therapeutic target for diabetic nephropathy orchestrated renal fibrosis. <i>Metabolism: Clinical and Experimental</i> , 2019, 96, 33-45.	3.4	29
8	Quercetin Attenuates Podocyte Apoptosis of Diabetic Nephropathy Through Targeting EGFR Signaling. <i>Frontiers in Pharmacology</i> , 2021, 12, 792777.	3.5	13
9	Insulin exacerbated high glucose-induced epithelial-mesenchymal transition in prostatic epithelial cells BPH-1 and prostate cancer cells PC-3 via MEK/ERK signaling pathway. <i>Experimental Cell Research</i> , 2020, 394, 112145.	2.6	12
10	Sphingosine 1-phosphate receptor-1 specific agonist SEW2871 ameliorates ANIT-induced dysregulation of bile acid homeostasis in mice plasma and liver. <i>Toxicology Letters</i> , 2020, 331, 242-253.	0.8	11
11	SEW2871 attenuates ANIT-induced hepatotoxicity by protecting liver barrier function via sphingosine 1-phosphate receptor-1-mediated AMPK signaling pathway. <i>Cell Biology and Toxicology</i> , 2021, 37, 595-609.	5.3	11
12	YY1 inactivated transcription co-regulator PGC-1 β to promote mitochondrial dysfunction of early diabetic nephropathy-associated tubulointerstitial fibrosis. <i>Cell Biology and Toxicology</i> , 2023, 39, 391-413.	5.3	8
13	Comparison of bile acids profiles in the enterohepatic circulation system of mice and rats. <i>Journal of Steroid Biochemistry and Molecular Biology</i> , 2022, 220, 106100.	2.5	5
14	Simultaneous quantification of oestrogens and androgens in the serum of patients with benign prostatic hyperplasia by liquid chromatography-Tandem mass spectrometry. <i>Andrologia</i> , 2020, 52, e13611.	2.1	4