

# Sittisak Honsawek

## List of Publications by Year in Descending Order

**Source:** <https://exaly.com/author-pdf/9025752/sittisak-honsawek-publications-by-year.pdf>

**Version:** 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

120  
papers

1,840  
citations

24  
h-index

38  
g-index

130  
ext. papers

2,241  
ext. citations

3.9  
avg, IF

5.16  
L-index

#	Paper	IF	Citations
120	Safety and Immunogenicity of the Third Booster Dose with Inactivated, Viral Vector, and mRNA COVID-19 Vaccines in Fully Immunized Healthy Adults with Inactivated Vaccine.. <i>Vaccines</i> , <b>2022</b> , 10,	5.3	12
119	Neutralizing Activities against the Omicron Variant after a Heterologous Booster in Healthy Adults Receiving Two Doses of CoronaVac Vaccination.. <i>Journal of Infectious Diseases</i> , <b>2022</b> ,	7	7
118	Systemic cytokine profiles in biliary atresia.. <i>PLoS ONE</i> , <b>2022</b> , 17, e0267363	3.7	
117	Plasma and Joint Fluid Glypican-3 Are Inversely Correlated with the Severity of Knee Osteoarthritis. <i>Cartilage</i> , <b>2021</b> , 12, 505-511	3	3
116	Cellular senescence in liver fibrosis: Implications for age-related chronic liver diseases. <i>Expert Opinion on Therapeutic Targets</i> , <b>2021</b> , 25, 799-813	6.4	0
115	The Effects of Andrographolide on the Enhancement of Chondrogenesis and Osteogenesis in Human Suprapatellar Fat Pad Derived Mesenchymal Stem Cells. <i>Molecules</i> , <b>2021</b> , 26,	4.8	4
114	A novel gelatin/chitooligosaccharide/demineralized bone matrix composite scaffold and periosteum-derived mesenchymal stem cells for bone tissue engineering. <i>Biomaterials Research</i> , <b>2021</b> , 25, 19	16.8	4
113	Internalization of Neutrophil-Derived Microvesicles Modulates TNF $\alpha$ -stimulated Proinflammatory Cytokine Production in Human Fibroblast-Like Synoviocytes. <i>International Journal of Molecular Sciences</i> , <b>2021</b> , 22,	6.3	3
112	Isolation of Microvesicles from Human Circulating Neutrophils. <i>Bio-protocol</i> , <b>2021</b> , 11, e3119	0.9	0
111	Carboxymethyl Cellulose Entrapped in a Poly(vinyl) Alcohol Network: Plant-Based Scaffolds for Cartilage Tissue Engineering. <i>Molecules</i> , <b>2021</b> , 26,	4.8	8
110	Diagnostic Value of Interleukin-34 as a Novel Biomarker for Severity of Knee Osteoarthritis. <i>Cartilage</i> , <b>2021</b> , 1947603521990866	3	2
109	Cartilage oligomeric matrix protein as a marker of progressive liver fibrosis in biliary atresia. <i>Scientific Reports</i> , <b>2021</b> , 11, 16695	4.9	1
108	Global DNA hypomethylation of Alu and LINE-1 transposable elements as an epigenetic biomarker of anti-tuberculosis drug-induced liver injury. <i>Emerging Microbes and Infections</i> , <b>2021</b> , 10, 1862-1872	18.9	1
107	Interleukin-34 overexpression mediated through tumor necrosis factor-alpha reflects severity of synovitis in knee osteoarthritis. <i>Scientific Reports</i> , <b>2020</b> , 10, 7987	4.9	3
106	Decreased Serum Adiponectin Reflects Low Vitamin D, High Interleukin 6, and Poor Physical Performance in Knee Osteoarthritis. <i>Archivum Immunologiae Et Therapiae Experimentalis</i> , <b>2020</b> , 68, 16	4	1
105	Positive impact of lockdown on COVID-19 outbreak in Thailand. <i>Travel Medicine and Infectious Disease</i> , <b>2020</b> , 36, 101802	8.4	21
104	Hepatic expression of HGF/C-met and native liver survival in biliary atresia. <i>Pediatric Surgery International</i> , <b>2020</b> , 36, 597-602	2.1	0

103	Neutrophil-to-lymphocyte ratio predicts the severity of motor impairment in cerebral palsy children living at home and the rehabilitation center: A comparative study. <i>Biomedical Reports</i> , <b>2020</b> , 13, 63	1.8	
102	Neutrophil-to-lymphocyte ratio predicts the severity of motor impairment in cerebral palsy children living at home and the rehabilitation center: A comparative study. <i>Biomedical Reports</i> , <b>2020</b> , 13, 1-1	1.8	1
101	Telomere shortening is associated with poor physical performance in knee osteoarthritis. <i>Biomedical Reports</i> , <b>2020</b> , 13, 27	1.8	3
100	No Difference in Pain After Spine Surgery with Local Wound Filtration of Morphine and Ketorolac: A Randomized Controlled Trial. <i>Clinical Orthopaedics and Related Research</i> , <b>2020</b> , 478, 2823-2829	2.2	3
99	Anatomical Differences Between Foot and Hand Web Space Reconstruction. <i>Journal of Foot and Ankle Surgery</i> , <b>2020</b> , 59, 31-37	1.6	0
98	Hepatic glypican-3 and alpha-smooth muscle actin overexpressions reflect severity of liver fibrosis and predict outcome after successful portoenterostomy in biliary atresia. <i>Surgery</i> , <b>2020</b> , 167, 560-568	3.6	7
97	Relationships between blood leukocyte mitochondrial DNA copy number and inflammatory cytokines in knee osteoarthritis. <i>Journal of Zhejiang University: Science B</i> , <b>2020</b> , 21, 42-52	4.5	3
96	Decreased circulating clusterin reflects severe liver complications after hepatoportoenterostomy of biliary atresia. <i>Scientific Reports</i> , <b>2020</b> , 10, 19736	4.9	3
95	Clusterin Is Associated with Systemic and Synovial Inflammation in Knee Osteoarthritis. <i>Cartilage</i> , <b>2020</b> , 1947603520958149	3	5
94	Circulating and Synovial Fluid Heat Shock Protein 70 Are Correlated with Severity in Knee Osteoarthritis. <i>Cartilage</i> , <b>2020</b> , 11, 323-328	3	3
93	Blood leukocyte LINE-1 hypomethylation and oxidative stress in knee osteoarthritis. <i>Heliyon</i> , <b>2019</b> , 5, e01774	3.6	3
92	Interleukin-34 as a promising clinical biomarker and therapeutic target for inflammatory arthritis. <i>Cytokine and Growth Factor Reviews</i> , <b>2019</b> , 47, 43-53	17.9	10
91	Relative telomere length and oxidative stress in musculoskeletal tumors. <i>Molecular Biology Reports</i> , <b>2019</b> , 46, 4009-4016	2.8	
90	Leukocyte mitochondrial DNA copy number as a potential biomarker indicating poor outcome in biliary atresia and its association with oxidative DNA damage and telomere length. <i>Mitochondrion</i> , <b>2019</b> , 47, 1-9	4.9	11
89	Increased serum glypican-3 is associated with liver stiffness and hepatic dysfunction in children with biliary atresia. <i>Clinical and Experimental Hepatology</i> , <b>2019</b> , 5, 48-54	2.2	2
88	Serum adipokine levels, bodyweight and functional status in children with cerebral palsy. <i>Obesity Medicine</i> , <b>2019</b> , 16, 100154	2.6	1
87	Preparation of an injectable modified chitosan-based hydrogel approaching for bone tissue engineering. <i>International Journal of Biological Macromolecules</i> , <b>2019</b> , 123, 167-173	7.9	40
86	Do-It-Yourself Microsuture from Human Hair for Basic Microsurgical Training. <i>Journal of Reconstructive Microsurgery</i> , <b>2019</b> , 35, 315-321	2.5	2

85	Association between leukocyte telomere length and angiogenic cytokines in knee osteoarthritis. <i>International Journal of Rheumatic Diseases</i> , <b>2018</b> , 21, 118-125	2.3	18
84	Relative telomere length and oxidative DNA damage in hypertrophic ligamentum flavum of lumbar spinal stenosis. <i>PeerJ</i> , <b>2018</b> , 6, e5381	3.1	9
83	Alu hypermethylation and high oxidative stress in patients with musculoskeletal tumors. <i>PeerJ</i> , <b>2018</b> , 6, e5492	3.1	4
82	Hepatic autotaxin overexpression in infants with biliary atresia. <i>PeerJ</i> , <b>2018</b> , 6, e5224	3.1	2
81	Adiponectin as a novel biomarker for liver fibrosis. <i>World Journal of Hepatology</i> , <b>2018</b> , 10, 708-718	3.4	13
80	Elevated serum leptin levels are associated with low vitamin D, sarcopenic obesity, poor muscle strength, and physical performance in knee osteoarthritis. <i>Biomarkers</i> , <b>2017</b> , 22, 723-730	2.6	16
79	Noradrenaline, Serotonin, GABA, and Glycine in Cerebrospinal Fluid during Labor Pain: A Cross-Sectional Prospective Study. <i>Pain Research and Management</i> , <b>2017</b> , 2017, 2752658	2.6	2
78	Association between Promoter Hypomethylation and Overexpression of Autotaxin with Outcome Parameters in Biliary Atresia. <i>PLoS ONE</i> , <b>2017</b> , 12, e0169306	3.7	4
77	Mesenchymal stem cells for cartilage regeneration in osteoarthritis. <i>World Journal of Orthopedics</i> , <b>2017</b> , 8, 674-680	2.2	24
76	Elevated serum heat shock protein 70 and liver stiffness reflect hepatic dysfunction and severity in postoperative biliary atresia. <i>Pediatric Surgery International</i> , <b>2017</b> , 33, 893-899	2.1	3
75	Effect of vitamin E on oxidative stress level in blood, synovial fluid, and synovial tissue in severe knee osteoarthritis: a randomized controlled study. <i>BMC Musculoskeletal Disorders</i> , <b>2017</b> , 18, 281	2.8	16
74	Alteration of inflammatory cytokine production in primary total knee arthroplasty using antibiotic-loaded bone cement. <i>Journal of Orthopaedics and Traumatology</i> , <b>2017</b> , 18, 51-57	5	3
73	Vitamin D Supplementation Improves Quality of Life and Physical Performance in Osteoarthritis Patients. <i>Nutrients</i> , <b>2017</b> , 9,	6.7	52
72	Low bone mineral density and the severity of cholestasis in biliary atresia. <i>World Journal of Hepatology</i> , <b>2017</b> , 9, 746-751	3.4	3
71	Association of adiponectin gene polymorphisms with knee osteoarthritis. <i>World Journal of Orthopedics</i> , <b>2017</b> , 8, 719-725	2.2	7
70	Global methylation, oxidative stress, and relative telomere length in biliary atresia patients. <i>Scientific Reports</i> , <b>2016</b> , 6, 26969	4.9	31
69	Increased expression of vascular endothelial growth factor is associated with hypertrophic ligamentum flavum in lumbar spinal canal stenosis. <i>Journal of Investigative Medicine</i> , <b>2016</b> , 64, 882-7	2.9	7
68	Elevation of serum urokinase plasminogen activator receptor and liver stiffness in postoperative biliary atresia. <i>World Journal of Hepatology</i> , <b>2016</b> , 8, 1471-1477	3.4	3

67	Vitamin D and spine surgery. <i>World Journal of Orthopedics</i> , <b>2016</b> , 7, 726-730	2.2	12
66	Glomus tumors of the fingers: Expression of vascular endothelial growth factor. <i>World Journal of Orthopedics</i> , <b>2016</b> , 7, 843-846	2.2	2
65	Isolation and Characterization of Human Mesenchymal Stem Cells From Facet Joints and Interspinous Ligaments. <i>Spine</i> , <b>2016</b> , 41, E1-7	3.3	12
64	Serum Leptin as a Nutritional Biomarker in Children with Cerebral Palsy. <i>Tohoku Journal of Experimental Medicine</i> , <b>2016</b> , 239, 139-46	2.4	4
63	Role of pronator release in revision carpal tunnel surgery. <i>Sicot-j</i> , <b>2016</b> , 2, 9	2.2	7
62	Association between ADAM12 polymorphism and knee osteoarthritis in Thai population. <i>Knee</i> , <b>2016</b> , 23, 357-61	2.6	18
61	Plasma and synovial fluid inflammatory cytokine profiles in primary knee osteoarthritis. <i>Biomarkers</i> , <b>2016</b> , 21, 639-44	2.6	35
60	Association de la périostine plasmatique et synoviale à l'arthrose radiographique du genou : Étude transversale. <i>Revue Du Rhumatisme (Edition Francaise)</i> , <b>2016</b> , 83, 428-432	0.1	
59	A Water-Based Chitosan-Maleimide Precursor for Bioconjugation: An Example of a Rapid Pathway for an In Situ Injectable Adhesive Gel. <i>Macromolecular Rapid Communications</i> , <b>2016</b> , 37, 1618-1622	4.8	16
58	Serum autotaxin levels correlate with hepatic dysfunction and severity in postoperative biliary atresia. <i>Biomarkers</i> , <b>2015</b> , 20, 89-94	2.6	9
57	Increased serum sclerostin in postoperative biliary atresia. <i>Clinica Chimica Acta</i> , <b>2015</b> , 442, 136-40	6.2	2
56	Elevated serum periostin is associated with liver stiffness and clinical outcome in biliary atresia. <i>Biomarkers</i> , <b>2015</b> , 20, 157-61	2.6	10
55	Association of plasma and synovial fluid periostin with radiographic knee osteoarthritis: Cross-sectional study. <i>Joint Bone Spine</i> , <b>2015</b> , 82, 352-5	2.9	17
54	Cytokines as biochemical markers for knee osteoarthritis. <i>World Journal of Orthopedics</i> , <b>2015</b> , 6, 95-105	2.2	12.2
53	Characteristics and Osteoconductivity of Bone Composite Scaffolds Made of Thai Silk Fibroin, Gelatin and Inorganic Compounds: A Comparative Study of Tricalcium Phosphate and Hydroxyapatite. <i>Macromolecular Symposia</i> , <b>2015</b> , 354, 258-264	0.8	1
52	Telomere Length in Peripheral Blood Leukocytes Is Associated with Severity of Biliary Atresia. <i>PLoS ONE</i> , <b>2015</b> , 10, e0134689	3.7	13
51	Role of Vitamin D in Osteoarthritis: Molecular, Cellular, and Clinical Perspectives. <i>International Journal of Endocrinology</i> , <b>2015</b> , 2015, 383918	2.7	37
50	Plasma and synovial fluid autotaxin correlate with severity in knee osteoarthritis. <i>Clinica Chimica Acta</i> , <b>2015</b> , 444, 72-7	6.2	16

49	Angiogenic cytokine expression profiles in plasma and synovial fluid of primary knee osteoarthritis. <i>International Orthopaedics</i> , <b>2014</b> , 38, 1885-92	3.8	38
48	Adipokines: Biomarkers for osteoarthritis?. <i>World Journal of Orthopedics</i> , <b>2014</b> , 5, 319-27	2.2	67
47	Association of adiponectin +276G/T polymorphism with knee osteoarthritis. <i>Biomedical Reports</i> , <b>2014</b> , 2, 229-232	1.8	6
46	Current perspectives in mesenchymal stem cell therapies for osteoarthritis. <i>Stem Cells International</i> , <b>2014</b> , 2014, 194318	5	55
45	Plasma and synovial fluid sclerostin are inversely associated with radiographic severity of knee osteoarthritis. <i>Clinical Biochemistry</i> , <b>2014</b> , 47, 547-51	3.5	21
44	Relationship of plasma and synovial fluid vascular endothelial growth factor with radiographic severity in primary knee osteoarthritis. <i>International Orthopaedics</i> , <b>2014</b> , 38, 1099-104	3.8	41
43	Hypertrophy of the ligamentum flavum in lumbar spinal canal stenosis is associated with increased bFGF expression. <i>International Orthopaedics</i> , <b>2013</b> , 37, 1387-92	3.8	17
42	Association of MMP-3 (-1612 5A/6A) polymorphism with knee osteoarthritis in Thai population. <i>Rheumatology International</i> , <b>2013</b> , 33, 435-9	3.6	18
41	The effect of laminar air flow and door openings on operating room contamination. <i>Journal of Arthroplasty</i> , <b>2013</b> , 28, 1482-5	4.4	77
40	Soluble receptor for advanced glycation end products and liver stiffness in postoperative biliary atresia. <i>Clinical Biochemistry</i> , <b>2013</b> , 46, 214-8	3.5	11
39	Development of collagen/demineralized bone powder scaffolds and periosteum-derived cells for bone tissue engineering application. <i>International Journal of Molecular Sciences</i> , <b>2013</b> , 14, 2056-71	6.3	33
38	Oxidative stress, vitamin e, and antioxidant capacity in knee osteoarthritis. <i>Journal of Clinical and Diagnostic Research JCDR</i> , <b>2013</b> , 7, 1855-9	0	25
37	Correlation of connective tissue growth factor with liver stiffness measured by transient elastography in biliary atresia. <i>Hepatology Research</i> , <b>2013</b> , 43, 795-800	5.1	8
36	+276 G/T single nucleotide polymorphism of the adiponectin gene is associated with the susceptibility to biliary atresia. <i>World Journal of Pediatrics</i> , <b>2012</b> , 8, 328-34	4.6	14
35	Correlation between plasma and synovial fluid basic fibroblast growth factor with radiographic severity in primary knee osteoarthritis. <i>International Orthopaedics</i> , <b>2012</b> , 36, 981-5	3.8	22
34	Plasma and synovial fluid connective tissue growth factor levels are correlated with disease severity in patients with knee osteoarthritis. <i>Biomarkers</i> , <b>2012</b> , 17, 303-8	2.6	21
33	Association of X-prolyl aminopeptidase 1 rs17095355 polymorphism with biliary atresia in Thai children. <i>Hepatology Research</i> , <b>2011</b> , 41, 1249-52	5.1	23
32	Association of plasma and synovial fluid interferon- $\gamma$ -inducible protein-10 with radiographic severity in knee osteoarthritis. <i>Clinical Biochemistry</i> , <b>2011</b> , 44, 1218-22	3.5	19

31	Association of the IL-6 -174G/C gene polymorphism with knee osteoarthritis in a Thai population. <i>Genetics and Molecular Research</i> , <b>2011</b> , 10, 1674-80	1.2	26
30	Association of circulating osteopontin levels with clinical outcomes in postoperative biliary atresia. <i>Pediatric Surgery International</i> , <b>2011</b> , 27, 283-8	2.1	6
29	High serum matrix metalloproteinase-3 and liver stiffness in postoperative biliary atresia. <i>Pediatric Surgery International</i> , <b>2011</b> , 27, 681-7	2.1	6
28	Relationship of serum IL-6, C-reactive protein, erythrocyte sedimentation rate, and knee skin temperature after total knee arthroplasty: a prospective study. <i>International Orthopaedics</i> , <b>2011</b> , 35, 31-5	3.8	53
27	Serum retinol binding protein 4 and clinical outcome in postoperative biliary atresia. <i>Hepatology International</i> , <b>2011</b> , 5, 906-12	8.8	5
26	Serum adiponectin and transient elastography as non-invasive markers for postoperative biliary atresia. <i>BMC Gastroenterology</i> , <b>2011</b> , 11, 16	3	16
25	Elevation of serum galectin-3 and liver stiffness measured by transient elastography in biliary atresia. <i>European Journal of Pediatric Surgery</i> , <b>2011</b> , 21, 250-4	1.9	14
24	Gene expression analysis of demineralized bone matrix-induced osteogenesis in human periosteal cells using cDNA array technology. <i>Genetics and Molecular Research</i> , <b>2011</b> , 10, 2093-103	1.2	4
23	RhBMP-2 and -7 combined with absorbable collagen sponge carrier enhance ectopic bone formation: An in vivo bioassay. <i>Asian Biomedicine</i> , <b>2011</b> , 5, 85-92	0.4	
22	Dickkopf-1 (Dkk-1) in plasma and synovial fluid is inversely correlated with radiographic severity of knee osteoarthritis patients. <i>BMC Musculoskeletal Disorders</i> , <b>2010</b> , 11, 257	2.8	58
21	Correlation of plasma and synovial fluid adiponectin with knee osteoarthritis severity. <i>Archives of Medical Research</i> , <b>2010</b> , 41, 593-8	6.6	80
20	Soluble receptor for advanced glycation end products (sRAGE) in plasma and synovial fluid is inversely associated with disease severity of knee osteoarthritis. <i>Clinical Biochemistry</i> , <b>2010</b> , 43, 1133-7	3.5	24
19	Osteoinductive potential of small intestinal submucosa/ demineralized bone matrix as composite scaffolds for bone tissue engineering. <i>Asian Biomedicine</i> , <b>2010</b> , 4, 913-922	0.4	3
18	Increased osteopontin and liver stiffness measurement by transient elastography in biliary atresia. <i>World Journal of Gastroenterology</i> , <b>2010</b> , 16, 5467-73	5.6	16
17	Stem cell factor in gingival crevicular fluid in periodontal health and disease. <i>Asian Biomedicine</i> , <b>2010</b> , 4, 281-287	0.4	1
16	Elevated circulating and synovial fluid endoglin are associated with primary knee osteoarthritis severity. <i>Archives of Medical Research</i> , <b>2009</b> , 40, 590-4	6.6	11
15	Relationships between OPG, RANKL, bone metabolism, and bone mineral density in biliary atresia. <i>Pediatric Surgery International</i> , <b>2009</b> , 25, 261-7	2.1	5
14	Inhibin A levels and severity of preeclampsia. <i>Archives of Gynecology and Obstetrics</i> , <b>2009</b> , 280, 183-6	2.5	5



13	Relationship of plasma and synovial fluid BMP-7 with disease severity in knee osteoarthritis patients: a pilot study. <i>International Orthopaedics</i> , <b>2009</b> , 33, 1171-5	3.8	41
12	Correlation of plasma and synovial fluid osteopontin with disease severity in knee osteoarthritis. <i>Clinical Biochemistry</i> , <b>2009</b> , 42, 808-12	3.5	52
11	Circulating leptin levels and bone mineral density in children with biliary atresia. <i>Acta Paediatrica, International Journal of Paediatrics</i> , <b>2008</b> , 97, 206-11	3.1	7
10	Rosiglitazone effect on radioiodine uptake in thyroid carcinoma patients with high thyroglobulin but negative total body scan: a correlation with the expression of peroxisome proliferator-activated receptor-gamma. <i>Thyroid</i> , <b>2008</b> , 18, 697-704	6.2	55
9	First trimester serum inhibin A in normal pregnant women. <i>Archives of Gynecology and Obstetrics</i> , <b>2008</b> , 277, 307-10	2.5	3
8	High levels of serum basic fibroblast growth factor in children with biliary atresia. <i>Hepato-Gastroenterology</i> , <b>2008</b> , 55, 1184-8		6
7	Elevation of serum stem-cell factor in postoperative biliary atresia. <i>Pediatrics International</i> , <b>2007</b> , 49, 888-93	1.2	7
6	Increased levels of serum hyaluronan in patients with dengue infection. <i>Journal of Infection</i> , <b>2007</b> , 54, 225-9	18.9	11
5	Association of serum levels of tissue inhibitors of metalloproteinase-1 with clinical outcome in children with biliary atresia. <i>Asian Pacific Journal of Allergy and Immunology</i> , <b>2006</b> , 24, 161-6	5.4	4
4	Effects of demineralized bone matrix on proliferation and osteogenic differentiation of mesenchymal stem cells from human umbilical cord. <i>Journal of the Medical Association of Thailand = Chotmai het Thangphaet</i> , <b>2006</b> , 89 Suppl 3, S189-95		8
3	Serum interleukin-8 in children with biliary atresia: relationship with disease stage and biochemical parameters. <i>Pediatric Surgery International</i> , <b>2005</b> , 21, 73-7	2.1	36
2	Extractable bone morphogenetic protein and correlation with induced new bone formation in an in vivo assay in the athymic mouse model. <i>Cell and Tissue Banking</i> , <b>2005</b> , 6, 13-23	2.2	32
1	Content of bone morphogenetic protein-4 in human demineralized bone: relationship to donor age and ability to induce new bone formation. <i>Journal of the Medical Association of Thailand = Chotmai het Thangphaet</i> , <b>2005</b> , 88 Suppl 4, S260-5		5