Dagmar M Ouweneel

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/9028026/publications.pdf

Version: 2024-02-01



#	Article	IF	CITATIONS
1	Percutaneous Mechanical Circulatory Support Versus Intra-Aortic Balloon PumpÂin Cardiogenic Shock After AcuteÂMyocardial Infarction. Journal of the American College of Cardiology, 2017, 69, 278-287.	1.2	612
2	Extracorporeal life support during cardiac arrest and cardiogenic shock: a systematic review and meta-analysis. Intensive Care Medicine, 2016, 42, 1922-1934.	3.9	405
3	Percutaneous short-term active mechanical support devices in cardiogenic shock: a systematic review and collaborative meta-analysis of randomized trials. European Heart Journal, 2017, 38, 3523-3531.	1.0	280
4	Epinephrine and short-term survival in cardiogenic shock: an individual data meta-analysis of 2583 patients. Intensive Care Medicine, 2018, 44, 847-856.	3.9	106
5	Percutaneous Mechanical Circulatory Support Versus Intra-Aortic Balloon Pump for Treating Cardiogenic Shock. Journal of the American College of Cardiology, 2017, 69, 358-360.	1.2	98
6	The ICM research agenda on extracorporeal life support. Intensive Care Medicine, 2017, 43, 1306-1318.	3.9	94
7	Mechanical circulatory support in cardiogenic shock from acute myocardial infarction: Impella CP/5.0 versus ECMO. European Heart Journal: Acute Cardiovascular Care, 2020, 9, 164-172.	0.4	72
8	Percutaneous cardiac support devices for cardiogenic shock: current indications and recommendations. Heart, 2012, 98, 1246-1254.	1.2	62
9	Limitations and Opportunities of Transcutaneous Bilirubin Measurements. Pediatrics, 2012, 129, 689-694.	1.0	60
10	Real-life use of left ventricular circulatory support with Impella in cardiogenic shock after acute myocardial infarction: 12 years AMC experience. European Heart Journal: Acute Cardiovascular Care, 2019, 8, 338-349.	0.4	55
11	Improved recovery of regional left ventricular function after PCI of chronic total occlusion in STEMI patients: a cardiovascular magnetic resonance study of the randomized controlled EXPLORE trial. Journal of Cardiovascular Magnetic Resonance, 2017, 19, 53.	1.6	41
12	Prognostic Value of Access Site and Nonaccess Site Bleeding After Percutaneous Coronary Intervention. JACC: Cardiovascular Interventions, 2014, 7, 622-630.	1.1	34
13	Evaluating the learning curve in the prospective Randomized Clinical Trial of hemodynamic support with Impella 2.5 versus Intra-Aortic Balloon Pump in patients undergoing high-risk percutaneous coronary intervention: a prespecified subanalysis of the PROTECT II study. American Heart Journal, 2014 167 472-479 e5	1.2	34
14	Long-term 5-year outcome of the randomized IMPRESS in severe shock trial: percutaneous mechanical circulatory support vs. intra-aortic balloon pump in cardiogenic shock after acute myocardial infarction. European Heart Journal: Acute Cardiovascular Care, 2021, 10, 1009-1015.	0.4	30
15	Lactate is a Prognostic Factor in Patients Admitted With Suspected ST-Elevation Myocardial Infarction. Shock, 2019, 51, 321-327.	1.0	28
16	Vasopressors and Inotropes in Acute Myocardial Infarction Related Cardiogenic Shock: A Systematic Review and Meta-Analysis. Journal of Clinical Medicine, 2020, 9, 2051.	1.0	21
17	Arterial Pressure Variation as a Biomarker of Preload Dependency in Spontaneously Breathing Subjects – A Proof of Principle. PLoS ONE, 2015, 10, e0137364.	1.1	17
18	Pre-PCI versus immediate post-PCI Impella initiation in acute myocardial infarction complicated by cardiogenic shock. PLoS ONE, 2020, 15, e0235762.	1.1	14

DAGMAR M OUWENEEL

#	Article	IF	CITATIONS
19	Recovery and prognostic value of myocardial strain in ST-segment elevation myocardial infarction patients with a concurrent chronic total occlusion. European Radiology, 2020, 30, 600-608.	2.3	13
20	Impact of collateralisation to a concomitant chronic total occlusion in patients with ST-elevation myocardial infarction: a subanalysis of the EXPLORE randomised controlled trial. Open Heart, 2018, 5, e000810.	0.9	11
21	The impact of the location of a chronic total occlusion in a non-infarct-related artery on long-term mortality in ST-elevation myocardial infarction patients. EuroIntervention, 2016, 12, 423-430.	1.4	8
22	The effect of revascularization of a chronic total coronary occlusion on electrocardiographic variables. A sub-study of the EXPLORE trial. Journal of Electrocardiology, 2018, 51, 906-912.	0.4	6
23	Value of the SYNTAX Score in ST-Elevation Myocardial Infarction Patients With a Concomitant Chronic Total Coronary Occlusion(from the EXPLORE Trial). American Journal of Cardiology, 2019, 123, 1035-1043.	0.7	6
24	Assessment of Cardiac Device Position on Supine Chest Radiograph in the ICU. Critical Care Medicine, 2016, 44, e957-e963.	0.4	3
25	Predictors and outcomes of procedural failure of percutaneous coronary intervention of a chronic total occlusion—A subanalysis of the EXPLORE trial. Catheterization and Cardiovascular Interventions, 2021, 97, 1176-1183.	0.7	2
26	The Role of Percutaneous Haemodynamic Support in High-risk Percutaneous Coronary Intervention and Cardiogenic Shock. Interventional Cardiology Review, 2015, 10, 39.	0.7	2
27	Collateral Quality Decay Several Days After Primary Percutaneous Coronary Intervention. JACC: Cardiovascular Interventions, 2018, 11, 511-512.	1.1	0