

# Hou-Qun Ying

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9028390/publications.pdf>

Version: 2024-02-01

53  
papers

1,897  
citations

331538

21  
h-index

265120

42  
g-index

56  
all docs

56  
docs citations

56  
times ranked

3065  
citing authors

#	ARTICLE	IF	CITATIONS
1	Upregulated lncRNA-UCA1 contributes to progression of hepatocellular carcinoma through inhibition of miR-216b and activation of FGFR1/ERK signaling pathway. <i>Oncotarget</i> , 2015, 6, 7899-7917.	0.8	329
2	The prognostic value of preoperative NLR, d-NLR, PLR and LMR for predicting clinical outcome in surgical colorectal cancer patients. <i>Medical Oncology</i> , 2014, 31, 305.	1.2	239
3	Prognostic value of pre-operative inflammatory response biomarkers in gastric cancer patients and the construction of a predictive model. <i>Journal of Translational Medicine</i> , 2015, 13, 66.	1.8	172
4	Prognostic value of neutrophil-lymphocyte ratio in breast cancer. <i>FEBS Open Bio</i> , 2015, 5, 502-507.	1.0	104
5	Albumin-fibrinogen ratio as a promising biomarker to predict clinical outcome of non-small cell lung cancer individuals. <i>Cancer Medicine</i> , 2018, 7, 1221-1231.	1.3	74
6	Up-Regulation of 91H Promotes Tumor Metastasis and Predicts Poor Prognosis for Patients with Colorectal Cancer. <i>PLoS ONE</i> , 2014, 9, e103022.	1.1	72
7	Two new inflammatory markers associated with disease activity score-28 in patients with rheumatoid arthritis: Albumin to fibrinogen ratio and C-reactive protein to albumin ratio. <i>International Immunopharmacology</i> , 2018, 62, 293-298.	1.7	61
8	Genome-Wide Profiling of Prognostic Alternative Splicing Signature in Colorectal Cancer. <i>Frontiers in Oncology</i> , 2018, 8, 537.	1.3	54
9	Neutrophil-lymphocyte ratio and platelet-lymphocyte ratio are 2 new inflammatory markers associated with pulmonary involvement and disease activity in patients with dermatomyositis. <i>Clinica Chimica Acta</i> , 2017, 465, 11-16.	0.5	50
10	MiR-608, pre-miR-124-1 and pre-miR26a-1 polymorphisms modify susceptibility and recurrence-free survival in surgically resected CRC individuals. <i>Oncotarget</i> , 2016, 7, 75865-75873.	0.8	44
11	Combination of preoperative NLR, PLR and CEA could increase the diagnostic efficacy for I-III stage CRC. <i>Journal of Clinical Laboratory Analysis</i> , 2017, 31, e22075.	0.9	39
12	Circulating vitamin D binding protein, total, free and bioavailable 25-hydroxyvitamin D and risk of colorectal cancer. <i>Scientific Reports</i> , 2015, 5, 7956.	1.6	38
13	Prognostic value of a novel FPR biomarker in patients with surgical stage II and III gastric cancer. <i>Oncotarget</i> , 2017, 8, 75195-75205.	0.8	35
14	Prognostic Value of Long Non-Coding RNA HOTAIR in Various Cancers. <i>PLoS ONE</i> , 2014, 9, e110059.	1.1	32
15	Circulating fibrinogen to prealbumin ratio is a promising biomarker for diagnosis of colorectal cancer. <i>Journal of Clinical Laboratory Analysis</i> , 2019, 33, e22635.	0.9	32
16	The predictive and prognostic role of a novel ADS score in esophageal squamous cell carcinoma patients undergoing esophagectomy. <i>Cancer Cell International</i> , 2018, 18, 153.	1.8	31
17	Preoperative circulating FPR and CCF score are promising biomarkers for predicting clinical outcome of stage II&ndash;III colorectal cancer patients. <i>Cancer Management and Research</i> , 2018, Volume 10, 2151-2161.	0.9	30
18	Circulating miR-148/152 family as potential biomarkers in hepatocellular carcinoma. <i>Tumor Biology</i> , 2016, 37, 4945-4953.	0.8	27

#	ARTICLE	IF	CITATIONS
19	Platelet-to-lymphocyte ratio could be a promising prognostic biomarker for survival of colorectal cancer: a systematic review and meta-analysis. <i>FEBS Open Bio</i> , 2016, 6, 742-750.	1.0	26
20	Preoperative fibrinogen to prealbumin ratio as a novel predictor for clinical outcome of hepatocellular carcinoma. <i>Future Oncology</i> , 2019, 15, 13-22.	1.1	25
21	Relationship of SNP of H2BFWT gene to male infertility in a Chinese population with idiopathic spermatogenesis impairment. <i>Biomarkers</i> , 2012, 17, 402-406.	0.9	23
22	The effect of BIM deletion polymorphism on intrinsic resistance and clinical outcome of cancer patient with kinase inhibitor therapy. <i>Scientific Reports</i> , 2015, 5, 11348.	1.6	23
23	Significance of combined preoperative serum Alb and dNLR for diagnosis of pancreatic cancer. <i>Future Oncology</i> , 2018, 14, 229-239.	1.1	21
24	The Role of Cancer-Elicited Inflammatory Biomarkers in Predicting Early Recurrence Within Stage II-III Colorectal Cancer Patients After Curable Resection. <i>Journal of Inflammation Research</i> , 2021, Volume 14, 115-129.	1.6	20
25	Elevated FPR confers to radiochemoresistance and predicts clinical efficacy and outcome of metastatic colorectal cancer patients. <i>Aging</i> , 2019, 11, 1716-1732.	1.4	20
26	CCL5-403, CCR5-59029, and Delta32 polymorphisms and cancer risk: a meta-analysis based on 20,625 subjects. <i>Tumor Biology</i> , 2014, 35, 5895-5904.	0.8	18
27	Chemotherapy plus bevacizumab as an optimal first-line therapeutic treatment for patients with right-sided metastatic colon cancer: a meta-analysis of first-line clinical trials. <i>ESMO Open</i> , 2020, 5, e000605.	2.0	18
28	Exome sequencing identified a novel SMAD2 mutation in a Chinese family with early onset aortic aneurysms. <i>Clinica Chimica Acta</i> , 2017, 468, 211-214.	0.5	17
29	The involvement of Kras gene 3'-UTR polymorphisms in risk of cancer and influence on patient response to anti-EGFR therapy in metastatic colorectal cancer: a meta-analysis. <i>OncoTargets and Therapy</i> , 2014, 7, 1487.	1.0	16
30	Genetic variants of eNOS gene may modify the susceptibility to idiopathic male infertility. <i>Biomarkers</i> , 2013, 18, 412-417.	0.9	15
31	Null genotype of GSTM1 and GSTT1 may contribute to susceptibility to male infertility with impaired spermatogenesis in Chinese population. <i>Biomarkers</i> , 2013, 18, 151-154.	0.9	13
32	A systematic review on the association between the <i>Helicobacter pylori vacA</i> genotype and gastric disease. <i>FEBS Open Bio</i> , 2016, 6, 409-417.	1.0	13
33	Primary Tumor Sidedness Predicts Bevacizumab Benefit in Metastatic Colorectal Cancer Patients. <i>Frontiers in Oncology</i> , 2019, 9, 723.	1.3	13
34	The value of circulating fibrinogen-to-pre-albumin ratio in predicting survival and benefit from chemotherapy in colorectal cancer. <i>Therapeutic Advances in Medical Oncology</i> , 2021, 13, 175883592110228.	1.4	13
35	Gene therapy for human colorectal cancer cell lines with recombinant adenovirus 5 based on loss of the insulin-like growth factor 2 imprinting. <i>International Journal of Oncology</i> , 2015, 46, 1759-1767.	1.4	12
36	Evaluation the susceptibility of five polymorphisms in microRNA-binding sites to female breast cancer risk in Chinese population. <i>Gene</i> , 2015, 573, 160-165.	1.0	11

#	ARTICLE	IF	CITATIONS
37	The diagnostic role of circulating inflammation-based biomarker in gallbladder carcinoma. <i>Biomarkers in Medicine</i> , 2018, 12, 1095-1103.	0.6	11
38	Gene therapy for colorectal cancer by adenovirus-mediated siRNA targeting CD147 based on loss of the IGF2 imprinting system. <i>International Journal of Oncology</i> , 2015, 47, 1881-1889.	1.4	10
39	Albumin to fibrinogen ratio and fibrinogen to pre-albumin ratio are economical, simple and promising prognostic factors for solid malignancy. <i>Journal of Thoracic Disease</i> , 2019, 11, S2036-S2038.	0.6	10
40	FCGR2A, FCGR3A polymorphisms and therapeutic efficacy of anti-EGFR monoclonal antibody in metastatic colorectal cancer. <i>Oncotarget</i> , 2015, 6, 28071-28083.	0.8	10
41	Meta-analysis of prognostic value of inflammation parameter in breast cancer. <i>Journal of Cancer Research and Therapeutics</i> , 2018, 14, S85-S89.	0.3	10
42	Inhibition of CD147 expression by RNA interference reduces proliferation, invasion and increases chemosensitivity in cancer stem cell-like HT-29 cells. <i>International Journal of Oncology</i> , 2015, 47, 1476-1484.	1.4	9
43	Association of GSTM1 and GSTT1 Genes with the Susceptibility to Male Infertility: Result from a Meta-Analysis. <i>Genetic Testing and Molecular Biomarkers</i> , 2013, 17, 535-542.	0.3	8
44	Trends of incidence and survival in patients with gastrointestinal mucinous adenocarcinoma. <i>Oncology Letters</i> , 2018, 16, 5791-5798.	0.8	7
45	Cancer-elicited inflammation attenuates response and outcome in tyrosine kinase inhibitor naive patients with advanced NSCLC. <i>Pharmacological Research</i> , 2021, 170, 105734.	3.1	7
46	Analysis of Diagnostic Value of YKL-40 in Ovarian Cancer. <i>International Journal of Gynecological Cancer</i> , 2016, , 1.	1.2	6
47	Methyltransferase DNMT3B in leukemia. <i>Leukemia and Lymphoma</i> , 2020, 61, 263-273.	0.6	6
48	MAD2 Combined with Mitotic Spindle Apparatus (MSA) and Anticentromere Antibody (ACA) for Diagnosis of Small Cell Lung Cancer (SCLC). <i>Medical Science Monitor</i> , 2018, 24, 7541-7547.	0.5	6
49	Role of Chronic Inflammatory Ratios in Predicting Recurrence of Resected Patients with Stage III Mucinous Colorectal Adenocarcinoma. <i>Cancer Management and Research</i> , 2021, Volume 13, 3455-3464.	0.9	5
50	Quantification of fibrinogen-to-pre-albumin ratio provides an integrating parameter for differential diagnosis and risk stratification of early-stage colorectal cancer. <i>Cancer Cell International</i> , 2022, 22, 137.	1.8	5
51	High-Grade Inflammation Attenuates Chemosensitivity and Confers to Poor Survival of Surgical Stage III CRC Patients. <i>Frontiers in Oncology</i> , 2021, 11, 580455.	1.3	3
52	Different effects of the three polymorphisms on 15q25.1 on lung cancer risk: Evidence from published literatures. <i>Journal of Cancer Research and Therapeutics</i> , 2016, 12, 12.	0.3	3
53	A Novel Prognostic Score Based on ZG16 for Predicting CRC Survival. <i>Pharmacogenomics and Personalized Medicine</i> , 2020, Volume 13, 735-747.	0.4	1