Ana Neves

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/9041236/publications.pdf

Version: 2024-02-01

687363 713466 45 543 13 21 citations h-index g-index papers 45 45 45 629 citing authors all docs docs citations times ranked

#	Article	IF	CITATIONS
1	Diet comparison of four ray species (Raja clavata, Raja brachyura, Raja montaguiandLeucoraja naevus) caught along the Portuguese continental shelf. Aquatic Living Resources, 2006, 19, 105-114.	1.2	41
2	Molecular barcoding of north-east Atlantic deep-water sharks: species identification and application to fisheries management and conservation. Marine and Freshwater Research, 2008, 59, 214.	1.3	40
3	Reproductive strategy of leafscale gulper shark <i>Centrophorus squamosus</i> and the Portuguese dogfish <i>Centroscymnus coelolepis</i> on the Portuguese continental slope. Journal of Fish Biology, 2008, 73, 206-225.	1.6	37
4	Macroparasites as biological tags for stock identification of the bluemouth, Helicolenus dactylopterus (Delaroche, 1809) in Portuguese waters. Fisheries Research, 2010, 106, 321-328.	1.7	28
5	Otolith shape analysis as a tool for stock discrimination of forkbeard (Phycis phycis) in the Northeast Atlantic. Hydrobiologia, 2014, 728, 103-110.	2.0	27
6	Age and growth of bluemouth, Helicolenus dactylopterus, from the Portuguese continental slope. ICES Journal of Marine Science, 2009, 66, 524-531.	2.5	26
7	Reproductive strategies in black scabbardfish (<i>Aphanopus carbo</i> Lowe, 1839) from the NE Atlantic. Scientia Marina, 2009, 73, 19-31.	0.6	25
8	The use of caudal thorns for ageing Raja undulata from the Portuguese continental shelf, with comments on its reproductive cycle. Marine and Freshwater Research, 2007, 58, 983.	1.3	22
9	Discriminating bluemouth, Helicolenus dactylopterus (Pisces: Sebastidae), stocks in Portuguese waters by means of otolith shape analysis. Journal of the Marine Biological Association of the United Kingdom, 2011, 91, 1237-1242.	0.8	21
10	Age and growth of black scabbardfish (<i>Aphanopus carbo</i> Lowe, 1839) in the southern NE Atlantic. Scientia Marina, 2009, 73, 33-46.	0.6	19
11	Evidence for temporal changes in ray and skate populations in the Portuguese coast (1998–2003) – its implications in the ecosystem. Aquatic Living Resources, 2007, 20, 85-93.	1.2	15
12	Distribution and abundance patterns of decapod crustaceans in the Sado estuary, Portugal. Crustaceana, 2007, 80, 97-112.	0.3	14
13	Distribution patterns and reproduction of the cuttlefish, <i>Sepia officinalis</i> in the Sado estuary (Portugal). Journal of the Marine Biological Association of the United Kingdom, 2009, 89, 579-584.	0.8	14
14	Using body geometric morphometrics to identify bluemouth, Helicolenus dactylopterus (Delaroche,) Tj ETQq0 0	0 rgBT /0	verlack 10 Tf S
15	The gelatinous matrix of the teleost <i>Helicolenus dactylopterus dactylopterus</i> (Delaroche, 1809) in the context of its reproductive strategy. Marine Biology Research, 2011, 7, 478-487.	0.7	14
16	Life history parameters as possible discriminators of bluemouth Helicolenus dactylopterus (Delaroche, 1809) populations in Portuguese waters. Fisheries Research, 2012, 125-126, 69-76.	1.7	14
17	Feeding habits of the bluemouth, Helicolenus dactylopterus dactylopterus (Delaroche, 1809) (Pisces:) Tj ETQq1	1 0.78431 1.3	14 rgBT /Overla 14
18	Genetic and Morphological Variation of the Forkbeard, Phycis phycis (Pisces, Phycidae): Evidence of Panmixia and Recent Population Expansion along Its Distribution Area. PLoS ONE, 2016, 11, e0167045.	2.5	14

#	Article	IF	CITATIONS
19	Feeding habits of the cuttlefish SepiaÂofficinalis during its life cycle in the Sado estuary (Portugal). Hydrobiologia, 2009, 636, 479-488.	2.0	11
20	Reproductive strategy of the female deep-water shark birdbeak dogfish, <i>Deania calcea < /i> lecithotrophy or matrotrophy?. Journal of the Marine Biological Association of the United Kingdom, 2012, 92, 387-394.</i>	0.8	10
21	Adding Value to Bycatch Fish Species Captured in the Portuguese Coast—Development of New Food Products. Foods, 2021, 10, 68.	4.3	9
22	Age, growth and reproduction of the protandrous hermaphrodite fish, Sarpa salpa, from the Portuguese continental coast. Journal of the Marine Biological Association of the United Kingdom, 2018, 98, 269-281.	0.8	8
23	Age and growth of forkbeard, Phycis phycis, in Portuguese continental waters. Journal of the Marine Biological Association of the United Kingdom, 2014, 94, 623-630.	0.8	7
24	Reproductive strategy of forkbeard, Phycis phycis, from the Portuguese coast. Helgoland Marine Research, 2016, 70, .	1.3	7
25	Otolith shape and isotopic ratio analyses as a tool to study Spondyliosoma cantharus population structure. Marine Environmental Research, 2019, 143, 93-100.	2.5	7
26	Stock structure of black scabbardfish (<i>Aphanopus carbo</i> Lowe, 1839) in the southern northeast Atlantic. Scientia Marina, 2009, 73, 89-101.	0.6	7
27	Age, growth and mortality of the comber <i>Serranus cabrilla</i> (Linnaeus, 1758) in the Eastern Atlantic. Marine Biology Research, 2016, 12, 656-662.	0.7	6
28	Modelling the growth of a protogynous sparid species, Spondyliosoma cantharus (Teleostei:) Tj ETQq0 0 0 rgBT	Overlock 2.0	10 ₆ Tf 50 382
29	Phenotypic changes in the body of black seabream, Spondyliosoma cantharus (Teleostei: Sparidae), along the eastern Atlantic. Estuarine, Coastal and Shelf Science, 2018, 214, 31-37.	2.1	6
30	Age, growth and mortality of <i>Pontinus kuhlii</i> (Bowdich, 1825) (Scorpaeniformes:) Tj ETQq0 0 0	rgBT/Ove	rlock 10 Tf 50
31	Risks and benefits' consumption of birdbeak dogfishDeania calcea. British Food Journal, 2012, 114, 826-839.	2.9	5
32	Reproductive patterns of blacktail comber (Serranus atricauda, Serranidae) from south-west Portugal seamounts. Helgoland Marine Research, 2014, 68, 133-142.	1.3	5
33	Diversity of sexual strategies of fish species caught by bottom trawl off the western Iberian Peninsula. Marine Biology Research, 2015, 11, 361-374.	0.7	5
34	Insight on reproductive strategy in Portuguese waters of a commercial protogynous species, the black seabream Spondyliosoma cantharus (Sparidae). Fisheries Research, 2018, 206, 85-95.	1.7	5
35	Seasonal Sensory Evaluation of Low Commercial Value or Unexploited Fish Species from the Portuguese Coast. Foods, 2020, 9, 1880.	4.3	5
36	Highly regional population structure of Spondyliosoma cantharus depicted by nuclear and mitochondrial DNA data. Scientific Reports, 2020, 10, 4063.	3.3	4

#	Article	IF	CITATIONS
37	Age and growth of small red scorpionfish, Scorpaena notata (Actinopterygii: Scorpaeniformes:) Tj ETQq1 1 0.7843 Piscatoria, 2015, 45, 13-20.	314 rgBT / 0.7	Overlock 1 4
38	Modelling Fish Growth with Imperfect Data: The Case of Trachurus picturatus. Fishes, 2022, 7, 52.	1.7	4
39	Seasonal study of the nutritional composition of unexploited and low commercial value fish species from the Portuguese coast. Food Science and Nutrition, 2022, 10, 3368-3379.	3.4	4
40	Whole or sectioned otoliths? Choosing the best method for aging bluemouth, Helicolenus dactyloterus (Delaroche, 1809). Fisheries Research, 2013, 147, 235-239.	1.7	3
41	Estimating fecundity in the zygoparous species Helicolenus dactylopterus (Actinopterygii,) Tj ETQq1 1 0.784314 r	·gBT Ove	rlgck 10 Tf
42	Zygoparity and sex steroid hormone profiles in bluemouth <i>Helicolenus dactylopterus </i> . Journal of Fish Biology, 2017, 90, 2157-2169.	1.6	3
43	Diet and feeding strategy of the forkbeard Phycis phycis (Pisces: Phycidae) from the Portuguese continental coast. Journal of the Marine Biological Association of the United Kingdom, 2018, 98, 1757-1765.	0.8	2
44	Fecundity and sex steroid profile in boarfish Capros aper. Marine and Freshwater Research, 2021, 72, 140.	1.3	1
45	Enhancing knowledge on lowâ€value fishing species: the distinct reproductive strategy of two gurnard species. Journal of Fish Biology, 2021, 99, 1403-1414.	1.6	1