

Pingyang Wang

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9052351/publications.pdf>

Version: 2024-02-01

10
papers

39
citations

1937685

4
h-index

1872680

6
g-index

10
all docs

10
docs citations

10
times ranked

78
citing authors

#	ARTICLE	IF	CITATIONS
1	The silkworm (<i>Bombyx mori</i>) neuropeptide orcokinin is involved in the regulation of pigmentation. <i>Insect Biochemistry and Molecular Biology</i> , 2019, 114, 103229.	2.7	11
2	Mutation of a Cuticle Protein Gene, <i>BmCPG10</i> , Is Responsible for Silkworm Non-Moulting in the 2nd Instar Mutant. <i>PLoS ONE</i> , 2016, 11, e0153549.	2.5	10
3	Transcriptome analysis of the epidermis of the purple quail-like (q-lp) mutant of silkworm, <i>Bombyx mori</i> . <i>PLoS ONE</i> , 2017, 12, e0175994.	2.5	5
4	Differentially expressed genes in the head of the 2nd instar pre-molting larvae of the nm2 mutant of the silkworm, <i>Bombyx mori</i> . <i>PLoS ONE</i> , 2017, 12, e0180160.	2.5	4
5	A Review of Pedal Peptide/Orcokinin-type Neuropeptides. <i>Current Protein and Peptide Science</i> , 2021, 22, 41-49.	1.4	3
6	Downregulation of aldose reductase is responsible for developmental abnormalities of the silkworm purple quail-like mutant (q-l p). <i>Gene</i> , 2018, 665, 96-104.	2.2	2
7	The expression analysis of cysteine proteinase-like protein in wild-type and nm2 mutant silkworm (<i>Lepidoptera: Bombyx mori</i>). <i>Gene</i> , 2016, 586, 170-175.	2.2	1
8	Semaphorin-1a-like gene plays an important role in the embryonic development of silkworm, <i>Bombyx mori</i> . <i>PLoS ONE</i> , 2020, 15, e0240193.	2.5	1
9	Transcriptome analysis of the eggs of the silkworm pale red egg (rep-1) mutant at 36 hours after oviposition. <i>PLoS ONE</i> , 2020, 15, e0237242.	2.5	1
10	phytanoyl-CoA dioxygenase domain-containing protein 1 plays an important role in egg shell formation of silkworm (<i>Bombyx mori</i>). <i>PLoS ONE</i> , 2021, 16, e0261918.	2.5	1