Denis E Naumov

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/9058604/publications.pdf Version: 2024-02-01



DENIS E NAUMOV

#	Article	IF	CITATIONS
1	Effect of TRPM8 and TRPA1 Polymorphisms on COPD Predisposition and Lung Function in COPD Patients. Journal of Personalized Medicine, 2021, 11, 108.	1.1	5
2	Thermosensory Transient Receptor Potential Ion Channels and Asthma. Biomedicines, 2021, 9, 816.	1.4	7
3	Functional activity of bronchial granulocytes in the cytokine profile formation in asthma patients during airway reaction to cold stimulus. Immunologiya, 2020, 41, 432-440.	0.1	0
4	POLYMORPHISM OF TRPM8 GENE AS AN INDEPENDENT FACTOR OF BRONCHIAL OBSUCTION IN ASTHMA. Bulletin Physiology and Pathology of Respiration, 2019, 1, 31-36.	0.0	1
5	PECULIARITIES OF TRPM8 RECEPTOR EXPRESSION IN THE RESPIRATORY TRACT OF ASTHMA PATIENTS. Bulletin Physiology and Pathology of Respiration, 2018, 1, 19-24.	0.0	3
6	TRPM8 is overexpressed in the respiratory tract of steroid-naive asthma patients. Asian Pacific Journal of Tropical Medicine, 2018, 11, 16.	0.4	2
7	ROLE OF TRPM8 GENE POLYMORPHISMS IN THE FORMATION OF ASTHMA PHENOTYPE WITH COLD AIRWAY HYPERRESPONSIVENESS. Bulletin Physiology and Pathology of Respiration, 2017, 1, 16-23.	0.0	2
8	TRPM8 GENE POLYMORPHISM AND SMOKING AS THE FACTORS OF SEVERE BRONCHIAL OBSTRUCTION IN PATIENTS WITH ASTHMA. Bulletin Physiology and Pathology of Respiration, 2017, 1, 24-30.	0.0	4
9	Influence of TRPV4 gene polymorphisms on the development of osmotic airway hyperresponsiveness in patients with bronchial asthma. Doklady Biochemistry and Biophysics, 2016, 469, 260-263.	0.3	11
10	Transient receptor potential melastatin 8 gene polymorphism is associated with coldâ€induced airway hyperresponsiveness in bronchial asthma. Respirology, 2015, 20, 1192-1197.	1.3	24
11	Role of β2 Adrenoreceptor Gene Polymorphism in the Formation of Cold Hyperreactivity of the Airways in Asthmatics. Bulletin of Experimental Biology and Medicine, 2012, 154, 73-76.	0.3	2
12	A metabolite of prostaglandin D2, 11β-prostaglandin F2α (11β-PGF2α), in exhaled breath condensate and serum of asthmatics with airway hyperresponsiveness to distilled water. F1000Research, 0, 5, 307.	0.8	0