

Guruprasad P Aithal

List of Publications by Citations

Source: <https://exaly.com/author-pdf/9059840/guruprasad-p-aithal-publications-by-citations.pdf>

Version: 2024-04-23

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

205
papers

11,812
citations

50
h-index

107
g-index

232
ext. papers

14,815
ext. citations

6.6
avg. IF

6.36
L-index

| # | Paper | IF | Citations |
|-----|---|------|-----------|
| 205 | Association of polymorphisms in the cytochrome P450 CYP2C9 with warfarin dose requirement and risk of bleeding complications. <i>Lancet, The</i> , 1999 , 353, 717-9 | 40 | 1066 |
| 204 | Liraglutide safety and efficacy in patients with non-alcoholic steatohepatitis (LEAN): a multicentre, double-blind, randomised, placebo-controlled phase 2 study. <i>Lancet, The</i> , 2016 , 387, 679-690 | 40 | 935 |
| 203 | HLA-B*5701 genotype is a major determinant of drug-induced liver injury due to flucloxacillin. <i>Nature Genetics</i> , 2009 , 41, 816-9 | 36.3 | 818 |
| 202 | Randomized, placebo-controlled trial of pioglitazone in nondiabetic subjects with nonalcoholic steatohepatitis. <i>Gastroenterology</i> , 2008 , 135, 1176-84 | 13.3 | 516 |
| 201 | Noninvasive markers of fibrosis in nonalcoholic fatty liver disease: Validating the European Liver Fibrosis Panel and exploring simple markers. <i>Hepatology</i> , 2008 , 47, 455-60 | 11.2 | 494 |
| 200 | A randomized, placebo-controlled trial of cenicriviroc for treatment of nonalcoholic steatohepatitis with fibrosis. <i>Hepatology</i> , 2018 , 67, 1754-1767 | 11.2 | 376 |
| 199 | TM6SF2 rs58542926 influences hepatic fibrosis progression in patients with non-alcoholic fatty liver disease. <i>Nature Communications</i> , 2014 , 5, 4309 | 17.4 | 362 |
| 198 | Susceptibility to amoxicillin-clavulanate-induced liver injury is influenced by multiple HLA class I and II alleles. <i>Gastroenterology</i> , 2011 , 141, 338-47 | 13.3 | 359 |
| 197 | EASL Clinical Practice Guidelines: Drug-induced liver injury. <i>Journal of Hepatology</i> , 2019 , 70, 1222-1261 | 13.4 | 327 |
| 196 | Genetic susceptibility to diclofenac-induced hepatotoxicity: contribution of UGT2B7, CYP2C8, and ABCC2 genotypes. <i>Gastroenterology</i> , 2007 , 132, 272-81 | 13.3 | 277 |
| 195 | Drug-induced liver injury: recent advances in diagnosis and risk assessment. <i>Gut</i> , 2017 , 66, 1154-1164 | 19.2 | 250 |
| 194 | Hepatotoxicity Related to Anti-tuberculosis Drugs: Mechanisms and Management. <i>Journal of Clinical and Experimental Hepatology</i> , 2013 , 3, 37-49 | 4.1 | 225 |
| 193 | Hepatic adducts, circulating antibodies, and cytokine polymorphisms in patients with diclofenac hepatotoxicity. <i>Hepatology</i> , 2004 , 39, 1430-40 | 11.2 | 188 |
| 192 | Incidence and Etiology of Drug-Induced Liver Injury in Mainland China. <i>Gastroenterology</i> , 2019 , 156, 2230-2241 | 13.4 | 181 |
| 191 | Technical aspects of endoscopic ultrasound (EUS)-guided sampling in gastroenterology: European Society of Gastrointestinal Endoscopy (ESGE) Technical Guideline - March 2017. <i>Endoscopy</i> , 2017 , 49, 989-1006 | 3.4 | 182 |
| 190 | Indications, results, and clinical impact of endoscopic ultrasound (EUS)-guided sampling in gastroenterology: European Society of Gastrointestinal Endoscopy (ESGE) Clinical Guideline - Updated January 2017. <i>Endoscopy</i> , 2017 , 49, 695-714 | 3.4 | 166 |
| 189 | Drug-induced liver injury. <i>Nature Reviews Disease Primers</i> , 2019 , 5, 58 | 51.1 | 148 |

| | | | |
|-----|---|------|-----|
| 188 | Association of Liver Injury From Specific Drugs, or Groups of Drugs, With Polymorphisms in HLA and Other Genes in a Genome-Wide Association Study. <i>Gastroenterology</i> , 2017 , 152, 1078-1089 | 13.3 | 137 |
| 187 | No difference between high-fructose and high-glucose diets on liver triacylglycerol or biochemistry in healthy overweight men. <i>Gastroenterology</i> , 2013 , 145, 1016-1025.e2 | 13.3 | 131 |
| 186 | Incidence and prevalence of cirrhosis in the United Kingdom, 1992-2001: a general population-based study. <i>Journal of Hepatology</i> , 2008 , 49, 732-8 | 13.4 | 128 |
| 185 | Nonsteroidal anti-inflammatory drug-induced hepatotoxicity. <i>Clinics in Liver Disease</i> , 2007 , 11, 563-75, vi-vii | 4.6 | 117 |
| 184 | Genecriviroc Treatment for Adults With Nonalcoholic Steatohepatitis and Fibrosis: Final Analysis of the Phase 2b CENTAUR Study. <i>Hepatology</i> , 2020 , 72, 892-905 | 11.2 | 116 |
| 183 | Human leucocyte antigen class II genotype in susceptibility and resistance to co-amoxiclav-induced liver injury. <i>Journal of Hepatology</i> , 2010 , 53, 1049-53 | 13.4 | 114 |
| 182 | Genome-wide association study of non-alcoholic fatty liver and steatohepatitis in a histologically characterised cohort. <i>Journal of Hepatology</i> , 2020 , 73, 505-515 | 13.4 | 113 |
| 181 | Endoscopic treatment of chronic pancreatitis: European Society of Gastrointestinal Endoscopy (ESGE) Guideline - Updated August 2018. <i>Endoscopy</i> , 2019 , 51, 179-193 | 3.4 | 109 |
| 180 | Clinical diagnostic scale: a useful tool in the evaluation of suspected hepatotoxic adverse drug reactions. <i>Journal of Hepatology</i> , 2000 , 33, 949-52 | 13.4 | 108 |
| 179 | Hepatotoxicity related to antirheumatic drugs. <i>Nature Reviews Rheumatology</i> , 2011 , 7, 139-50 | 8.1 | 106 |
| 178 | Accuracy of hepatic adverse drug reaction reporting in one English health region. <i>BMJ: British Medical Journal</i> , 1999 , 319, 1541 | | 102 |
| 177 | ADAPT: An Algorithm Incorporating PRO-C3 Accurately Identifies Patients With NAFLD and Advanced Fibrosis. <i>Hepatology</i> , 2019 , 69, 1075-1086 | 11.2 | 100 |
| 176 | Efficacy, safety, and predictive factors for a positive yield of EUS-guided Trucut biopsy: a large tertiary referral center experience. <i>American Journal of Gastroenterology</i> , 2009 , 104, 584-91 | 0.7 | 100 |
| 175 | Limited contribution of common genetic variants to risk for liver injury due to a variety of drugs. <i>Pharmacogenetics and Genomics</i> , 2012 , 22, 784-95 | 1.9 | 96 |
| 174 | Relationship of polymorphism in CYP2C9 to genetic susceptibility to diclofenac-induced hepatitis. <i>Pharmacogenetics and Genomics</i> , 2000 , 10, 511-8 | | 96 |
| 173 | Genetic basis of drug-induced liver injury: present and future. <i>Seminars in Liver Disease</i> , 2014 , 34, 123-33 | 7.3 | 95 |
| 172 | Helical CT versus EUS with fine needle aspiration for celiac nodal assessment in patients with esophageal cancer. <i>Gastrointestinal Endoscopy</i> , 2002 , 55, 648-54 | 5.2 | 94 |
| 171 | All-cause mortality in people with cirrhosis compared with the general population: a population-based cohort study. <i>Liver International</i> , 2012 , 32, 79-84 | 7.9 | 89 |

| | | | |
|-----|--|------|----|
| 170 | 1 and 5 year survival estimates for people with cirrhosis of the liver in England, 1998-2009: a large population study. <i>Journal of Hepatology</i> , 2014 , 60, 282-9 | 13.4 | 82 |
| 169 | Upstream and coding region CYP2C9 polymorphisms: correlation with warfarin dose and metabolism. <i>Pharmacogenetics and Genomics</i> , 2004 , 14, 813-22 | | 82 |
| 168 | Validation of terminal peptide of procollagen III for the detection and assessment of nonalcoholic steatohepatitis in patients with nonalcoholic fatty liver disease. <i>Hepatology</i> , 2013 , 57, 103-11 | 11.2 | 81 |
| 167 | Association of non-alcoholic steatohepatitis without significant fibrosis with hepatocellular carcinoma. <i>Journal of Hepatology</i> , 2004 , 41, 685-6 | 13.4 | 81 |
| 166 | Prevalence of clinically significant liver disease within the general population, as defined by non-invasive markers of liver fibrosis: a systematic review. <i>The Lancet Gastroenterology and Hepatology</i> , 2017 , 2, 288-297 | 18.8 | 80 |
| 165 | Minocycline hepatotoxicity: Clinical characterization and identification of HLA-B*35:02 as a risk factor. <i>Journal of Hepatology</i> , 2017 , 67, 137-144 | 13.4 | 78 |
| 164 | Fracture risk in people with primary biliary cirrhosis: a population-based cohort study. <i>Gastroenterology</i> , 2006 , 131, 1752-7 | 13.3 | 71 |
| 163 | A study of T1ρ relaxation time as a measure of liver fibrosis and the influence of confounding histological factors. <i>NMR in Biomedicine</i> , 2015 , 28, 706-14 | 4.4 | 69 |
| 162 | Direct targeting of risk factors significantly increases the detection of liver cirrhosis in primary care: a cross-sectional diagnostic study utilising transient elastography. <i>BMJ Open</i> , 2015 , 5, e007516 | 3 | 66 |
| 161 | Influence of ursodeoxycholic acid on the mortality and malignancy associated with primary biliary cirrhosis: a population-based cohort study. <i>Hepatology</i> , 2007 , 46, 1131-7 | 11.2 | 63 |
| 160 | Non-invasive assessment of portal hypertension using quantitative magnetic resonance imaging. <i>Journal of Hepatology</i> , 2016 , 65, 1131-1139 | 13.4 | 61 |
| 159 | A Missense Variant in PTPN22 is a Risk Factor for Drug-induced Liver Injury. <i>Gastroenterology</i> , 2019 , 156, 1707-1716.e2 | 13.3 | 59 |
| 158 | Pharmacogenetic testing in idiosyncratic drug-induced liver injury: current role in clinical practice. <i>Liver International</i> , 2015 , 35, 1801-8 | 7.9 | 53 |
| 157 | Transient elastography for screening of liver fibrosis: Cost-effectiveness analysis from six prospective cohorts in Europe and Asia. <i>Journal of Hepatology</i> , 2019 , 71, 1141-1151 | 13.4 | 51 |
| 156 | A role for the pregnane X receptor in flucloxacillin-induced liver injury. <i>Hepatology</i> , 2010 , 51, 1656-64 | 11.2 | 50 |
| 155 | HLA-DRB1*16: 01-DQB1*05: 02 is a novel genetic risk factor for flupirtine-induced liver injury. <i>Pharmacogenetics and Genomics</i> , 2016 , 26, 218-24 | 1.9 | 50 |
| 154 | Lower gut microbiome diversity and higher abundance of proinflammatory genus are associated with biopsy-proven nonalcoholic steatohepatitis. <i>Gut Microbes</i> , 2020 , 11, 569-580 | 8.8 | 49 |
| 153 | N-acetyltransferase 2 (NAT2) genotype as a risk factor for development of drug-induced liver injury relating to antituberculosis drug treatment in a mixed-ethnicity patient group. <i>European Journal of Clinical Pharmacology</i> , 2014 , 70, 1079-86 | 2.8 | 48 |

| | | | |
|-----|--|------|----|
| 152 | Genetic Variation in HSD17B13 Reduces the Risk of Developing Cirrhosis and Hepatocellular Carcinoma in Alcohol Misusers. <i>Hepatology</i> , 2020 , 72, 88-102 | 11.2 | 46 |
| 151 | Pharmacogenomics of drug-induced liver injury (DILI): Molecular biology to clinical applications. <i>Journal of Hepatology</i> , 2018 , 69, 948-957 | 13.4 | 45 |
| 150 | Risk of cardiovascular and cerebrovascular events in primary biliary cirrhosis: a population-based cohort study. <i>American Journal of Gastroenterology</i> , 2008 , 103, 2784-8 | 0.7 | 44 |
| 149 | Guidelines on the management of ascites in cirrhosis. <i>Gut</i> , 2021 , 70, 9-29 | 19.2 | 43 |
| 148 | The pathological response to neoadjuvant chemotherapy with FOLFOX-4 for colorectal liver metastases: a comparative study. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 2007 , 451, 943-8 | 5.1 | 41 |
| 147 | Accuracy of EUS for detection of intraductal papillary mucinous tumor of the pancreas. <i>Gastrointestinal Endoscopy</i> , 2002 , 56, 701-7 | 5.2 | 41 |
| 146 | Human leukocyte antigen genetic risk factors of drug-induced liver toxicology. <i>Expert Opinion on Drug Metabolism and Toxicology</i> , 2015 , 11, 395-409 | 5.5 | 40 |
| 145 | Mass spectrometric characterization of circulating covalent protein adducts derived from a drug acyl glucuronide metabolite: multiple albumin adductions in diclofenac patients. <i>Journal of Pharmacology and Experimental Therapeutics</i> , 2014 , 350, 387-402 | 4.7 | 39 |
| 144 | Role of polymorphisms in the interleukin-10 gene in determining disease susceptibility and phenotype in inflammatory bowel disease. <i>Digestive Diseases and Sciences</i> , 2001 , 46, 1520-5 | 4 | 39 |
| 143 | Role of Drugs Used for Chronic Disease Management on Susceptibility and Severity of COVID-19: A Large Case-Control Study. <i>Clinical Pharmacology and Therapeutics</i> , 2020 , 108, 1185-1194 | 6.1 | 39 |
| 142 | Natural history of histologically proven alcohol-related liver disease: A systematic review. <i>Journal of Hepatology</i> , 2019 , 71, 586-593 | 13.4 | 38 |
| 141 | Drug-Induced Liver Injury due to Flucloxacillin: Relevance of Multiple Human Leukocyte Antigen Alleles. <i>Clinical Pharmacology and Therapeutics</i> , 2019 , 106, 245-253 | 6.1 | 35 |
| 140 | Shared Genetic Risk Factors Across Carbamazepine-Induced Hypersensitivity Reactions. <i>Clinical Pharmacology and Therapeutics</i> , 2019 , 106, 1028-1036 | 6.1 | 34 |
| 139 | Causality assessment for suspected DILI during clinical phases of drug development. <i>Drug Safety</i> , 2014 , 37 Suppl 1, S47-56 | 5.1 | 34 |
| 138 | Polygenic architecture informs potential vulnerability to drug-induced liver injury. <i>Nature Medicine</i> , 2020 , 26, 1541-1548 | 50.5 | 34 |
| 137 | Economic evaluation of a community-based diagnostic pathway to stratify adults for non-alcoholic fatty liver disease: a Markov model informed by a feasibility study. <i>BMJ Open</i> , 2017 , 7, e015659 | 3 | 33 |
| 136 | Monitoring liver function during methotrexate therapy for psoriasis: are routine biopsies really necessary?. <i>American Journal of Clinical Dermatology</i> , 2005 , 6, 357-63 | 7.1 | 33 |
| 135 | Increased liver fat and glycogen stores after consumption of high versus low glycaemic index food: A randomized crossover study. <i>Diabetes, Obesity and Metabolism</i> , 2017 , 19, 70-77 | 6.7 | 31 |

| | | | |
|-----|---|------|----|
| 134 | Diagnostic accuracy of non-invasive tests for advanced fibrosis in patients with NAFLD: an individual patient data meta-analysis. <i>Gut</i> , 2021 , | 19.2 | 31 |
| 133 | Unacceptable failures: the final report of the Lancet Commission into liver disease in the UK. <i>Lancet, The</i> , 2020 , 395, 226-239 | 40 | 28 |
| 132 | The European NAFLD Registry: A real-world longitudinal cohort study of nonalcoholic fatty liver disease. <i>Contemporary Clinical Trials</i> , 2020 , 98, 106175 | 2.3 | 28 |
| 131 | Prevalence and natural history of histologically proven chronic liver disease in a longitudinal cohort of patients with type 1 diabetes. <i>Hepatology</i> , 2014 , 60, 158-68 | 11.2 | 27 |
| 130 | Obesity and type 2 diabetes are important risk factors underlying previously undiagnosed cirrhosis in general practice: a cross-sectional study using transient elastography. <i>Alimentary Pharmacology and Therapeutics</i> , 2018 , 47, 504-515 | 6.1 | 27 |
| 129 | Volixibat in adults with non-alcoholic steatohepatitis: 24-week interim analysis from a randomized, phase II study. <i>Journal of Hepatology</i> , 2020 , 73, 231-240 | 13.4 | 26 |
| 128 | The influence of adiposity and acute exercise on circulating hepatokines in normal-weight and overweight/obese men. <i>Applied Physiology, Nutrition and Metabolism</i> , 2018 , 43, 482-490 | 3 | 25 |
| 127 | Multi-organ assessment of compensated cirrhosis patients using quantitative magnetic resonance imaging. <i>Journal of Hepatology</i> , 2018 , 69, 1015-1024 | 13.4 | 24 |
| 126 | Efficacy of recombinant human interleukin-10 in prevention of post-endoscopic retrograde cholangiopancreatography pancreatitis in subjects with increased risk. <i>Pancreas</i> , 2009 , 38, 267-74 | 2.6 | 24 |
| 125 | Multicenter experience from the UK and Ireland of use of lumen-apposing metal stent for transluminal drainage of pancreatic fluid collections. <i>Endoscopy International Open</i> , 2018 , 6, E259-E265 | 3 | 23 |
| 124 | Effect of exercise intensity on circulating hepatokine concentrations in healthy men. <i>Applied Physiology, Nutrition and Metabolism</i> , 2019 , 44, 1065-1072 | 3 | 19 |
| 123 | CYP2C9 polymorphism and warfarin dose requirements. <i>British Journal of Clinical Pharmacology</i> , 2002 , 53, 408-9 | 3.8 | 19 |
| 122 | Obesity Is the Most Common Risk Factor for Chronic Liver Disease: Results From a Risk Stratification Pathway Using Transient Elastography. <i>American Journal of Gastroenterology</i> , 2019 , 114, 1744-1752 | 0.7 | 19 |
| 121 | Human Leukocyte Antigen B*14:01 and B*35:01 Are Associated With Trimethoprim-Sulfamethoxazole Induced Liver Injury. <i>Hepatology</i> , 2021 , 73, 268-281 | 11.2 | 19 |
| 120 | Diagnostic accuracy of elastography and magnetic resonance imaging in patients with NAFLD: A systematic review and meta-analysis. <i>Journal of Hepatology</i> , 2021 , 75, 770-785 | 13.4 | 19 |
| 119 | Dynamics of 5-carboxylcytosine during hepatic differentiation: Potential general role for active demethylation by DNA repair in lineage specification. <i>Epigenetics</i> , 2017 , 12, 277-286 | 5.7 | 18 |
| 118 | The utility of scoring systems in predicting early and late mortality in alcoholic hepatitis: whose score is it anyway?. <i>International Journal of Hepatology</i> , 2012 , 2012, 624675 | 2.7 | 18 |
| 117 | Dangerous liaisons: drug, host and the environment. <i>Journal of Hepatology</i> , 2007 , 46, 995-8 | 13.4 | 18 |

| | | | |
|-----|--|------|----|
| 116 | EUS-guided Trucut mural biopsies in the investigation of unexplained thickening of the esophagogastric wall. <i>Gastrointestinal Endoscopy</i> , 2005 , 62, 624-9 | 5.2 | 18 |
| 115 | Genome-wide Association Study and Meta-analysis on Alcohol-Associated Liver Cirrhosis Identifies Genetic Risk Factors. <i>Hepatology</i> , 2021 , 73, 1920-1931 | 11.2 | 18 |
| 114 | Safety and long term efficacy of porfimer sodium photodynamic therapy in locally advanced biliary tract carcinoma. <i>Photodiagnosis and Photodynamic Therapy</i> , 2012 , 9, 287-92 | 3.5 | 17 |
| 113 | Genetic regulation of warfarin metabolism and response. <i>Seminars in Vascular Medicine</i> , 2003 , 3, 231-8 | | 17 |
| 112 | Warfarin dose requirement and CYP2C9 polymorphisms. <i>Lancet, The</i> , 1999 , 353, 1972-1973 | 40 | 17 |
| 111 | Genome-Wide Association Studies in Drug-Induced Liver Injury: Step Change in Understanding the Pathogenesis. <i>Seminars in Liver Disease</i> , 2015 , 35, 421-31 | 7.3 | 16 |
| 110 | Two doses of the SARS-CoV-2 BNT162b2 vaccine enhance antibody responses to variants in individuals with prior SARS-CoV-2 infection. <i>Science Translational Medicine</i> , 2021 , 13, eabj0847 | 17.5 | 16 |
| 109 | Effects of short-term energy restriction on liver lipid content and inflammatory status in severely obese adults: Results of a randomized controlled trial using 2 dietary approaches. <i>Diabetes, Obesity and Metabolism</i> , 2017 , 19, 1179-1183 | 6.7 | 15 |
| 108 | Care standards for non-alcoholic fatty liver disease in the United Kingdom 2016: a cross-sectional survey. <i>Frontline Gastroenterology</i> , 2017 , 8, 252-259 | 2.6 | 15 |
| 107 | Drug Development for Nonalcoholic Fatty Liver Disease: Landscape and Challenges. <i>Journal of Clinical and Experimental Hepatology</i> , 2019 , 9, 515-521 | 4.1 | 15 |
| 106 | Drug-induced liver injury: Asia Pacific Association of Study of Liver consensus guidelines. <i>Hepatology International</i> , 2021 , 15, 258-282 | 8.8 | 15 |
| 105 | Economic modelling of early transjugular intrahepatic portosystemic shunt insertion for acute variceal haemorrhage. <i>European Journal of Gastroenterology and Hepatology</i> , 2013 , 25, 201-7 | 2.2 | 14 |
| 104 | Efficacy, Safety, and Predictive Factors for a Positive Yield of EUS-Guided Trucut Biopsy. <i>American Journal of Gastroenterology</i> , 2009 , 104, 584-591 | 0.7 | 14 |
| 103 | Validation of a Model for Identification of Patients With Compensated Cirrhosis at High Risk of Decompensation. <i>Clinical Gastroenterology and Hepatology</i> , 2019 , 17, 2330-2338.e1 | 6.9 | 13 |
| 102 | Acute Hyperenergetic, High-Fat Feeding Increases Circulating FGF21, LECT2, and Fetuin-A in Healthy Men. <i>Journal of Nutrition</i> , 2020 , 150, 1076-1085 | 4.1 | 12 |
| 101 | Diagnosis, presentation and initial severity of Autoimmune Hepatitis (AIH) in patients attending 28 hospitals in the UK. <i>Liver International</i> , 2018 , 38, 1686-1695 | 7.9 | 12 |
| 100 | Effects of sprint interval training on ectopic lipids and tissue-specific insulin sensitivity in men with non-alcoholic fatty liver disease. <i>European Journal of Applied Physiology</i> , 2018 , 118, 817-828 | 3.4 | 12 |
| 99 | Visual morphometry and three non-invasive markers in the evaluation of liver fibrosis in chronic liver disease. <i>Scandinavian Journal of Gastroenterology</i> , 2017 , 52, 107-115 | 2.4 | 12 |

| | | | |
|----|---|------|----|
| 98 | Glycaemic, gastrointestinal and appetite responses to breakfast porridges from ancient cereal grains: A MRI pilot study in healthy humans. <i>Food Research International</i> , 2019 , 118, 49-57 | 7 | 12 |
| 97 | Genetic Risk Factors in Drug-Induced Liver Injury Due to Isoniazid-Containing Antituberculosis Drug Regimens. <i>Clinical Pharmacology and Therapeutics</i> , 2021 , 109, 1125-1135 | 6.1 | 12 |
| 96 | Non-alcoholic fatty liver disease: Not time for an obituary just yet!. <i>Journal of Hepatology</i> , 2021 , 74, 972-974 | 3.4 | 12 |
| 95 | Safety and efficacy of hydrothermal duodenal mucosal resurfacing in patients with type 2 diabetes: the randomised, double-blind, sham-controlled, multicentre REVITA-2 feasibility trial. <i>Gut</i> , 2021 , | 19.2 | 11 |
| 94 | Clinical and microbiological features of infection in alcoholic hepatitis: an international cohort study. <i>Journal of Gastroenterology</i> , 2017 , 52, 1192-1200 | 6.9 | 10 |
| 93 | When is a herb a drug?. <i>European Journal of Gastroenterology and Hepatology</i> , 2005 , 17, 391-3 | 2.2 | 10 |
| 92 | Immune dysfunction in patients with obstructive jaundice before and after endoscopic retrograde cholangiopancreatography. <i>Clinical Science</i> , 2016 , 130, 1535-44 | 6.5 | 10 |
| 91 | Advanced preclinical models for evaluation of drug-induced liver injury - consensus statement by the European Drug-Induced Liver Injury Network [PRO-EURO-DILI-NET]. <i>Journal of Hepatology</i> , 2021 , 75, 935-959 | 13.4 | 10 |
| 90 | Glycaemic, gastrointestinal, hormonal and appetitive responses to pearl millet or oats porridge breakfasts: a randomised, crossover trial in healthy humans. <i>British Journal of Nutrition</i> , 2019 , 122, 1142-1154 | 3.6 | 9 |
| 89 | A Revised Electronic Version of RUCAM for the Diagnosis of Drug Induced Liver Injury.. <i>Hepatology</i> , 2022 , | 11.2 | 9 |
| 88 | Prevention and management of idiosyncratic drug-induced liver injury: Systematic review and meta-analysis of randomised clinical trials. <i>Pharmacological Research</i> , 2021 , 164, 105404 | 10.2 | 9 |
| 87 | HLA associations with infliximab-induced liver injury. <i>Pharmacogenomics Journal</i> , 2020 , 20, 681-686 | 3.5 | 8 |
| 86 | Evaluation of laboratory tests for cirrhosis and for alcohol use, in the context of alcoholic cirrhosis. <i>Alcohol</i> , 2018 , 66, 1-7 | 2.7 | 8 |
| 85 | The detection of oesophageal varices using a novel, disposable, probe-based transnasal endoscope: a prospective diagnostic pilot study. <i>Liver International</i> , 2016 , 36, 1639-1648 | 7.9 | 8 |
| 84 | Hepatotoxicity Related to Methotrexate 2013 , 593-604 | | 8 |
| 83 | Longitudinal assessment of symptoms and risk of SARS-CoV-2 infection in healthcare workers across 5 hospitals to understand ethnic differences in infection risk. <i>EClinicalMedicine</i> , 2021 , 34, 100835 | 11.3 | 8 |
| 82 | Current and Future Magnetic Resonance Technologies for Assessing Liver Disease in Clinical and Experimental Medicine. <i>Digestive Diseases</i> , 2017 , 35, 314-322 | 3.2 | 7 |
| 81 | Cohort profile: the Trivandrum non-alcoholic fatty liver disease (NAFLD) cohort. <i>BMJ Open</i> , 2019 , 9, e027244 | 2.44 | 7 |

| | | | |
|----|--|------|---|
| 80 | Genetic variants of hepatic transporters and susceptibility to drug induced liver injury. <i>Toxicology</i> , 2008 , 253, 10 | 4.4 | 7 |
| 79 | Using MRI to study the alterations in liver blood flow, perfusion, and oxygenation in response to physiological stress challenges: Meal, hyperoxia, and hypercapnia. <i>Journal of Magnetic Resonance Imaging</i> , 2019 , 49, 1577-1586 | 5.6 | 7 |
| 78 | The role of hepatic lipid composition in obesity-related metabolic disease. <i>Liver International</i> , 2021 , | 7.9 | 7 |
| 77 | Incidence of post-ERCP pancreatitis from direct pancreatic juice collection in hereditary pancreatitis and familial pancreatic cancer before and after the introduction of prophylactic pancreatic stents and rectal diclofenac. <i>Pancreas</i> , 2015 , 44, 260-5 | 2.6 | 6 |
| 76 | Zanubrutinib-induced liver injury: a case report and literature review. <i>BMC Gastroenterology</i> , 2021 , 21, 244 | 3 | 6 |
| 75 | Evaluating the Sensitivity and Specificity of Promising Circulating Biomarkers to Diagnose Liver Injury in Humans. <i>Toxicological Sciences</i> , 2021 , 181, 23-34 | 4.4 | 6 |
| 74 | Elevated bilirubin, alkaline phosphatase at onset, and drug metabolism are associated with prolonged recovery from DILI. <i>Journal of Hepatology</i> , 2021 , 75, 333-341 | 13.4 | 6 |
| 73 | Healthcare costs of transarterial chemoembolization in the treatment of hepatocellular carcinoma. <i>Journal of Hepatocellular Carcinoma</i> , 2017 , 4, 123-130 | 5.3 | 5 |
| 72 | The performance of transient elastography compared to clinical acumen and routine tests - what is the incremental diagnostic value?. <i>Liver International</i> , 2013 , 33, 172-9 | 7.9 | 5 |
| 71 | Defining Acute on chronic liver failure: An identity crisis!. <i>Indian Journal of Gastroenterology</i> , 2010 , 29, 177-80 | 1.9 | 5 |
| 70 | Investigation of Oxidative Stress-Related Candidate Genes as Risk Factors for Drug-Induced Liver Injury due to Co-Amoxiclav. <i>DNA and Cell Biology</i> , 2020 , 39, 349-354 | 3.6 | 5 |
| 69 | Primary liver cancer in the UK: Incidence, incidence-based mortality, and survival by subtype, sex, and nation. <i>JHEP Reports</i> , 2021 , 3, 100232 | 10.3 | 5 |
| 68 | Accurate non-invasive diagnosis and staging of non-alcoholic fatty liver disease using the urinary steroid metabolome. <i>Alimentary Pharmacology and Therapeutics</i> , 2020 , 51, 1188-1197 | 6.1 | 5 |
| 67 | Determining a healthy reference range and factors potentially influencing PRO-C3 - A biomarker of liver fibrosis. <i>JHEP Reports</i> , 2021 , 3, 100317 | 10.3 | 5 |
| 66 | Low accuracy of FIB-4 and NAFLD fibrosis scores for screening for liver fibrosis in the population.. <i>Clinical Gastroenterology and Hepatology</i> , 2021 , | 6.9 | 5 |
| 65 | Jaundice: applying lessons from physiology. <i>Surgery</i> , 2009 , 27, 11-18 | 0.3 | 4 |
| 64 | Diffuse nesidioblastosis causing hyperinsulinemic hypoglycemia: the importance of pancreatic sampling on EUS. <i>Gastrointestinal Endoscopy</i> , 2008 , 68, 571-2; discussion 572 | 5.2 | 4 |
| 63 | Macrophage Scavenger Receptor 1 mediates lipid-induced inflammation in non-alcoholic fatty liver disease.. <i>Journal of Hepatology</i> , 2021 , | 13.4 | 4 |

| | | | |
|----|---|------|---|
| 62 | Increased serum miR-193a-5p during non-alcoholic fatty liver disease progression: Diagnostic and mechanistic relevance.. <i>JHEP Reports</i> , 2022 , 4, 100409 | 10.3 | 4 |
| 61 | □Monozygotic twins with NASH cirrhosis: cumulative effect of multiple single nucleotide polymorphisms?. <i>Annals of Hepatology</i> , 2016 , 15, 277-82 | 3.1 | 4 |
| 60 | A genetic risk score and diabetes predict development of alcohol-related cirrhosis in drinkers. <i>Journal of Hepatology</i> , 2021 , | 13.4 | 4 |
| 59 | Validation of the AASLD recommendations for classification of oesophageal varices in clinical practice. <i>Liver International</i> , 2020 , 40, 905-912 | 7.9 | 4 |
| 58 | HLA-DR polymorphism in SARS-CoV-2 infection and susceptibility to symptomatic COVID-19.. <i>Immunology</i> , 2022 , | 7.8 | 4 |
| 57 | Systematic Review with Meta-Analysis: Diagnostic Accuracy of Pro-C3 for Hepatic Fibrosis in Patients with Non-Alcoholic Fatty Liver Disease.. <i>Biomedicines</i> , 2021 , 9, | 4.8 | 4 |
| 56 | Long-Term Pioglitazone Treatment for Patients With Nonalcoholic Steatohepatitis. <i>Annals of Internal Medicine</i> , 2017 , 166, 229-230 | 8 | 3 |
| 55 | PS-112-Endoscopic duodenal mucosal resurfacing improves hepatic fat fraction, glycemic and lipid profiles in type 2 diabetes. <i>Journal of Hepatology</i> , 2019 , 70, e70-e71 | 13.4 | 3 |
| 54 | Immune-Mediated Drug-Induced Liver Injury 2014 , 401-412 | | 3 |
| 53 | Mind the gap. <i>ATLA Alternatives To Laboratory Animals</i> , 2010 , 38 Suppl 1, 1-4 | 2.1 | 3 |
| 52 | Concurrent mucinous cystadenoma of the liver and bile duct: a rare cause of jaundice. <i>Gastrointestinal Endoscopy</i> , 2006 , 63, 870-2 | 5.2 | 3 |
| 51 | Incidence and risk factors of anti-tuberculosis drug induced liver injury (DILI): Large cohort study involving 4652 Chinese adult tuberculosis patients. <i>Liver International</i> , 2021 , 41, 1565-1575 | 7.9 | 3 |
| 50 | MR Measures of Small Bowel Wall T2 Are Associated With Increased Permeability. <i>Journal of Magnetic Resonance Imaging</i> , 2021 , 53, 1422-1431 | 5.6 | 3 |
| 49 | Metabolic Imaging in Non-Alcoholic Fatty Liver Disease: Applications of Magnetic Resonance Spectroscopy. <i>Journal of Clinical Medicine</i> , 2021 , 10, | 5.1 | 3 |
| 48 | The XL probe: A luxury or a necessity? Risk stratification in an obese community cohort using transient elastography. <i>United European Gastroenterology Journal</i> , 2018 , 6, 1372-1379 | 5.3 | 3 |
| 47 | Drug-induced liver injury. <i>Medicine</i> , 2015 , 43, 590-593 | 0.6 | 2 |
| 46 | The Evaluation and Use of a Food Frequency Questionnaire Among the Population in Trivandrum, South Kerala, India. <i>Nutrients</i> , 2020 , 12, | 6.7 | 2 |
| 45 | Analysis of genotyping for predicting liver injury marker, procollagen III in persons at risk of non-alcoholic fatty liver disease. <i>Liver International</i> , 2018 , 38, 1832-1838 | 7.9 | 2 |

| | | | |
|----|--|------|---|
| 44 | Pretreatment Lesions on Magnetic Resonance Imaging in Patients With Hepatitis C Virus Infection Diagnosed With Hepatocellular Carcinoma After Initiating Direct-Acting Antiviral Therapy. <i>Gastroenterology</i> , 2018 , 154, 1848-1850 | 13.3 | 2 |
| 43 | Drug-induced liver injury. <i>Medicine</i> , 2019 , 47, 734-739 | 0.6 | 2 |
| 42 | Steering azathioprine safely: lessons from pharmacogenetics. <i>American Journal of Gastroenterology</i> , 2001 , 96, 3202-3 | 0.7 | 2 |
| 41 | EUS-guided choledochoduodenostomy with electrocautery-enhanced lumen-apposing metal stents in patients with malignant distal biliary obstruction: multicenter collaboration from the United Kingdom and Ireland. <i>Gastrointestinal Endoscopy</i> , 2021 , | 5.2 | 2 |
| 40 | What is the incidence of methotrexate or leflunomide discontinuation related to cytopenia, liver enzyme elevation or kidney function decline?. <i>Rheumatology</i> , 2021 , 60, 5785-5794 | 3.9 | 2 |
| 39 | Risk-stratified screening for chronic liver disease using vibration-controlled transient elastography (Fibroscan). <i>Gastrointestinal Nursing</i> , 2018 , 16, S15-S22 | 0.2 | 2 |
| 38 | Utility of palliative EUS-guided biliary drainage using lumen-apposing metal stents: a prospective multicenter feasibility study (with video). <i>Gastrointestinal Endoscopy</i> , 2021 , 94, 321-328 | 5.2 | 2 |
| 37 | Economic evaluation of covered stents for transjugular intrahepatic portosystemic stent shunt in patients with variceal bleeding and refractory ascites secondary to cirrhosis. <i>BMJ Open Gastroenterology</i> , 2021 , 8, | 3.9 | 2 |
| 36 | Drug Interactions in the Liver 2020 , 1050-1057 | | 1 |
| 35 | Jaundice: applying lessons from physiology. <i>Surgery</i> , 2011 , 29, 603-609 | 0.3 | 1 |
| 34 | Portal hypertension and ascites. <i>Surgery</i> , 2007 , 25, 28-33 | 0.3 | 1 |
| 33 | Acceptability of chronic liver disease screening in a UK primary care setting: a qualitative evaluation. <i>BMJ Open</i> , 2020 , 10, e041574 | 3 | 1 |
| 32 | Biomarkers of idiosyncratic drug-induced liver injury (DILI) - a systematic review. <i>Expert Opinion on Drug Metabolism and Toxicology</i> , 2021 , 17, 1327-1343 | 5.5 | 1 |
| 31 | Chronic liver disease in homeless individuals and performance of non-invasive liver fibrosis and injury markers: VALID study. <i>Liver International</i> , 2021 , | 7.9 | 1 |
| 30 | Macrophage Scavenger Receptor 1 mediates lipid-induced inflammation in non-alcoholic fatty liver disease | | 1 |
| 29 | Provision and standards of care for treatment and follow-up of patients with Autoimmune Hepatitis (AIH).. <i>Frontline Gastroenterology</i> , 2022 , 13, 126-132 | 2.6 | 1 |
| 28 | Reply to: "Hip fracture risk in patients with alcoholic cirrhosis: Do comorbidities and complications matter?". <i>Journal of Hepatology</i> , 2019 , 70, 213-214 | 13.4 | 1 |
| 27 | The Effect of Covid-19 on Alcohol Use Disorder and the Role of Universal Alcohol Screening in an Inpatient Setting: A Retrospective Cohort Control Study. <i>Alcohol and Alcoholism</i> , 2021 , | 3.5 | 1 |

| | | | |
|----|---|------|---|
| 26 | Hepatic ADRs429-443 | | 1 |
| 25 | Epidemiology of Cirrhosis and Hepatocellular Carcinoma344-356 | | 1 |
| 24 | Pro-adrenomedullin in acute decompensation of liver cirrhosis: relationship with acute-on-chronic liver failure and short-term survival. <i>Scandinavian Journal of Gastroenterology</i> , 2020 , 55, 606-614 | 2.4 | 0 |
| 23 | Jaundice: applying lessons from physiology. <i>Surgery</i> , 2014 , 32, 627-634 | 0.3 | 0 |
| 22 | Effects of an isoenergetic low Glycaemic Index (GI) diet on liver fat accumulation and gut microbiota composition in patients with non-alcoholic fatty liver disease (NAFLD): a study protocol of an efficacy mechanism evaluation. <i>BMJ Open</i> , 2021 , 11, e045802 | 3 | 0 |
| 21 | Gut Microbial Profile Is Associated With Residential Settings and Not Nutritional Status in Adults in Karnataka, India. <i>Frontiers in Nutrition</i> , 2021 , 8, 595756 | 6.2 | 0 |
| 20 | Large volume paracentesis: to do or where to do?. <i>Gut</i> , 2021 , 70, 2401-2402 | 19.2 | 0 |
| 19 | Estimating the clinical prevalence of Wilson's disease in the UK. <i>JHEP Reports</i> , 2021 , 3, 100329 | 10.3 | 0 |
| 18 | Long-term outcomes of liver transplant recipients followed up in non-transplant centres: Care closer to home. <i>Clinical Medicine</i> , 2021 , 21, e32-e38 | 1.9 | 0 |
| 17 | The effect of exercise training on adipose tissue insulin sensitivity: A systematic review and meta-analysis.. <i>Obesity Reviews</i> , 2022 , e13445 | 10.6 | 0 |
| 16 | Prognostic non-invasive biomarkers for all-cause mortality in non-alcoholic fatty liver disease: A systematic review and meta-analysis. <i>World Journal of Hepatology</i> , 2022 , 14, 1025-1037 | 3.4 | 0 |
| 15 | Drug-Induced Cholestasis: Mechanisms and Importance 2017 , 117-128 | | |
| 14 | Commentary: non-alcoholic steatohepatitis-finding and minding the fire behind the smoke. <i>Alimentary Pharmacology and Therapeutics</i> , 2020 , 51, 209-211 | 6.1 | |
| 13 | Portal hypertension and ascites. <i>Surgery</i> , 2014 , 32, 661-667 | 0.3 | |
| 12 | Hepatic Adverse Drug Reactions 2014 , 539-555 | | |
| 11 | To feed or what to feed in cirrhosis?. <i>Hepatology</i> , 2012 , 56, 1590 | 11.2 | |
| 10 | Portal hypertension and ascites. <i>Surgery</i> , 2011 , 29, 640-646 | 0.3 | |
| 9 | Portal hypertension and ascites. <i>Surgery</i> , 2010 , 28, 234-240 | 0.3 | |

- 8 Comparison of the reverse bevel versus Franseen type endoscopic ultrasound needle. *World Journal of Gastrointestinal Endoscopy*, **2020**, 12, 266-275 2.2
- 7 Pharmacogenetics in Gastroenterology **2006**, 273-294
- 6 Immune-Mediated Drug-Induced Liver Injury **2020**, 491-504
- 5 Hepatic Adverse Drug Reactions 459-474
- 4 Magnetic Resonance Imaging Methods for Assessing Cirrhosis and Portal Hypertension **2018**, 211-223
- 3 Monoclonal antibody BTT1023 targeting vascular adhesion protein 1 for treating primary sclerosing cholangitis: BUTE0 single-arm Phase II trial. *Efficacy and Mechanism Evaluation*, **2022**, 9, 1-54 1.7
- 2 Characteristics of alcohol recovery narratives: Systematic review and narrative synthesis.. *PLoS ONE*, **2022**, 17, e0268034 3.7
- 1 Incidence and prevalence of venous thromboembolism in chronic liver disease: A systematic review and meta-analysis. *Thrombosis Research*, **2022**, 215, 19-29 8.2