

# Zhening Liu

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/906008/publications.pdf>

Version: 2024-02-01

137  
papers

7,128  
citations

93792

39  
h-index

73587

79  
g-index

142  
all docs

142  
docs citations

142  
times ranked

8740  
citing authors

#	ARTICLE	IF	CITATIONS
1	The centrality of working memory networks in differentiating bipolar type I depression from unipolar depression: A task-fMRI study. <i>Canadian Journal of Psychiatry</i> , 2023, 68, 22-32.	0.9	4
2	Imbalance Between Prefronto-Thalamic and Sensorimotor-Thalamic Circuitries Associated with Working Memory Deficit in Schizophrenia. <i>Schizophrenia Bulletin</i> , 2022, 48, 251-261.	2.3	15
3	Abnormal network properties and fiber connections of DMN across major mental disorders: a probability tracing and graph theory study. <i>Cerebral Cortex</i> , 2022, 32, 3127-3136.	1.6	7
4	Dynamical regional activity in putamen distinguishes bipolar type I depression and unipolar depression. <i>Journal of Affective Disorders</i> , 2022, 297, 94-101.	2.0	6
5	Mental health characteristics and their associations with childhood trauma among subgroups of people living with HIV in China. <i>BMC Psychiatry</i> , 2022, 22, 13.	1.1	4
6	Overprotection and overcontrol in childhood: An evaluation on reliability and validity of 33-item expanded Childhood Trauma Questionnaire (CTQ-33), Chinese version. <i>Asian Journal of Psychiatry</i> , 2022, 68, 102962.	0.9	17
7	Changes in brain network properties in major depressive disorder following electroconvulsive therapy: a combined static and dynamic functional magnetic resonance imaging study. <i>Annals of Palliative Medicine</i> , 2022, 11, 1969-1980.	0.5	6
8	Factors influencing medication adherence among patients with severe mental disorders from the perspective of mental health professionals. <i>BMC Psychiatry</i> , 2022, 22, 22.	1.1	14
9	Aberrant integrity of the cortico-limbic-striatal circuit in major depressive disorder with suicidal ideation. <i>Journal of Psychiatric Research</i> , 2022, 148, 277-285.	1.5	9
10	Increasing Trends in Mental Health Problems Among Urban Chinese Adolescents: Results From Repeated Cross-Sectional Data in Changsha 2016â€“2020. <i>Frontiers in Public Health</i> , 2022, 10, 829674.	1.3	8
11	Association Between Wisdom and Psychotic-Like Experiences in the General Population: A Cross-Sectional Study. <i>Frontiers in Psychiatry</i> , 2022, 13, 814242.	1.3	2
12	Effectiveness and safety of blonanserin for improving social and cognitive functions in patients with first-episode schizophrenia: a study protocol for a prospective, multicentre, single-arm clinical trial. <i>BMJ Open</i> , 2022, 12, e054079.	0.8	2
13	Multimetric structural covariance in first-episode major depressive disorder: a graph theoretical analysis. <i>Journal of Psychiatry and Neuroscience</i> , 2022, 47, E176-E185.	1.4	6
14	Treatment Effect of Long-Term Antipsychotics on Default-Mode Network Dysfunction in Drug-Naïve Patients With First-Episode Schizophrenia: A Longitudinal Study. <i>Frontiers in Pharmacology</i> , 2022, 13, .	1.6	2
15	Associations between polygenic risk, negative symptoms, and functional connectome topology during a working memory task in early-onset schizophrenia. <i>NPJ Schizophrenia</i> , 2022, 8, .	2.0	3
16	Drug-naïve first-episode schizophrenia spectrum disorders: Pharmacological treatment practices in inpatient units in Hunan Province, China. <i>Microbial Biotechnology</i> , 2021, 15, 1010-1018.	0.9	2
17	Abnormal Thalamocortical Circuit in Adolescents With Early-Onset Schizophrenia. <i>Journal of the American Academy of Child and Adolescent Psychiatry</i> , 2021, 60, 479-489.	0.3	31
18	Decreased integration of the frontoparietal network during a working memory task in major depressive disorder. <i>Australian and New Zealand Journal of Psychiatry</i> , 2021, 55, 577-587.	1.3	24

#	ARTICLE	IF	CITATIONS
19	Gray matter reorganization underpinnings of antidepressant treatment of persistent depressive disorder. <i>European Neuropsychopharmacology</i> , 2021, 43, 129-138.	0.3	6
20	Abnormal Large-Scale Network Activation Present in Bipolar Mania and Bipolar Depression Under Resting State. <i>Frontiers in Psychiatry</i> , 2021, 12, 634299.	1.3	9
21	Online Real-Time Monitoring of Exhaled Breath Particles Reveals Unnoticed Transport of Nonvolatile Drugs from Blood to Breath. <i>Analytical Chemistry</i> , 2021, 93, 5005-5008.	3.2	13
22	The Role of Military Training in Improving Psychological Resilience and Reducing Depression Among College Freshmen. <i>Frontiers in Psychiatry</i> , 2021, 12, 641396.	1.3	9
23	Evaluation on Long-Term Test-Retest Reliability of the Short-Form Childhood Trauma Questionnaire in Patients with Schizophrenia. <i>Psychology Research and Behavior Management</i> , 2021, Volume 14, 1033-1040.	1.3	14
24	Childhood trauma is linked to decreased temporal stability of functional brain networks in young adults. <i>Journal of Affective Disorders</i> , 2021, 290, 23-30.	2.0	27
25	Acute Effect of Betel Quid Chewing on Brain Network Dynamics: A Resting-State Functional Magnetic Resonance Imaging Study. <i>Frontiers in Psychiatry</i> , 2021, 12, 701420.	1.3	5
26	The integrity of the white matter in first-episode schizophrenia patients with auditory verbal hallucinations: An atlas-based DTI analysis. <i>Psychiatry Research - Neuroimaging</i> , 2021, 315, 111328.	0.9	7
27	Associations between brain abnormalities and common genetic variants for schizophrenia: a narrative review of structural and functional neuroimaging findings. <i>Annals of Palliative Medicine</i> , 2021, 10, 10031-10052.	0.5	6
28	Changes of psychotic-like experiences and their association with anxiety/depression among young adolescents before COVID-19 and after the lockdown in China. <i>Schizophrenia Research</i> , 2021, 237, 40-46.	1.1	41
29	Family functioning as a moderator in the relation between perceived stress and psychotic-like experiences among adolescents during COVID-19. <i>Comprehensive Psychiatry</i> , 2021, 111, 152274.	1.5	18
30	Decreased Cortical Folding of the Fusiform Gyrus and Its Hypoconnectivity with Sensorimotor Areas in Major Depressive Disorder. <i>Journal of Affective Disorders</i> , 2021, 295, 657-664.	2.0	23
31	Age-Related Decrease in Default-Mode Network Functional Connectivity Is Accelerated in Patients With Major Depressive Disorder. <i>Frontiers in Aging Neuroscience</i> , 2021, 13, 809853.	1.7	8
32	Differential Dynamical Pattern of Regional Homogeneity in Bipolar and Unipolar Depression: A Preliminary Resting-State fMRI Study. <i>Frontiers in Psychiatry</i> , 2021, 12, 764932.	1.3	8
33	Altered resting-state dynamic functional brain networks in major depressive disorder: Findings from the REST-meta-MDD consortium. <i>NeuroImage: Clinical</i> , 2020, 26, 102163.	1.4	76
34	Brain-Wide Functional Dysconnectivity in Schizophrenia: Parsing Diathesis, Resilience, and the Effects of Clinical Expression. <i>Canadian Journal of Psychiatry</i> , 2020, 65, 21-29.	0.9	14
35	Morphological Profiling of Schizophrenia: Cluster Analysis of MRI-Based Cortical Thickness Data. <i>Schizophrenia Bulletin</i> , 2020, 46, 623-632.	2.3	35
36	S209. DRUG-NAÏVE FIRST-EPISEODE SCHIZOPHRENIA SPECTRUM DISORDERS: PHARMACOLOGICAL TREATMENT PRACTICES IN INPATIENT UNITS IN HUNAN PROVINCE, CHINA. <i>Schizophrenia Bulletin</i> , 2020, 46, S118-S118.	2.3	0

#	ARTICLE	IF	CITATIONS
37	Reduced Cortical Thickness in the Right Caudal Middle Frontal Is Associated With Symptom Severity in Betel Quid-Dependent Chewers. <i>Frontiers in Psychiatry</i> , 2020, 11, 654.	1.3	3
38	A multivariate pattern analysis of resting-state functional MRI data in Na <sup>+</sup> and chronic betel quid chewers. <i>Brain Imaging and Behavior</i> , 2020, 15, 1222-1234.	1.1	4
39	Bipolar Mood State Reflected in Functional Connectivity of the Hate Circuit: A Resting-State Functional Magnetic Resonance Imaging Study. <i>Frontiers in Psychiatry</i> , 2020, 11, 556126.	1.3	8
40	Altered Temporal Variability of Local and Large-Scale Resting-State Brain Functional Connectivity Patterns in Schizophrenia and Bipolar Disorder. <i>Frontiers in Psychiatry</i> , 2020, 11, 422.	1.3	38
41	Connectomic signatures of working memory deficits in depression, mania, and euthymic states of bipolar disorder. <i>Journal of Affective Disorders</i> , 2020, 274, 190-198.	2.0	22
42	Medication adherence and its correlates among patients affected by schizophrenia with an episodic course: A large-scale multi-center cross-sectional study in China. <i>Asian Journal of Psychiatry</i> , 2020, 53, 102198.	0.9	8
43	Connectomic Underpinnings of Working Memory Deficits in Schizophrenia: Evidence From a replication fMRI study. <i>Schizophrenia Bulletin</i> , 2020, 46, 916-926.	2.3	27
44	Acute and Chronic Effects of Betel Quid Chewing on Brain Functional Connectivity. <i>Frontiers in Psychiatry</i> , 2020, 11, 198.	1.3	9
45	Saliency-thalamic circuit uncouples in major depressive disorder, but not in bipolar depression. <i>Journal of Affective Disorders</i> , 2020, 269, 43-50.	2.0	9
46	“Brain Connectivity Deviates by Sex and Hemisphere in the First Episode of Schizophrenia” A Route to the Genetic Basis of Language and Psychosis?. <i>Schizophrenia Bulletin</i> , 2019, 45, 484-494.	2.3	19
47	Crosstalk Between Inflammation and Glutamate System in Depression: Signaling Pathway and Molecular Biomarkers for Ketamine’s Antidepressant Effect. <i>Molecular Neurobiology</i> , 2019, 56, 3484-3500.	1.9	59
48	A Qualitative Study of Implementation Challenges of Mental Health Clubhouse Rehabilitation Services in China’s Hunan Province. <i>Psychiatric Services</i> , 2019, 70, 674-680.	1.1	4
49	Structural and Functional Alterations in Betel-Quid Chewers: A Systematic Review of Neuroimaging Findings. <i>Frontiers in Psychiatry</i> , 2019, 10, 16.	1.3	15
50	Longitudinal Trajectories of Psychotic-Like Experiences and Their Relationship to Emergent Mental Disorders Among Adolescents. <i>Journal of Clinical Psychiatry</i> , 2019, 80, .	1.1	14
51	Psychological resilience negatively correlates with resting-state brain network flexibility in young healthy adults: a dynamic functional magnetic resonance imaging study. <i>Annals of Translational Medicine</i> , 2019, 7, 809-809.	0.7	39
52	Abnormal Connectivity Within Anterior Cortical Midline Structures in Bipolar Disorder: Evidence From Integrated MRI and Functional MRI. <i>Frontiers in Psychiatry</i> , 2019, 10, 788.	1.3	10
53	Association between migrant worker experience, limitations on insurance coverage, and hospitalization for schizophrenia in Hunan Province, China. <i>Schizophrenia Research</i> , 2018, 197, 93-97.	1.1	12
54	Multi-Site Diagnostic Classification of Schizophrenia Using Discriminant Deep Learning with Functional Connectivity MRI. <i>EBioMedicine</i> , 2018, 30, 74-85.	2.7	206

#	ARTICLE	IF	CITATIONS
55	Attitudes towards mental illness among medical students in China: Impact of medical education on stigma. <i>Asia-Pacific Psychiatry</i> , 2018, 10, e12294.	1.2	19
56	The instability of functional connectivity in patients with schizophrenia and their siblings: A dynamic connectivity study. <i>Schizophrenia Research</i> , 2018, 195, 183-189.	1.1	28
57	Associations Among Suicidal Ideation, White Matter Integrity and Cognitive Deficit in First-Episode Schizophrenia. <i>Frontiers in Psychiatry</i> , 2018, 9, 391.	1.3	43
58	Effects of DISC1 Polymorphisms on Resting-State Spontaneous Neuronal Activity in the Early-Stage of Schizophrenia. <i>Frontiers in Psychiatry</i> , 2018, 9, 137.	1.3	12
59	Resilience and Cognitive Function in Patients With Schizophrenia and Bipolar Disorder, and Healthy Controls. <i>Frontiers in Psychiatry</i> , 2018, 9, 279.	1.3	41
60	The Psychiatry Major: A Curricular Innovation to Improve Undergraduate Psychiatry Education in China. <i>Academic Psychiatry</i> , 2018, 42, 376-381.	0.4	8
61	Brain-Wide Analysis of Functional Connectivity in First-Episode and Chronic Stages of Schizophrenia. <i>Schizophrenia Bulletin</i> , 2017, 43, sbw099.	2.3	142
62	Betel quid chewing alters functional connectivity in frontal and default networks: A resting-state fMRI study. <i>Journal of Magnetic Resonance Imaging</i> , 2017, 45, 157-166.	1.9	24
63	Task and resting-state fMRI studies in first-episode schizophrenia: A systematic review. <i>Schizophrenia Research</i> , 2017, 189, 9-18.	1.1	99
64	Working Memory Modulation of Frontoparietal Network Connectivity in First-Episode Schizophrenia. <i>Cerebral Cortex</i> , 2017, 27, 3832-3841.	1.6	49
65	Psychotic-like experiences and correlation with childhood trauma and other socio-demographic factors: A cross-sectional survey in adolescence and early adulthood in China. <i>Psychiatry Research</i> , 2017, 255, 272-277.	1.7	35
66	Expanding the Mental Health Workforce in China: Narrowing the Mental Health Service Gap. <i>Psychiatric Services</i> , 2017, 68, 987-989.	1.1	31
67	Psychotic-like experiences, trauma and related risk factors among "left-behind" children in China. <i>Schizophrenia Research</i> , 2017, 181, 43-48.	1.1	51
68	Altered Brain Functional Connectivity in Betel Quid-Dependent Chewers. <i>Frontiers in Psychiatry</i> , 2017, 8, 239.	1.3	16
69	Decreased Left Putamen and Thalamus Volume Correlates with Delusions in First-Episode Schizophrenia Patients. <i>Frontiers in Psychiatry</i> , 2017, 8, 245.	1.3	40
70	Gray matter morphological anomalies in the cerebellar vermis in first-episode schizophrenia patients with cognitive deficits. <i>BMC Psychiatry</i> , 2017, 17, 374.	1.1	15
71	Identification of RELN variation p.Thr3192Ser in a Chinese family with schizophrenia. <i>Scientific Reports</i> , 2016, 6, 24327.	1.6	27
72	Inefficient DMN Suppression in Schizophrenia Patients with Impaired Cognitive Function but not Patients with Preserved Cognitive Function. <i>Scientific Reports</i> , 2016, 6, 21657.	1.6	51

#	ARTICLE	IF	CITATIONS
73	Heredity characteristics of schizophrenia shown by dynamic functional connectivity analysis of resting-state functional MRI scans of unaffected siblings. <i>NeuroReport</i> , 2016, 27, 843-848.	0.6	34
74	Bipolar diagnosis in China: Evaluating diagnostic confidence using the Bipolarity Index. <i>Journal of Affective Disorders</i> , 2016, 202, 247-253.	2.0	9
75	Abnormally increased and incoherent resting-state activity is shared between patients with schizophrenia and their unaffected siblings. <i>Schizophrenia Research</i> , 2016, 171, 158-165.	1.1	44
76	Decreased functional connectivity in the language regions in bipolar patients during depressive episodes but not remission. <i>Journal of Affective Disorders</i> , 2016, 197, 116-124.	2.0	45
77	Perceived barriers on utilization of mental health services among adults in Dodoma Municipality " Tanzania. <i>Journal of Public Mental Health</i> , 2015, 14, 79-93.	0.8	6
78	Psychotic-like experiences and associated socio-demographic factors among adolescents in China. <i>Schizophrenia Research</i> , 2015, 166, 49-54.	1.1	37
79	Grey matter morphological anomalies in the caudate head in first-episode psychosis patients with delusions of reference. <i>Psychiatry Research - Neuroimaging</i> , 2015, 233, 57-63.	0.9	25
80	Major depressive disorder in hemodialysis patients in China. <i>Asia-Pacific Psychiatry</i> , 2015, 7, 78-84.	1.2	7
81	Reduced cortical thickness in right Heschl's gyrus associated with auditory verbal hallucinations severity in first-episode schizophrenia. <i>BMC Psychiatry</i> , 2015, 15, 152.	1.1	38
82	Anatomical Distance Affects Functional Connectivity in Patients With Schizophrenia and Their Siblings. <i>Schizophrenia Bulletin</i> , 2014, 40, 449-459.	2.3	33
83	A Brain-wide association study of DISC1 genetic variants reveals a relationship with the structure and functional connectivity of the precuneus in schizophrenia. <i>Human Brain Mapping</i> , 2014, 35, 5414-5430.	1.9	27
84	Aberrant functional connectivity for diagnosis of major depressive disorder: A discriminant analysis. <i>Psychiatry and Clinical Neurosciences</i> , 2014, 68, 110-119.	1.0	46
85	Altered functional connectivity links in neuroleptic-naïve and neuroleptic-treated patients with schizophrenia, and their relation to symptoms including volition. <i>NeuroImage: Clinical</i> , 2014, 6, 463-474.	1.4	24
86	Internetwork dynamic connectivity effectively differentiates schizophrenic patients from healthy controls. <i>NeuroReport</i> , 2014, 25, 1344-1349.	0.6	24
87	Altered default mode and fronto-parietal network subsystems in patients with schizophrenia and their unaffected siblings. <i>Brain Research</i> , 2014, 1562, 87-99.	1.1	71
88	Hyper-coupling between working memory task-evoked activations and amplitude of spontaneous fluctuations in first-episode schizophrenia. <i>Schizophrenia Research</i> , 2014, 159, 80-89.	1.1	18
89	Effective connectivity of the posterior cingulate and medial prefrontal cortices relates to working memory impairment in schizophrenic and bipolar patients. <i>Schizophrenia Research</i> , 2014, 158, 85-90.	1.1	41
90	A comparative diffusion tensor imaging study of corpus callosum subregion integrity in bipolar disorder and schizophrenia. <i>Psychiatry Research - Neuroimaging</i> , 2014, 221, 58-62.	0.9	50

#	ARTICLE	IF	CITATIONS
91	Brain-wide functional inter-hemispheric disconnection is a potential biomarker for schizophrenia and distinguishes it from depression. <i>NeuroImage: Clinical</i> , 2013, 2, 818-826.	1.4	62
92	Detection of malingering: psychometric evaluation of the Chinese version of the structured interview of reported symptoms-2. <i>BMC Psychiatry</i> , 2013, 13, 254.	1.1	9
93	Long- and short-term inpatients with schizophrenia in China: Implications for community-based service development. <i>Asia-Pacific Psychiatry</i> , 2013, 5, E39-46.	1.2	6
94	Comparison of first-episode and chronic patients diagnosed with schizophrenia: symptoms and childhood trauma. <i>Microbial Biotechnology</i> , 2013, 7, 23-30.	0.9	42
95	Comparison of Psychosocial Determinants in Inpatients With First-Episode and Chronic Schizophrenia in China. <i>Archives of Psychiatric Nursing</i> , 2013, 27, 32-41.	0.7	6
96	Abnormal amplitude low-frequency oscillations in medication-naive, first-episode patients with major depressive disorder: A resting-state fMRI study. <i>Journal of Affective Disorders</i> , 2013, 146, 401-406.	2.0	231
97	Decreased left middle temporal gyrus volume in antipsychotic drug-naive, first-episode schizophrenia patients and their healthy unaffected siblings. <i>Schizophrenia Research</i> , 2013, 144, 37-42.	1.1	84
98	The diminished interhemispheric connectivity correlates with negative symptoms and cognitive impairment in first-episode schizophrenia. <i>Schizophrenia Research</i> , 2013, 150, 144-150.	1.1	36
99	Abnormal resting-state cerebellar-cerebral functional connectivity in treatment-resistant depression and treatment sensitive depression. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2013, 44, 51-57.	2.5	160
100	Impaired interhemispheric connectivity in medication-naive patients with major depressive disorder. <i>Journal of Psychiatry and Neuroscience</i> , 2013, 38, 43-48.	1.4	56
101	Investigation on the influence of a didactic course in psychiatry on attitudes of mental illness in Chinese college students. <i>Asia-Pacific Psychiatry</i> , 2013, 5, 147-151.	1.2	8
102	Opposite Effective Connectivity in the Posterior Cingulate and Medial Prefrontal Cortex between First-Episode Schizophrenic Patients with Suicide Risk and Healthy Controls. <i>PLoS ONE</i> , 2013, 8, e63477.	1.1	32
103	Decreased Interhemispheric Coordination in Treatment-Resistant Depression: A Resting-State fMRI Study. <i>PLoS ONE</i> , 2013, 8, e71368.	1.1	68
104	Schizophrenic Patients and Their Unaffected Siblings Share Increased Resting-State Connectivity in the Task-Negative Network but Not Its Anticorrelated Task-Positive Network. <i>Schizophrenia Bulletin</i> , 2012, 38, 285-294.	2.3	116
105	Potential risk for healthy siblings to develop schizophrenia. <i>NeuroReport</i> , 2012, 23, 265-269.	0.6	21
106	Right lateralized white matter abnormalities in first-episode, drug-naive paranoid schizophrenia. <i>Neuroscience Letters</i> , 2012, 531, 5-9.	1.0	85
107	Effects of antipsychotic medications on quality of life and psychosocial functioning in patients with early-stage schizophrenia: 1-year follow-up naturalistic study. <i>Comprehensive Psychiatry</i> , 2012, 53, 1006-1012.	1.5	13
108	Morphological and functional abnormalities of salience network in the early-stage of paranoid schizophrenia. <i>Schizophrenia Research</i> , 2012, 141, 15-21.	1.1	57

#	ARTICLE	IF	CITATIONS
109	Genetic variation in the calcium/calmodulin-dependent protein kinase (CaMK) pathway is associated with antidepressant response in females. <i>Journal of Affective Disorders</i> , 2012, 136, 558-566.	2.0	26
110	Classification of Different Therapeutic Responses of Major Depressive Disorder with Multivariate Pattern Analysis Method Based on Structural MR Scans. <i>PLoS ONE</i> , 2012, 7, e40968.	1.1	125
111	Discriminant analysis of functional connectivity patterns on Grassmann manifold. <i>NeuroImage</i> , 2011, 56, 2058-2067.	2.1	78
112	Schizophrenic Patients and Their Unaffected Siblings Show Similar Abnormalities in Resting-State Functional Connectivity. <i>Procedia Environmental Sciences</i> , 2011, 8, 715-722.	1.3	1
113	A study of schizophrenia inheritance through pattern classification. , 2011, , .		2
114	Complementary diffusion tensor imaging study of the corpus callosum in patients with first-episode and chronic schizophrenia. <i>Journal of Psychiatry and Neuroscience</i> , 2011, 36, 120-125.	1.4	61
115	Bilateral functional asymmetry disparity in positive and negative schizophrenia revealed by resting-state fMRI. <i>Psychiatry Research - Neuroimaging</i> , 2010, 182, 30-39.	0.9	34
116	Effect of Antipsychotic Medication Alone vs Combined With Psychosocial Intervention on Outcomes of Early-Stage Schizophrenia. <i>Archives of General Psychiatry</i> , 2010, 67, 895.	13.8	124
117	Discriminant analysis of resting-state functional connectivity patterns on the Grassmann manifold. , 2010, , .		4
118	Frontal and cingulate gray matter volume reduction in heroin dependence: Optimized voxel-based morphometry. <i>Psychiatry and Clinical Neurosciences</i> , 2009, 63, 563-568.	1.0	91
119	Schizophrenia patients and their healthy siblings share disruption of white matter integrity in the left prefrontal cortex and the hippocampus but not the anterior cingulate cortex. <i>Schizophrenia Research</i> , 2009, 114, 128-135.	1.1	81
120	Abnormal neural activity in the patients with remitted geriatric depression: A resting-state functional magnetic resonance imaging study. <i>Journal of Affective Disorders</i> , 2008, 111, 145-152.	2.0	122
121	Altered resting-state functional connectivity and anatomical connectivity of hippocampus in schizophrenia. <i>Schizophrenia Research</i> , 2008, 100, 120-132.	1.1	289
122	Regional Gray Matter Changes Are Associated with Cognitive Deficits in Remitted Geriatric Depression: An Optimized Voxel-Based Morphometry Study. <i>Biological Psychiatry</i> , 2008, 64, 541-544.	0.7	80
123	Distinct neurobehavioral consequences of prenatal exposure to sulpiride (SUL) and risperidone (RIS) in rats. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2008, 32, 387-397.	2.5	28
124	Disrupted White Matter Integrity in Heroin Dependence: A Controlled Study Utilizing Diffusion Tensor Imaging. <i>American Journal of Drug and Alcohol Abuse</i> , 2008, 34, 562-575.	1.1	76
125	Disrupted small-world networks in schizophrenia. <i>Brain</i> , 2008, 131, 945-961.	3.7	944
126	Functional dysconnectivity of the dorsolateral prefrontal cortex in first-episode schizophrenia using resting-state fMRI. <i>Neuroscience Letters</i> , 2007, 417, 297-302.	1.0	286



#	ARTICLE	IF	CITATIONS
127	Functional disintegration in paranoid schizophrenia using resting-state fMRI. <i>Schizophrenia Research</i> , 2007, 97, 194-205.	1.1	384
128	Regional Homogeneity and Anatomical Parcellation for fMRI Image Classification: Application to Schizophrenia and Normal Controls. , 2007, 10, 136-143.		19
129	Antipsychotic Combination with Psychosocial Intervention on Outcome of Schizophrenia (ACPIOS): Rationale and Design of the Clinical Trial. <i>Clinical Schizophrenia and Related Psychoses</i> , 2007, 1, 185-192.	1.4	2
130	The Effects of Clozapine on Cognitive Function and Regional Cerebral Blood Flow in the Negative Symptom Profile Schizophrenia. <i>International Journal of Psychiatry in Medicine</i> , 2006, 36, 171-181.	0.8	21
131	White matter integrity of the whole brain is disrupted in first-episode schizophrenia. <i>NeuroReport</i> , 2006, 17, 23-26.	0.6	129
132	Decreased regional homogeneity in schizophrenia: a resting state functional magnetic resonance imaging study. <i>NeuroReport</i> , 2006, 17, 19-22.	0.6	237
133	Widespread functional disconnectivity in schizophrenia with resting-state functional magnetic resonance imaging. <i>NeuroReport</i> , 2006, 17, 209-213.	0.6	381
134	Positive and negative symptom profile schizophrenia and abnormalities in the P300 component of the event-related potential: a longitudinal controlled study. <i>Psychiatry Research - Neuroimaging</i> , 2004, 132, 131-139.	0.9	29
135	Comparison of Neurocognition Between Drug-Free Patients With Schizophrenia and Bipolar Disorder. <i>Journal of Nervous and Mental Disease</i> , 2004, 192, 464-470.	0.5	28
136	The relationship between regional cerebral blood flow and the Wisconsin Card Sorting Test in negative schizophrenia. <i>Psychiatry and Clinical Neurosciences</i> , 2002, 56, 3-7.	1.0	32
137	The Role of Family Function in Reducing Suicidal Ideation Mediated by COVID-Related Stress and Psychotic-Like Experiences Among Adolescents. <i>SSRN Electronic Journal</i> , 0, , .	0.4	1