

# Melanie Luppa

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9061359/publications.pdf>

Version: 2024-02-01

93  
papers

4,983  
citations

136950

32  
h-index

98798

67  
g-index

114  
all docs

114  
docs citations

114  
times ranked

6538  
citing authors

#	ARTICLE	IF	CITATIONS
1	Do oldest old individuals perceive receipt of informal care as a restriction or support of their autonomy?. <i>Aging and Mental Health</i> , 2022, 26, 1862-1873.	2.8	1
2	Factors Leading to Institutionalization among the Oldest Old: Longitudinal Findings from the AgeCoDe-AgeQualiDe Study. <i>Gerontology</i> , 2022, 68, 894-902.	2.8	6
3	Physical Activity Determinants in Older German Adults at Increased Dementia Risk with Multimorbidity: Baseline Results of the AgeWell.de Study. <i>International Journal of Environmental Research and Public Health</i> , 2022, 19, 3164.	2.6	6
4	Factors Contributing to Persistent Frequent Attendance in Primary Care Among the Oldest Old: Longitudinal Evidence From the AgeCoDe-AgeQualiDe Study. <i>Frontiers in Medicine</i> , 2022, 9, 815419.	2.6	3
5	Psychometric Evaluation of the German Version of the Social Functioning in Dementia Scale (SF-DEM). <i>Journal of Alzheimer's Disease</i> , 2022, 86, 1231-1241.	2.6	1
6	Incidence and risk factors of depressive symptoms in the highest age groups and competing mortality risk. Evidence from the AgeCoDe-AgeQualiDe prospective cohort study. <i>Journal of Affective Disorders</i> , 2022, 308, 494-501.	4.1	2
7	What Comes after the Trial? An Observational Study of the Real-World Uptake of an E-Mental Health Intervention by General Practitioners to Reduce Depressive Symptoms in Their Patients. <i>International Journal of Environmental Research and Public Health</i> , 2022, 19, 6203.	2.6	3
8	Depressive Symptoms and Frailty Among the Oldest Old: Evidence From a Multicenter Prospective Study. <i>Journal of the American Medical Directors Association</i> , 2021, 22, 577-582.e2.	2.5	4
9	Recruitment and Baseline Characteristics of Participants in the AgeWell.de Study—A Pragmatic Cluster-Randomized Controlled Lifestyle Trial against Cognitive Decline. <i>International Journal of Environmental Research and Public Health</i> , 2021, 18, 408.	2.6	15
10	GPs' awareness of car driving among oldest patients: exploratory results from a primary care cohort. <i>BJGP Open</i> , 2021, 5, BJGPO.2020.0145.	1.8	1
11	Frailty and Autonomy among the Oldest Old: Evidence from the Multicenter Prospective AgeCoDe-AgeQualiDe Study. <i>Gerontology</i> , 2021, 67, 591-598.	2.8	3
12	Using a brief web-based 5A intervention to improve weight management in primary care: results of a cluster-randomized controlled trial. <i>BMC Family Practice</i> , 2021, 22, 61.	2.9	9
13	Carers' distress. <i>International Psychogeriatrics</i> , 2021, 33, 311-313.	1.0	1
14	Correlates of institutionalization among the oldest old—Evidence from the multicenter AgeCoDe-AgeQualiDe study. <i>International Journal of Geriatric Psychiatry</i> , 2021, 36, 1095-1102.	2.7	13
15	Risk factors and protective factors of depression in older people 65+. A systematic review. <i>PLoS ONE</i> , 2021, 16, e0251326.	2.5	79
16	The Role of Social Isolation and the Development of Depression. A Comparison of the Widowed and Married Oldest Old in Germany. <i>International Journal of Environmental Research and Public Health</i> , 2021, 18, 6986.	2.6	16
17	Association of Alcohol and Tobacco Consumption with Depression Severity in the Oldest Old. Results from the Age Different Old Age Cohort Platform. <i>International Journal of Environmental Research and Public Health</i> , 2021, 18, 7959.	2.6	4
18	Mental Health Self-Stigma of Syrian Refugees With Posttraumatic Stress Symptoms: Investigating Sociodemographic and Psychopathological Correlates. <i>Frontiers in Psychiatry</i> , 2021, 12, 642618.	2.6	10

#	ARTICLE	IF	CITATIONS
19	Gender-specific design and effectiveness of non-pharmacological interventions against cognitive decline and dementia – protocol for a systematic review and meta-analysis. <i>PLoS ONE</i> , 2021, 16, e0256826.	2.5	1
20	Excess costs of dementia in old age (85+) in Germany: Results from the AgeCoDe-AgeQualiDe study. <i>Journal of the Economics of Ageing</i> , 2021, 20, 100346.	1.3	3
21	Loss and bereavement in late life (60+): Study protocol for a randomized controlled trial regarding an internet-based self-help intervention. <i>Internet Interventions</i> , 2021, 26, 100451.	2.7	3
22	Incidence of Anxiety in Latest Life and Risk Factors. Results of the AgeCoDe/AgeQualiDe Study. <i>International Journal of Environmental Research and Public Health</i> , 2021, 18, 12786.	2.6	7
23	Effectiveness and Feasibility of Internet-Based Interventions for Grief After Bereavement: Systematic Review and Meta-analysis. <i>JMIR Mental Health</i> , 2021, 8, e29661.	3.3	17
24	Association of mental demands in the workplace with cognitive function in older adults at increased risk for dementia. <i>BMC Geriatrics</i> , 2021, 21, 688.	2.7	1
25	Social support and health-related quality of life among the oldest old – longitudinal evidence from the multicenter prospective AgeCoDe-AgeQualiDe study. <i>Quality of Life Research</i> , 2021, , 1.	3.1	6
26	Late-Life Depressive Symptoms Are Associated With Functional Impairment Cross-sectionally and Over Time: Results of the AgeMooDe Study. <i>Journals of Gerontology - Series B Psychological Sciences and Social Sciences</i> , 2020, 75, 811-820.	3.9	6
27	Effectiveness and feasibility of internet-based and mobile-based interventions for individuals experiencing bereavement: a systematic review protocol. <i>BMJ Open</i> , 2020, 10, e036034.	1.9	2
28	Internet-Based Cognitive Behavior Therapy Only for the Young? A Secondary Analysis of a Randomized Controlled Trial of Depression Treatment. <i>Frontiers in Psychiatry</i> , 2020, 11, 735.	2.6	8
29	Attitudes and treatment practice of general practitioners towards patients with obesity in primary care. <i>BMC Family Practice</i> , 2020, 21, 169.	2.9	16
30	Frequent attenders in the German healthcare system: determinants of high utilization of primary care services. Results from the cross-sectional German health interview and examination survey for adults (DEGS). <i>BMC Family Practice</i> , 2020, 21, 10.	2.9	11
31	Internet Use in Old Age: Results of a German Population-Representative Survey. <i>Journal of Medical Internet Research</i> , 2020, 22, e15543.	4.3	50
32	Cognitive behavioural therapy for the treatment of late life depression: study protocol of a multicentre, randomized, observer-blinded, controlled trial (CBTlate). <i>BMC Psychiatry</i> , 2019, 19, 423.	2.6	11
33	A novel superior medication-based chronic disease score predicted all-cause mortality in independent geriatric cohorts. <i>Journal of Clinical Epidemiology</i> , 2019, 105, 112-124.	5.0	14
34	Five A – counseling in weight management of obese patients in primary care: a cluster-randomized controlled trial (INTERACT). <i>BMC Family Practice</i> , 2018, 19, 97.	2.9	13
35	Physician emigration from Germany: insights from a survey in Saxony, Germany. <i>BMC Health Services Research</i> , 2018, 18, 341.	2.2	24
36	Computerized cognitive behavior therapy for patients with mild to moderately severe depression in primary care: A pragmatic cluster randomized controlled trial (@ktiv). <i>Journal of Affective Disorders</i> , 2018, 238, 317-326.	4.1	53

#	ARTICLE	IF	CITATIONS
37	Which types of mental work demands may be associated with reduced risk of dementia?. <i>Alzheimer's and Dementia</i> , 2017, 13, 431-440.	0.8	33
38	Choosing the right rehabilitation setting after herniated disc surgery: Motives, motivations and expectations from the patientsâ€™ perspective. <i>PLoS ONE</i> , 2017, 12, e0183698.	2.5	1
39	Job satisfaction of foreign-national physicians working in patient care: a cross-sectional study in Saxony, Germany. <i>Journal of Occupational Medicine and Toxicology</i> , 2016, 11, 41.	2.2	31
40	Burnout among young physicians and its association with physiciansâ€™ wishes to leave: results of a survey in Saxony, Germany. <i>Journal of Occupational Medicine and Toxicology</i> , 2016, 11, 2.	2.2	73
41	The German version of the Camberwell Assessment of Need for the Elderly (CANE): evaluation of content validity and adaptation to the German-speaking context. <i>International Psychogeriatrics</i> , 2015, 27, 1919-1926.	1.0	19
42	Determinants of health-related quality of life in older primary care patients: results of the longitudinal observational AgeCoDe Study. <i>British Journal of General Practice</i> , 2015, 65, e716-e723.	1.4	59
43	Is the Short Form of the Mini-Mental State Examination (MMSE) a better screening instrument for dementia in older primary care patients than the original MMSE? Results of the German study on ageing, cognition, and dementia in primary care patients (AgeCoDe).. <i>Psychological Assessment</i> , 2015, 27, 895-904.	1.5	9
44	Association of the apolipoprotein E genotype with memory performance and executive functioning in cognitively intact elderly.. <i>Neuropsychology</i> , 2015, 29, 382-387.	1.3	26
45	Mild Depressive Symptoms Predict Mortality from Heart Failure in Elderly Men but Not Women. <i>Journal of the American Geriatrics Society</i> , 2015, 63, 2408-2410.	2.6	2
46	First evidence for glial pathology in late life minor depression: S100B is increased in males with minor depression. <i>Frontiers in Cellular Neuroscience</i> , 2015, 9, 406.	3.7	19
47	Risk of Dementia in Older Adults With Low Versus High Occupation-Based Motivational Processes. <i>Journal of Geriatric Psychiatry and Neurology</i> , 2015, 28, 126-135.	2.3	11
48	Cognitive performance before and after the onset of subjective cognitive decline in old age. <i>Alzheimer's and Dementia: Diagnosis, Assessment and Disease Monitoring</i> , 2015, 1, 194-205.	2.4	110
49	The association of BMI and social distance towards obese individuals is mediated by sympathy and understanding. <i>Social Science and Medicine</i> , 2015, 128, 25-30.	3.8	20
50	Age- and gender-specific norms for the German version of the Three-Factor Eating-Questionnaire (TFEQ). <i>Appetite</i> , 2015, 91, 241-247.	3.7	31
51	Cost-utility analysis of a preventive home visit program for older adults in Germany. <i>BMC Health Services Research</i> , 2015, 15, 141.	2.2	22
52	Differential effects of enriched environment at work on cognitive decline in old age. <i>Neurology</i> , 2015, 84, 2169-2176.	1.1	47
53	Inpatient or Outpatient Rehabilitation after Herniated Disc Surgery? â€” Setting-Specific Preferences, Participation and Outcome of Rehabilitation. <i>PLoS ONE</i> , 2014, 9, e89200.	2.5	5
54	Impact of Depression on Health Care Utilization and Costs among Multimorbid Patients â€” Results from the MultiCare Cohort Study. <i>PLoS ONE</i> , 2014, 9, e91973.	2.5	88

#	ARTICLE	IF	CITATIONS
55	Obesity and Associated Lifestyle in a Large Sample of Multi-Morbid German Primary Care Attendees. PLoS ONE, 2014, 9, e102587.	2.5	19
56	Associations between Dementia Outcomes and Depressive Symptoms, Leisure Activities, and Social Support. Dementia and Geriatric Cognitive Disorders Extra, 2014, 4, 481-493.	1.3	19
57	Social relations and depression in late life – A systematic review. International Journal of Geriatric Psychiatry, 2014, 29, 1-21.	2.7	203
58	Assessing met and unmet needs in the oldest-old and psychometric properties of the German version of the Camberwell Assessment of Need for the Elderly (CANE) – a pilot study. International Psychogeriatrics, 2014, 26, 285-295.	1.0	25
59	The role of spousal loss in the development of depressive symptoms in the elderly – Implications for diagnostic systems. Journal of Affective Disorders, 2014, 161, 97-103.	4.1	41
60	A comparison of GP and GDS diagnosis of depression in late life among multimorbid patients – Results of the MultiCare study. Journal of Affective Disorders, 2014, 168, 276-283.	4.1	12
61	Sexual Knowledge Among Adolescents with Physical Handicaps: A Systematic Review. Sexuality and Disability, 2014, 32, 429-441.	1.0	13
62	Association between mental demands at work and cognitive functioning in the general population – results of the health study of the Leipzig research center for civilization diseases (LIFE). Journal of Occupational Medicine and Toxicology, 2014, 9, 23.	2.2	34
63	The Costs of Dementia From the Societal Perspective: Is Care Provided in the Community Really Cheaper than Nursing Home Care?. Journal of the American Medical Directors Association, 2014, 15, 117-126.	2.5	35
64	Measuring Negative Attitudes towards Overweight and Obesity in the German Population – Psychometric Properties and Reference Values for the German Short Version of the Fat Phobia Scale (FPS). PLoS ONE, 2014, 9, e114641.	2.5	21
65	Attitudes Towards Bariatric Surgery in the General Public. Obesity Surgery, 2013, 23, 338-345.	2.1	30
66	Incidence and predictors of depression in non-demented primary care attenders aged 75 years and older: results from a 3-year follow-up study. Age and Ageing, 2013, 42, 173-180.	1.6	65
67	Direct costs associated with depressive symptoms in late life: a 4.5-year prospective study. International Psychogeriatrics, 2013, 25, 292-302.	1.0	33
68	Depression and Incident Dementia. An 8-Year Population-Based Prospective Study. PLoS ONE, 2013, 8, e59246.	2.5	28
69	Perception of overweight and obesity from different angles: A qualitative study. Scandinavian Journal of Public Health, 2012, 40, 271-277.	2.3	24
70	Economics of Medically Unexplained Symptoms: A Systematic Review of the Literature. Psychotherapy and Psychosomatics, 2012, 81, 265-275.	8.8	154
71	Predictors of Institutionalisation in Incident Dementia – Results of the German Study on Ageing, Cognition and Dementia in Primary Care Patients (AgeCoDe Study). Dementia and Geriatric Cognitive Disorders, 2012, 33, 282-288.	1.5	39
72	Health Service Utilization and Costs of Depressive Symptoms in Late Life - A Systematic Review. Current Pharmaceutical Design, 2012, 18, 5936-5957.	1.9	89

#	ARTICLE	IF	CITATIONS
73	Natural course of depressive symptoms in late life. An 8-year population-based prospective study. <i>Journal of Affective Disorders</i> , 2012, 142, 166-171.	4.1	48
74	Psychiatric comorbidity as predictor of costs in back pain patients undergoing disc surgery: a longitudinal observational study. <i>BMC Musculoskeletal Disorders</i> , 2012, 13, 165.	1.9	12
75	Incidence of late-life depression: A systematic review. <i>Journal of Affective Disorders</i> , 2012, 142, 172-179.	4.1	113
76	Obese Children, Adults and Senior Citizens in the Eyes of the General Public: Results of a Representative Study on Stigma and Causation of Obesity. <i>PLoS ONE</i> , 2012, 7, e46924.	2.5	80
77	Does GP training in depression care affect patient outcome? - A systematic review and meta-analysis. <i>BMC Health Services Research</i> , 2012, 12, 10.	2.2	63
78	Age- and gender-specific prevalence of depression in latest-life – Systematic review and meta-analysis. <i>Journal of Affective Disorders</i> , 2012, 136, 212-221.	4.1	613
79	Prevalence and risk factors of depressive symptoms in latest life – results of the Leipzig Longitudinal Study of the Aged (LEILA 75+). <i>International Journal of Geriatric Psychiatry</i> , 2012, 27, 286-295.	2.7	51
80	Age-related predictors of institutionalization: results of the German study on ageing, cognition and dementia in primary care patients (AgeCoDe). <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2012, 47, 263-270.	3.1	49
81	Effects of psychiatric comorbidity on costs in patients undergoing disc surgery: a cross-sectional study. <i>Spine Journal</i> , 2011, 11, 601-609.	1.3	12
82	The stigma of obesity in the general public and its implications for public health - a systematic review. <i>BMC Public Health</i> , 2011, 11, 661.	2.9	124
83	Predictors of nursing home admission of individuals without a dementia diagnosis before admission - results from the Leipzig Longitudinal Study of the Aged (LEILA 75+). <i>BMC Health Services Research</i> , 2010, 10, 186.	2.2	72
84	Prediction of institutionalization in the elderly. A systematic review. <i>Age and Ageing</i> , 2010, 39, 31-38.	1.6	690
85	Gender differences in predictors of nursing home placement in the elderly: a systematic review. <i>International Psychogeriatrics</i> , 2009, 21, 1015-1025.	1.0	51
86	Direct costs associated with mild cognitive impairment in primary care. <i>International Journal of Geriatric Psychiatry</i> , 2008, 23, 963-971.	2.7	33
87	Service Utilization and Health-Care Costs in the Advanced Elderly. <i>Value in Health</i> , 2008, 11, 611-620.	0.3	46
88	Direct costs associated with depression in old age in Germany. <i>Journal of Affective Disorders</i> , 2008, 105, 195-204.	4.1	82
89	Prevalence and risk factors for depression in non-demented primary care attenders aged 75 years and older. <i>Journal of Affective Disorders</i> , 2008, 111, 153-163.	4.1	156
90	Prediction of Institutionalisation in Dementia. <i>Dementia and Geriatric Cognitive Disorders</i> , 2008, 26, 65-78.	1.5	249

#	ARTICLE	IF	CITATIONS
91	Time until Institutionalization in Incident Dementia Cases &ndash; Results of the Leipzig Longitudinal Study of the Aged (LEILA 75+). <i>Neuroepidemiology</i> , 2008, 31, 100-108.	2.3	33
92	Healthcare costs associated with recognized and unrecognized depression in old age. <i>International Psychogeriatrics</i> , 2008, 20, 1219.	1.0	27
93	Cost-of-illness studies of depression. <i>Journal of Affective Disorders</i> , 2007, 98, 29-43.	4.1	296