Andrew Campbell

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/9062578/publications.pdf

Version: 2024-02-01

933447 1058476 1,157 14 10 14 citations g-index h-index papers 17 17 17 2568 docs citations times ranked citing authors all docs

#	Article	IF	Citations
1	The RAC1 Target NCKAP1 Plays a Crucial Role in the Progression of Braf;Pten-Driven Melanoma in Mice. Journal of Investigative Dermatology, 2021, 141, 628-637.e15.	0.7	8
2	CRISPR activation screen in mice identifies novel membrane proteins enhancing pulmonary metastatic colonisation. Communications Biology, 2021, 4, 395.	4.4	12
3	Oncogenic BRAF, unrestrained by TGFÎ ² -receptor signalling, drives right-sided colonic tumorigenesis. Nature Communications, 2021, 12, 3464.	12.8	33
4	The amino acid transporter SLC7A5 is required for efficient growth of KRAS-mutant colorectal cancer. Nature Genetics, 2021, 53, 16-26.	21.4	114
5	A RAC-GEF network critical for early intestinal tumourigenesis. Nature Communications, 2021, 12, 56.	12.8	11
6	Lef1 restricts ectopic crypt formation and tumor cell growth in intestinal adenomas. Science Advances, 2021, 7, eabj0512.	10.3	6
7	MCL1 Is Required for Maintenance of Intestinal Homeostasis and Prevention of Carcinogenesis in Mice. Gastroenterology, 2020, 159, 183-199.	1.3	22
8	Epithelial NOTCH Signaling Rewires the Tumor Microenvironment of Colorectal Cancer to Drive Poor-Prognosis Subtypes and Metastasis. Cancer Cell, 2019, 36, 319-336.e7.	16.8	278
9	Loss of BCL9/9l suppresses Wnt driven tumourigenesis in models that recapitulate human cancer. Nature Communications, 2019, 10, 723.	12.8	64
10	Systems level expression correlation of Ras GTPase regulators. Cell Communication and Signaling, 2018, 16, 46.	6.5	4
11	TGFÎ ² pathway limits dedifferentiation following WNT and MAPK pathway activation to suppress intestinal tumourigenesis. Cell Death and Differentiation, 2017, 24, 1681-1693.	11.2	48
12	Genome-wide in vivo screen identifies novel host regulators of metastatic colonization. Nature, 2017, 541, 233-236.	27.8	194
13	Loss of P53 Function Activates JAK2–STAT3 Signaling to Promote Pancreatic Tumor Growth, Stroma Modification, andÂGemcitabine Resistance in Mice and Is Associated WithÂPatient Survival. Gastroenterology, 2016, 151, 180-193.e12.	1.3	211
14	P-Rex1 is required for efficient melanoblast migration and melanoma metastasis. Nature Communications, 2011, 2, 555.	12.8	152