

Tsutomu Nakashima

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9062961/publications.pdf>

Version: 2024-02-01

110
papers

4,915
citations

81743

39
h-index

102304

66
g-index

110
all docs

110
docs citations

110
times ranked

1770
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 1 | Visualization of Endolymphatic Hydrops in Patients With Meniere's Disease. <i>Laryngoscope</i> , 2007, 117, 415-420. | 1.1 | 497 |
| 2 | Grading of endolymphatic hydrops using magnetic resonance imaging. <i>Acta Oto-Laryngologica</i> , 2009, 129, 5-8. | 0.3 | 300 |
| 3 | Disorders of cochlear blood flow. <i>Brain Research Reviews</i> , 2003, 43, 17-28. | 9.1 | 207 |
| 4 | Visualization of Endolymphatic Hydrops in Ménière's Disease with Single-dose Intravenous Gadolinium-based Contrast Media using Heavily T2-weighted 3D-FLAIR. <i>Magnetic Resonance in Medical Sciences</i> , 2010, 9, 237-242. | 1.1 | 153 |
| 5 | Separate visualization of endolymphatic space, perilymphatic space and bone by a single pulse sequence; 3D-inversion recovery imaging utilizing real reconstruction after intratympanic Gd-DTPA administration at 3 Tesla. <i>European Radiology</i> , 2008, 18, 920-924. | 2.3 | 133 |
| 6 | Visualization of endolymphatic hydrops with MR imaging in patients with Ménière's disease and related pathologies: current status of its methods and clinical significance. <i>Japanese Journal of Radiology</i> , 2014, 32, 191-204. | 1.0 | 127 |
| 7 | Three-Dimensional Fluid-Attenuated Inversion Recovery Magnetic Resonance Imaging Findings and Prognosis in Sudden Sensorineural Hearing Loss. <i>Laryngoscope</i> , 2008, 118, 1433-1437. | 1.1 | 124 |
| 8 | Endolymphatic hydrops and blood-labyrinth barrier in Ménière's disease. <i>Acta Oto-Laryngologica</i> , 2011, 131, 474-479. | 0.3 | 114 |
| 9 | Imaging of Ménière's Disease after Intravenous Administration of Single-dose Gadodiamide: Utility of Subtraction Images with Different Inversion Time. <i>Magnetic Resonance in Medical Sciences</i> , 2012, 11, 213-219. | 1.1 | 108 |
| 10 | Three-Dimensional Fluid-Attenuated Inversion Recovery Magnetic Resonance Imaging Findings in Patients with Sudden Sensorineural Hearing Loss. <i>Laryngoscope</i> , 2006, 116, 1451-1454. | 1.1 | 96 |
| 11 | Endolymphatic hydrops revealed by intravenous gadolinium injection in patients with Ménière's disease. <i>Acta Oto-Laryngologica</i> , 2010, 130, 338-343. | 0.3 | 95 |
| 12 | Individual Differences in the Permeability of the Round Window. <i>Otology and Neurotology</i> , 2009, 30, 645-648. | 0.7 | 94 |
| 13 | Imaging of Endolymphatic and Perilymphatic Fluid at 3T After Intratympanic Administration of Gadolinium-Diethylene-Triamine Pentaacetic Acid. <i>American Journal of Neuroradiology</i> , 2008, 29, 724-726. | 1.2 | 89 |
| 14 | Increased Sensitivity to Low Concentration Gadolinium Contrast by Optimized Heavily T2-weighted 3D-FLAIR to Visualize Endolymphatic Space. <i>Magnetic Resonance in Medical Sciences</i> , 2010, 9, 73-80. | 1.1 | 82 |
| 15 | Relationship between endolymphatic hydrops and vestibular-evoked myogenic potential. <i>Acta Oto-Laryngologica</i> , 2010, 130, 917-923. | 0.3 | 78 |
| 16 | Magnetic Resonance Imaging of the Inner Ear in Meniere's Disease. <i>Otolaryngologic Clinics of North America</i> , 2010, 43, 1059-1080. | 0.5 | 77 |
| 17 | Relationship between the Degree of Endolymphatic Hydrops and Electrocochleography. <i>Audiology and Neuro-Otology</i> , 2010, 15, 254-260. | 0.6 | 75 |
| 18 | Idiopathic sudden sensorineural hearing loss in Japan. <i>Acta Oto-Laryngologica</i> , 2014, 134, 1158-1163. | 0.3 | 71 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 19 | Endolymphatic Hydrops Revealed by Magnetic Resonance Imaging in Patients With Acute Low-Tone Sensorineural Hearing Loss. <i>Otology and Neurotology</i> , 2013, 34, 1241-1246. | 0.7 | 68 |
| 20 | Endolymphatic space size in patients with vestibular migraine and Ménière's disease. <i>Journal of Neurology</i> , 2014, 261, 2079-2084. | 1.8 | 65 |
| 21 | Comparison of Contrast Effect on the Cochlear Perilymph after Intratympanic and Intravenous Gadolinium Injection. <i>American Journal of Neuroradiology</i> , 2012, 33, 773-778. | 1.2 | 63 |
| 22 | Endolymphatic hydrops in superior canal dehiscence and large vestibular aqueduct syndromes. <i>Laryngoscope</i> , 2016, 126, 1446-1450. | 1.1 | 59 |
| 23 | Visualization of Endolymphatic Hydrops in Ménière's Disease after Single-dose Intravenous Gadolinium-based Contrast Medium: Timing of Optimal Enhancement. <i>Magnetic Resonance in Medical Sciences</i> , 2012, 11, 43-51. | 1.1 | 58 |
| 24 | Endolymphatic hydrops revealed by magnetic resonance imaging in patients with atypical Meniere's disease. <i>Acta Oto-Laryngologica</i> , 2013, 133, 123-129. | 0.3 | 58 |
| 25 | Imaging of Endolymphatic and Perilymphatic Fluid after Intravenous Administration of Single-dose Gadodiamide. <i>Magnetic Resonance in Medical Sciences</i> , 2012, 11, 145-150. | 1.1 | 56 |
| 26 | Imaging of the endolymphatic space in patients with Ménière's disease. <i>Auris Nasus Larynx</i> , 2018, 45, 33-38. | 0.5 | 55 |
| 27 | MR Imaging of the Cochlear Modiolus: Area Measurement in Healthy Subjects and in Patients with a Large Endolymphatic Duct and Sac. <i>Radiology</i> , 1999, 213, 819-823. | 3.6 | 53 |
| 28 | 3 Tesla magnetic resonance imaging obtained 4 hours after intravenous gadolinium injection in patients with sudden deafness. <i>Acta Oto-Laryngologica</i> , 2010, 130, 665-669. | 0.3 | 53 |
| 29 | Clinical significance of endolymphatic imaging after intratympanic gadolinium injection. <i>Acta Oto-Laryngologica</i> , 2009, 129, 9-14. | 0.3 | 51 |
| 30 | Imaging Endolymphatic Hydrops at 3 Tesla Using 3D-FLAIR with Intratympanic Gd-DTPA Administration. <i>Magnetic Resonance in Medical Sciences</i> , 2008, 7, 85-91. | 1.1 | 50 |
| 31 | Increased signal intensity of the cochlea on pre- and post-contrast enhanced 3D-FLAIR in patients with vestibular schwannoma. <i>Neuroradiology</i> , 2009, 51, 855-863. | 1.1 | 50 |
| 32 | Association Between Endolymphatic Hydrops as Revealed by Magnetic Resonance Imaging and Caloric Response. <i>Otology and Neurotology</i> , 2011, 32, 1480-1485. | 0.7 | 47 |
| 33 | Imaging of Ménière's Disease after Intravenous Administration of Single-dose Gadodiamide: Utility of Multiplication of MR Cisternography and HYDROPS Image. <i>Magnetic Resonance in Medical Sciences</i> , 2013, 12, 63-68. | 1.1 | 47 |
| 34 | Tympanometric Findings in Patients With Enlarged Vestibular Aqueducts. <i>Laryngoscope</i> , 2002, 112, 1642-1646. | 1.1 | 44 |
| 35 | Endolymphatic hydrops revealed by intravenous gadolinium injection in patients with Meniere's disease. <i>Acta Oto-Laryngologica</i> , 2010, 130, 1-6. | 0.3 | 44 |
| 36 | Inner Ear Hemorrhage in Systemic Lupus Erythematosus. <i>Laryngoscope</i> , 2006, 116, 826-828. | 1.1 | 43 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 37 | Endolymphatic hydrops in patients with vestibular schwannoma: visualization by non-contrast-enhanced 3D FLAIR. <i>Neuroradiology</i> , 2011, 53, 1009-1015. | 1.1 | 43 |
| 38 | MR imaging of the inner ear: comparison of a three-dimensional fast spin-echo sequence with use of a dedicated quadrature-surface coil with a gadolinium-enhanced spoiled gradient-recalled sequence.. <i>Radiology</i> , 1998, 208, 679-685. | 3.6 | 42 |
| 39 | Image evaluation of endolymphatic space in fluctuating hearing loss without vertigo. <i>European Archives of Oto-Rhino-Laryngology</i> , 2009, 266, 1871-1877. | 0.8 | 41 |
| 40 | Phenotypes associated with replacement of His by Arg in the Pendred syndrome gene. <i>European Journal of Endocrinology</i> , 2001, 145, 697-703. | 1.9 | 40 |
| 41 | Cutting edge of inner ear MRI. <i>Acta Oto-Laryngologica</i> , 2009, 129, 15-21. | 0.3 | 40 |
| 42 | A perspective from magnetic resonance imaging findings of the inner ear: Relationships among cerebrospinal, ocular and inner ear fluids. <i>Auris Nasus Larynx</i> , 2012, 39, 345-355. | 0.5 | 40 |
| 43 | Three-Dimensional Fluid-Attenuated Inversion Recovery Magnetic Resonance Imaging Investigation of Inner Ear Disturbances in Cases of Middle Ear Cholesteatoma With Labyrinthine Fistula. <i>Otology and Neurotology</i> , 2007, 28, 1029-1033. | 0.7 | 39 |
| 44 | Endolymphatic hydrops in patients with unilateral and bilateral Meniere's disease. <i>Acta Oto-Laryngologica</i> , 2017, 137, 23-28. | 0.3 | 37 |
| 45 | Three-dimensional (3D) visualization of endolymphatic hydrops after intratympanic injection of Gd-DTPA: Optimization of a 3D real inversion-recovery turbo spin-echo (TSE) sequence and application of a 32-channel head coil at 3T. <i>Journal of Magnetic Resonance Imaging</i> , 2010, 31, 210-214. | 1.9 | 36 |
| 46 | Endolymphatic space imaging in patients with delayed endolymphatic hydrops. <i>Acta Oto-Laryngologica</i> , 2009, 129, 1169-1174. | 0.3 | 33 |
| 47 | Imaging of Ménière's Disease by Subtraction of MR Cisternography from Positive Perilymph Image. <i>Magnetic Resonance in Medical Sciences</i> , 2012, 11, 303-309. | 1.1 | 33 |
| 48 | 3D-FLAIR magnetic resonance imaging in the evaluation of mumps deafness. <i>International Journal of Pediatric Otorhinolaryngology</i> , 2006, 70, 2115-2117. | 0.4 | 32 |
| 49 | Imaging analysis in cases with inflammation-induced sensorineural hearing loss. <i>Acta Oto-Laryngologica</i> , 2009, 129, 239-243. | 0.3 | 32 |
| 50 | Contrast enhancement of the inner ear in magnetic resonance images taken at 10 minutes or 4 hours after intravenous gadolinium injection. <i>Acta Oto-Laryngologica</i> , 2012, 132, 241-246. | 0.3 | 32 |
| 51 | Magnetic resonance imaging of the inner ear after both intratympanic and intravenous gadolinium injections. <i>Acta Oto-Laryngologica</i> , 2013, 133, 434-438. | 0.3 | 32 |
| 52 | Anatomical Details of the Brainstem and Cranial Nerves Visualized by High Resolution Readout-segmented Multi-shot Echo-planar Diffusion-weighted Images using Unidirectional MPG at 3T. <i>Magnetic Resonance in Medical Sciences</i> , 2011, 10, 269-275. | 1.1 | 31 |
| 53 | Peak Width in Multifrequency Tympanometry and Endolymphatic Hydrops Revealed by Magnetic Resonance Imaging. <i>Otology and Neurotology</i> , 2012, 33, 912-915. | 0.7 | 31 |
| 54 | MR Imaging of Ménière's Disease after Combined Intratympanic and Intravenous Injection of Gadolinium using HYDROPS2. <i>Magnetic Resonance in Medical Sciences</i> , 2014, 13, 133-137. | 1.1 | 31 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 55 | Contrast-enhanced MR imaging of the endolymphatic sac in patients with sudden hearing loss. <i>European Radiology</i> , 2002, 12, 1121-1126. | 2.3 | 30 |
| 56 | Contrast Enhancement of the Anterior Eye Segment and Subarachnoid Space: Detection in the Normal State by Heavily T2-weighted 3D FLAIR. <i>Magnetic Resonance in Medical Sciences</i> , 2011, 10, 193-199. | 1.1 | 29 |
| 57 | Gadolinium distribution in cochlear perilymph: differences between intratympanic and intravenous gadolinium injection. <i>Neuroradiology</i> , 2012, 54, 1161-1169. | 1.1 | 28 |
| 58 | Communication between cochlear perilymph and cerebrospinal fluid through the cochlear modiolus visualized after intratympanic administration of Gd-DTPA. <i>Radiation Medicine</i> , 2008, 26, 597-602. | 0.8 | 27 |
| 59 | Magnetic Resonance Imaging Evaluation of Endolymphatic Hydrops in Cases With Otosclerosis. <i>Otology and Neurotology</i> , 2015, 36, 1146-1150. | 0.7 | 27 |
| 60 | Changes in endolymphatic hydrops in patients with Ménière's disease treated conservatively for more than 1 year. <i>Acta Oto-Laryngologica</i> , 2015, 135, 866-870. | 0.3 | 25 |
| 61 | 3D-FLAIR MRI findings in a patient with Ramsay Hunt syndrome. <i>Acta Oto-Laryngologica</i> , 2007, 127, 547-549. | 0.3 | 24 |
| 62 | Detection of Presumed Hemorrhage in the Ampullar Endolymph of the Semicircular Canal: A Case Report. <i>Magnetic Resonance in Medical Sciences</i> , 2009, 8, 187-191. | 1.1 | 24 |
| 63 | 3D-FLAIR MRI in facial nerve paralysis with and without audio-vestibular disorder. <i>Acta Oto-Laryngologica</i> , 2010, 130, 632-636. | 0.3 | 23 |
| 64 | Sudden Sensorineural Hearing Loss Associated with Inner Ear Anomaly. <i>Otology and Neurotology</i> , 2005, 26, 241-246. | 0.7 | 22 |
| 65 | Effects of anterior inferior cerebellar artery occlusion on cochlear blood flow – a comparison between laser-Doppler and microsphere methods. <i>Hearing Research</i> , 2001, 162, 85-90. | 0.9 | 21 |
| 66 | Three-dimensional fluid-attenuated inversion recovery magnetic resonance imaging findings in a patient with cochlear otosclerosis. <i>Auris Nasus Larynx</i> , 2008, 35, 269-272. | 0.5 | 21 |
| 67 | Changes in endolymphatic hydrops in a patient with Meniere's disease observed using magnetic resonance imaging. <i>Auris Nasus Larynx</i> , 2010, 37, 220-222. | 0.5 | 21 |
| 68 | Accuracy of 3.0 Tesla magnetic resonance imaging in the diagnosis of intracochlear schwannoma. <i>Auris Nasus Larynx</i> , 2011, 38, 551-554. | 0.5 | 21 |
| 69 | Cochlear blood flow during occlusion and reperfusion of the anterior inferior cerebellar artery – effect of topical application of dexamethasone to the round window. <i>Acta Oto-Laryngologica</i> , 2009, 129, 127-131. | 0.3 | 20 |
| 70 | 18F-FDG-PET/CT predicts survival in hypopharyngeal squamous cell carcinoma. <i>Annals of Nuclear Medicine</i> , 2013, 27, 297-302. | 1.2 | 20 |
| 71 | Endolymphatic hydrops in patients with tinnitus as the major symptom. <i>European Archives of Oto-Rhino-Laryngology</i> , 2013, 270, 3043-3048. | 0.8 | 19 |
| 72 | Influence of dietary iodine deficiency on the thyroid gland in Slc26a4-null mutant mice. <i>Thyroid Research</i> , 2011, 4, 10. | 0.7 | 18 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 73 | Contrast enhancement of the inner ear after intravenous administration of a standard or double dose of gadolinium contrast agents. <i>Acta Oto-Laryngologica</i> , 2011, 131, 1025-1031. | 0.3 | 18 |
| 74 | Comparison of computed tomography and magnetic resonance imaging for evaluation of cholesteatoma with labyrinthine fistulae. <i>Laryngoscope</i> , 2012, 122, 1121-1125. | 1.1 | 18 |
| 75 | Contrast-enhanced MRI of the inner ear after intratympanic injection of meglumine gadopentetate or gadodiamide hydrate. <i>Acta Oto-Laryngologica</i> , 2011, 131, 130-135. | 0.3 | 17 |
| 76 | Visualization of Endolymphatic Hydrops after Intratympanic Injection of Gd-DTPA: Comparison of 2D and 3D Real Inversion Recovery Imaging. <i>Magnetic Resonance in Medical Sciences</i> , 2011, 10, 101-106. | 1.1 | 17 |
| 77 | Blood flow to the promontory in cochlear otosclerosis. <i>Clinical Otolaryngology</i> , 2006, 31, 110-115. | 0.0 | 16 |
| 78 | Simultaneous Three-dimensional Visualization of the Intra-parotid Facial Nerve and Parotid Duct using a Three-dimensional Reversed FISP Sequence with Diffusion Weighting. <i>Magnetic Resonance in Medical Sciences</i> , 2010, 9, 153-158. | 1.1 | 16 |
| 79 | Lateral Semicircular Canal and Vertigo in Patients With Large Vestibular Aqueduct Syndrome. <i>Otology and Neurotology</i> , 2006, 27, 788-792. | 0.7 | 15 |
| 80 | Audiological signs in pediatric cases with dehiscence of the bony labyrinth caused by a high jugular bulb. <i>International Journal of Pediatric Otorhinolaryngology</i> , 2012, 76, 447-451. | 0.4 | 15 |
| 81 | Numerical Assessment of Cholesteatoma by Signal Intensity on Non-EP-DWI and ADC Maps. <i>Otology and Neurotology</i> , 2014, 35, 1007-1010. | 0.7 | 15 |
| 82 | Magnetic resonance imaging evaluation of endolymphatic hydrops and post-operative findings in cases with otosclerosis. <i>Acta Oto-Laryngologica</i> , 2017, 137, 242-245. | 0.3 | 15 |
| 83 | Prompt Contrast Enhancement of Cerebrospinal Fluid Space in the Fundus of the Internal Auditory Canal: Observations in Patients with Meningeal Diseases on 3D-FLAIR Images at 3 Tesla. <i>Magnetic Resonance in Medical Sciences</i> , 2006, 5, 151-155. | 1.1 | 15 |
| 84 | MR Imaging of the Cochlear Modiolus after Intratympanic Administration of Gd-DTPA. <i>Magnetic Resonance in Medical Sciences</i> , 2010, 9, 23-29. | 1.1 | 14 |
| 85 | Prognostic value of 18F-fluorodeoxyglucose uptake before treatment for pharyngeal cancer. <i>Annals of Nuclear Medicine</i> , 2014, 28, 356-362. | 1.2 | 14 |
| 86 | Response of cochlear blood flow to prostaglandin E1 applied topically to the round window. <i>Acta Oto-Laryngologica</i> , 2006, 126, 232-236. | 0.3 | 12 |
| 87 | Cochlear modiolus and lateral semicircular canal in sudden deafness. <i>Acta Oto-Laryngologica</i> , 2007, 127, 1157-1161. | 0.3 | 12 |
| 88 | Signal Alteration of the Cochlear Perilymph on 3 Different Sequences after Intratympanic Gd-DTPA Administration at 3 Tesla: Comparison of 3D-FLAIR, 3D-T1-weighted Imaging, and 3D-CISS. <i>Magnetic Resonance in Medical Sciences</i> , 2010, 9, 65-71. | 1.1 | 12 |
| 89 | Imaging of a congenital perilymphatic fistula. <i>International Journal of Pediatric Otorhinolaryngology</i> , 2003, 67, 421-425. | 0.4 | 11 |
| 90 | Enlarged endolymphatic duct and sac syndrome: relationship between MR findings and genotype of mutation in pendred syndrome gene. <i>Magnetic Resonance Imaging</i> , 2004, 22, 25-30. | 1.0 | 11 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 91 | Estimation of Gadolinium-induced T1-shortening with Measurement of Simple Signal Intensity Ratio between the Cochlea and Brain Parenchyma on 3D-FLAIR: Correlation with T1 Measurement by TI Scout Sequence. <i>Magnetic Resonance in Medical Sciences</i> , 2010, 9, 17-22. | 1.1 | 11 |
| 92 | Laryngeal sensation and pharyngeal delay time after (chemo)radiotherapy. <i>European Archives of Oto-Rhino-Laryngology</i> , 2014, 271, 2299-2304. | 0.8 | 11 |
| 93 | Endolymphatic Hydrops of the Labyrinth Visualized on Noncontrast MR Imaging: A Case Report. <i>Magnetic Resonance in Medical Sciences</i> , 2009, 8, 43-46. | 1.1 | 10 |
| 94 | Progressive hearing loss following acquired cytomegalovirus infection in an immunocompromised child. <i>American Journal of Otolaryngology - Head and Neck Medicine and Surgery</i> , 2013, 34, 89-92. | 0.6 | 10 |
| 95 | Magnetic Resonance Imaging of Endolymphatic Sac in Acute Low-Tone Sensorineural Hearing Loss without Vertigo. <i>Orl</i> , 2003, 65, 254-260. | 0.6 | 8 |
| 96 | Evaluation of Vascular Activity in Otosclerosis by Laser Doppler Flowmetry. <i>Otology and Neurotology</i> , 2013, 34, 1559-1563. | 0.7 | 8 |
| 97 | Dietary habits and medical examination findings in Japanese adults middle-aged or older who live alone. <i>Nutrition</i> , 2021, 89, 111268. | 1.1 | 7 |
| 98 | Cochlear Blood Flow and Speech Perception Ability in Cochlear Implant Users. <i>Otology and Neurotology</i> , 2012, 33, 165-168. | 0.7 | 6 |
| 99 | Contrast enhancement of the cochlear aqueduct in MR imaging: its frequency and clinical significance. <i>Neuroradiology</i> , 2003, 45, 626-630. | 1.1 | 5 |
| 100 | Estimation of Perilymph Enhancement after Intratympanic Administration of Gd-DTPA by Fast T1-mapping with a Dual Flip Angle 3D Spoiled Gradient Echo Sequence. <i>Magnetic Resonance in Medical Sciences</i> , 2013, 12, 223-228. | 1.1 | 5 |
| 101 | Olfactory and gustatory dysfunction caused by SARS-CoV-2: Comparison with cases of infection with influenza and other viruses. <i>Infection Control and Hospital Epidemiology</i> , 2021, 42, 113-114. | 1.0 | 4 |
| 102 | Expression of midkine in the cochlea. <i>Hearing Research</i> , 2001, 160, 10-14. | 0.9 | 3 |
| 103 | Olfactory Function in Persons with Cerebral Palsy. <i>Journal of Policy and Practice in Intellectual Disabilities</i> , 2019, 16, 217-222. | 1.7 | 3 |
| 104 | Longitudinal associations between hearing aid usage and cognition in community-dwelling Japanese older adults with moderate hearing loss. <i>PLoS ONE</i> , 2021, 16, e0258520. | 1.1 | 3 |
| 105 | In Reference to <i>Visualization of Endolymphatic Hydrops in Patients With Meniere's Disease</i>. <i>Laryngoscope</i> , 2008, 118, 946-947. | 1.1 | 2 |
| 106 | Imaging findings in a case with cholesteatoma in complete aural atresia. <i>American Journal of Otolaryngology - Head and Neck Medicine and Surgery</i> , 2010, 31, 297-299. | 0.6 | 2 |
| 107 | Visualization of White Matter Tracts Using a Non-Diffusion Weighted Magnetic Resonance Imaging Method: Does Intravenous Gadolinium Injection Four Hours Prior to the Examination Affect the Visualization of White Matter Tracts?. <i>PLoS ONE</i> , 2014, 9, e91860. | 1.1 | 2 |
| 108 | Effect of an enlarged endolymphatic duct on bone conduction threshold. <i>Acta Oto-Laryngologica</i> , 2008, 128, 534-538. | 0.3 | 1 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 109 | Cerumen impaction shown by brain magnetic resonance imaging in patients with cognitive impairment. <i>Geriatrics and Gerontology International</i> , 2016, 16, 392-395. | 0.7 | 1 |
| 110 | Gustatory function in persons with cerebral palsy. <i>Journal of Oral Rehabilitation</i> , 2020, 47, 523-527. | 1.3 | 1 |