

Tsutomu Nakashima

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/9062961/tsutomu-nakashima-publications-by-year.pdf>

Version: 2024-04-24

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

110
papers

3,946
citations

35
h-index

58
g-index

110
ext. papers

4,466
ext. citations

3
avg, IF

4.99
L-index

#	Paper	IF	Citations
110	Longitudinal associations between hearing aid usage and cognition in community-dwelling Japanese older adults with moderate hearing loss. <i>PLoS ONE</i> , 2021 , 16, e0258520	3.7	0
109	Olfactory and gustatory dysfunction caused by SARS-CoV-2: Comparison with cases of infection with influenza and other viruses. <i>Infection Control and Hospital Epidemiology</i> , 2021 , 42, 113-114	2	1
108	Dietary habits and medical examination findings in Japanese adults middle-aged or older who live alone. <i>Nutrition</i> , 2021 , 89, 111268	4.8	0
107	Gustatory function in persons with cerebral palsy. <i>Journal of Oral Rehabilitation</i> , 2020 , 47, 523-527	3.4	1
106	Olfactory Function in Persons with Cerebral Palsy. <i>Journal of Policy and Practice in Intellectual Disabilities</i> , 2019 , 16, 217-222	1.8	2
105	Imaging of the endolymphatic space in patients with Ménière disease. <i>Auris Nasus Larynx</i> , 2018 , 45, 33-38	2.2	35
104	Magnetic resonance imaging evaluation of endolymphatic hydrops and post-operative findings in cases with otosclerosis. <i>Acta Oto-Laryngologica</i> , 2017 , 137, 242-245	1.6	9
103	Endolymphatic hydrops in patients with unilateral and bilateral Ménière disease. <i>Acta Oto-Laryngologica</i> , 2017 , 137, 23-28	1.6	24
102	Endolymphatic hydrops in superior canal dehiscence and large vestibular aqueduct syndromes. <i>Laryngoscope</i> , 2016 , 126, 1446-50	3.6	43
101	Cerumen impaction shown by brain magnetic resonance imaging in patients with cognitive impairment. <i>Geriatrics and Gerontology International</i> , 2016 , 16, 392-5	2.9	
100	Changes in endolymphatic hydrops in patients with Ménière disease treated conservatively for more than 1 year. <i>Acta Oto-Laryngologica</i> , 2015 , 135, 866-70	1.6	19
99	Magnetic Resonance Imaging Evaluation of Endolymphatic Hydrops in Cases With Otosclerosis. <i>Otology and Neurotology</i> , 2015 , 36, 1146-50	2.6	20
98	Visualization of endolymphatic hydrops with MR imaging in patients with Ménière disease and related pathologies: current status of its methods and clinical significance. <i>Japanese Journal of Radiology</i> , 2014 , 32, 191-204	2.9	102
97	Laryngeal sensation and pharyngeal delay time after (chemo)radiotherapy. <i>European Archives of Oto-Rhino-Laryngology</i> , 2014 , 271, 2299-304	3.5	10
96	Endolymphatic space size in patients with vestibular migraine and Ménière disease. <i>Journal of Neurology</i> , 2014 , 261, 2079-84	5.5	45
95	Prognostic value of (18)F-fluorodeoxyglucose uptake before treatment for pharyngeal cancer. <i>Annals of Nuclear Medicine</i> , 2014 , 28, 356-62	2.5	12
94	Visualization of white matter tracts using a non-diffusion weighted magnetic resonance imaging method: does intravenous gadolinium injection four hours prior to the examination affect the visualization of white matter tracts?. <i>PLoS ONE</i> , 2014 , 9, e91860	3.7	1

93	MR imaging of Ménière disease after combined intratympanic and intravenous injection of gadolinium using HYDROPS2. <i>Magnetic Resonance in Medical Sciences</i> , 2014 , 13, 133-7	2.9	27
92	Idiopathic sudden sensorineural hearing loss in Japan. <i>Acta Oto-Laryngologica</i> , 2014 , 134, 1158-63	1.6	46
91	Numerical assessment of cholesteatoma by signal intensity on non-EP-DWI and ADC maps. <i>Otology and Neurotology</i> , 2014 , 35, 1007-10	2.6	10
90	¹⁸ F-FDG-PET/CT predicts survival in hypopharyngeal squamous cell carcinoma. <i>Annals of Nuclear Medicine</i> , 2013 , 27, 297-302	2.5	17
89	Endolymphatic hydrops in patients with tinnitus as the major symptom. <i>European Archives of Oto-Rhino-Laryngology</i> , 2013 , 270, 3043-8	3.5	15
88	Endolymphatic hydrops revealed by magnetic resonance imaging in patients with atypical Ménière disease. <i>Acta Oto-Laryngologica</i> , 2013 , 133, 123-9	1.6	44
87	Progressive hearing loss following acquired cytomegalovirus infection in an immunocompromised child. <i>American Journal of Otolaryngology - Head and Neck Medicine and Surgery</i> , 2013 , 34, 89-92	2.8	7
86	Magnetic resonance imaging of the inner ear after both intratympanic and intravenous gadolinium injections. <i>Acta Oto-Laryngologica</i> , 2013 , 133, 434-8	1.6	28
85	Endolymphatic hydrops revealed by magnetic resonance imaging in patients with acute low-tone sensorineural hearing loss. <i>Otology and Neurotology</i> , 2013 , 34, 1241-6	2.6	51
84	Evaluation of vascular activity in otosclerosis by laser Doppler flowmetry: comparison with computed tomographic densitometry. <i>Otology and Neurotology</i> , 2013 , 34, 1559-63	2.6	6
83	Imaging of Ménière disease after intravenous administration of single-dose gadodiamide: utility of multiplication of MR cisternography and HYDROPS image. <i>Magnetic Resonance in Medical Sciences</i> , 2013 , 12, 63-8	2.9	33
82	Estimation of perilymph enhancement after intratympanic administration of Gd-DTPA by fast T ₂ mapping with a dual flip angle 3D spoiled gradient echo sequence. <i>Magnetic Resonance in Medical Sciences</i> , 2013 , 12, 223-8	2.9	5
81	A perspective from magnetic resonance imaging findings of the inner ear: Relationships among cerebrospinal, ocular and inner ear fluids. <i>Auris Nasus Larynx</i> , 2012 , 39, 345-55	2.2	32
80	Audiological signs in pediatric cases with dehiscence of the bony labyrinth caused by a high jugular bulb. <i>International Journal of Pediatric Otorhinolaryngology</i> , 2012 , 76, 447-51	1.7	14
79	Gadolinium distribution in cochlear perilymph: differences between intratympanic and intravenous gadolinium injection. <i>Neuroradiology</i> , 2012 , 54, 1161-9	3.2	26
78	Imaging of Ménière disease after intravenous administration of single-dose gadodiamide: utility of subtraction images with different inversion time. <i>Magnetic Resonance in Medical Sciences</i> , 2012 , 11, 213-9	2.9	85
77	Imaging of Ménière disease by subtraction of MR cisternography from positive perilymph image. <i>Magnetic Resonance in Medical Sciences</i> , 2012 , 11, 303-9	2.9	30
76	Comparison of computed tomography and magnetic resonance imaging for evaluation of cholesteatoma with labyrinthine fistulae. <i>Laryngoscope</i> , 2012 , 122, 1121-5	3.6	17

75	Contrast enhancement of the inner ear in magnetic resonance images taken at 10 minutes or 4 hours after intravenous gadolinium injection. <i>Acta Oto-Laryngologica</i> , 2012 , 132, 241-6	1.6	24
74	Comparison of contrast effect on the cochlear perilymph after intratympanic and intravenous gadolinium injection. <i>American Journal of Neuroradiology</i> , 2012 , 33, 773-8	4.4	49
73	Visualization of endolymphatic hydrops in Ménière disease after single-dose intravenous gadolinium-based contrast medium: timing of optimal enhancement. <i>Magnetic Resonance in Medical Sciences</i> , 2012 , 11, 43-51	2.9	53
72	Cochlear blood flow and speech perception ability in cochlear implant users. <i>Otology and Neurotology</i> , 2012 , 33, 165-8	2.6	5
71	Peak width in multifrequency tympanometry and endolymphatic hydrops revealed by magnetic resonance imaging. <i>Otology and Neurotology</i> , 2012 , 33, 912-5	2.6	24
70	Imaging of endolymphatic and perilymphatic fluid after intravenous administration of single-dose gadodiamide. <i>Magnetic Resonance in Medical Sciences</i> , 2012 , 11, 145-50	2.9	50
69	Contrast-enhanced MRI of the inner ear after intratympanic injection of meglumine gadopentetate or gadodiamide hydrate. <i>Acta Oto-Laryngologica</i> , 2011 , 131, 130-5	1.6	13
68	Accuracy of 3.0 Tesla magnetic resonance imaging in the diagnosis of intracochlear schwannoma. <i>Auris Nasus Larynx</i> , 2011 , 38, 551-4	2.2	20
67	Contrast enhancement of the anterior eye segment and subarachnoid space: detection in the normal state by heavily T2-weighted 3D FLAIR. <i>Magnetic Resonance in Medical Sciences</i> , 2011 , 10, 193-9	2.9	22
66	Visualization of endolymphatic hydrops after intratympanic injection of Gd-DTPA: comparison of 2D and 3D real inversion recovery imaging. <i>Magnetic Resonance in Medical Sciences</i> , 2011 , 10, 101-6	2.9	15
65	Anatomical details of the brainstem and cranial nerves visualized by high resolution readout-segmented multi-shot echo-planar diffusion-weighted images using unidirectional MPG at 3T. <i>Magnetic Resonance in Medical Sciences</i> , 2011 , 10, 269-75	2.9	24
64	Association between endolymphatic hydrops as revealed by magnetic resonance imaging and caloric response. <i>Otology and Neurotology</i> , 2011 , 32, 1480-5	2.6	35
63	Endolymphatic hydrops in patients with vestibular schwannoma: visualization by non-contrast-enhanced 3D FLAIR. <i>Neuroradiology</i> , 2011 , 53, 1009-15	3.2	33
62	Influence of dietary iodine deficiency on the thyroid gland in Slc26a4-null mutant mice. <i>Thyroid Research</i> , 2011 , 4, 10	2.4	16
61	Contrast enhancement of the inner ear after intravenous administration of a standard or double dose of gadolinium contrast agents. <i>Acta Oto-Laryngologica</i> , 2011 , 131, 1025-31	1.6	12
60	Endolymphatic hydrops and blood-labyrinth barrier in Ménière disease. <i>Acta Oto-Laryngologica</i> , 2011 , 131, 474-9	1.6	90
59	Estimation of gadolinium-induced T1-shortening with measurement of simple signal intensity ratio between the cochlea and brain parenchyma on 3D-FLAIR: correlation with T1 measurement by TI scout sequence. <i>Magnetic Resonance in Medical Sciences</i> , 2010 , 9, 17-22	2.9	11
58	Relationship between the degree of endolymphatic hydrops and electrocochleography. <i>Audiology and Neuro-Otology</i> , 2010 , 15, 254-60	2.2	61

57	Relationship between endolymphatic hydrops and vestibular-evoked myogenic potential. <i>Acta Oto-Laryngologica</i> , 2010 , 130, 917-23	1.6	64
56	3 Tesla magnetic resonance imaging obtained 4 hours after intravenous gadolinium injection in patients with sudden deafness. <i>Acta Oto-Laryngologica</i> , 2010 , 130, 665-9	1.6	45
55	Endolymphatic hydrops revealed by intravenous gadolinium injection in patients with Ménière's disease. <i>Acta Oto-Laryngologica</i> , 2010 , 130, 338-343	1.6	79
54	3D-FLAIR MRI in facial nerve paralysis with and without audio-vestibular disorder. <i>Acta Oto-Laryngologica</i> , 2010 , 130, 632-6	1.6	19
53	Magnetic resonance imaging of the inner ear in Meniere's disease. <i>Otolaryngologic Clinics of North America</i> , 2010 , 43, 1059-80	2	67
52	Imaging findings in a case with cholesteatoma in complete aural atresia. <i>American Journal of Otolaryngology - Head and Neck Medicine and Surgery</i> , 2010 , 31, 297-9	2.8	
51	Changes in endolymphatic hydrops in a patient with Meniere's disease observed using magnetic resonance imaging. <i>Auris Nasus Larynx</i> , 2010 , 37, 220-2	2.2	17
50	Visualization of endolymphatic hydrops in Ménière's disease with single-dose intravenous gadolinium-based contrast media using heavily T(2)-weighted 3D-FLAIR. <i>Magnetic Resonance in Medical Sciences</i> , 2010 , 9, 237-42	2.9	118
49	Increased sensitivity to low concentration gadolinium contrast by optimized heavily T2-weighted 3D-FLAIR to visualize endolymphatic space. <i>Magnetic Resonance in Medical Sciences</i> , 2010 , 9, 73-80	2.9	67
48	Simultaneous three-dimensional visualization of the intra-parotid facial nerve and parotid duct using a three-dimensional reversed FISP sequence with diffusion weighting. <i>Magnetic Resonance in Medical Sciences</i> , 2010 , 9, 153-8	2.9	14
47	MR imaging of the cochlear modiolus after intratympanic administration of Gd-DTPA. <i>Magnetic Resonance in Medical Sciences</i> , 2010 , 9, 23-9	2.9	10
46	Signal alteration of the cochlear perilymph on 3 different sequences after intratympanic Gd-DTPA administration at 3 tesla: comparison of 3D-FLAIR, 3D-T1-weighted imaging, and 3D-CISS. <i>Magnetic Resonance in Medical Sciences</i> , 2010 , 9, 65-71	2.9	10
45	Three-dimensional (3D) visualization of endolymphatic hydrops after intratympanic injection of Gd-DTPA: optimization of a 3D-real inversion-recovery turbo spin-echo (TSE) sequence and application of a 32-channel head coil at 3T. <i>Journal of Magnetic Resonance Imaging</i> , 2010 , 31, 210-4	5.6	26
44	Endolymphatic hydrops revealed by intravenous gadolinium injection in patients with Ménière's disease. <i>Acta Oto-Laryngologica</i> , 2010 , 130, 338-43	1.6	30
43	Detection of presumed hemorrhage in the ampullar endolymph of the semicircular canal: a case report. <i>Magnetic Resonance in Medical Sciences</i> , 2009 , 8, 187-91	2.9	20
42	Endolymphatic hydrops of the labyrinth visualized on noncontrast MR imaging: a case report. <i>Magnetic Resonance in Medical Sciences</i> , 2009 , 8, 43-6	2.9	9
41	Grading of endolymphatic hydrops using magnetic resonance imaging. <i>Acta Oto-Laryngologica</i> , 2009 , 5-8	1.6	229
40	Imaging analysis in cases with inflammation-induced sensorineural hearing loss. <i>Acta Oto-Laryngologica</i> , 2009 , 129, 239-43	1.6	28

39	Cochlear blood flow during occlusion and reperfusion of the anterior inferior cerebellar artery--effect of topical application of dexamethasone to the round window. <i>Acta Oto-Laryngologica</i> , 2009 , 129, 127-31	1.6	15
38	Endolymphatic space imaging in patients with delayed endolymphatic hydrops. <i>Acta Oto-Laryngologica</i> , 2009 , 129, 1169-74	1.6	27
37	Cutting edge of inner ear MRI. <i>Acta Oto-Laryngologica</i> , 2009 , 15-21	1.6	34
36	Image evaluation of endolymphatic space in fluctuating hearing loss without vertigo. <i>European Archives of Oto-Rhino-Laryngology</i> , 2009 , 266, 1871-7	3.5	36
35	Increased signal intensity of the cochlea on pre- and post-contrast enhanced 3D-FLAIR in patients with vestibular schwannoma. <i>Neuroradiology</i> , 2009 , 51, 855-63	3.2	39
34	Clinical significance of endolymphatic imaging after intratympanic gadolinium injection. <i>Acta Oto-Laryngologica</i> , 2009 , 9-14	1.6	45
33	Individual differences in the permeability of the round window: evaluating the movement of intratympanic gadolinium into the inner ear. <i>Otology and Neurotology</i> , 2009 , 30, 645-8	2.6	75
32	In Reference to Visualization of Endolymphatic Hydrops in Patients With Meniere's Disease. <i>Laryngoscope</i> , 2008 , 118, 946-947	3.6	1
31	Three-dimensional fluid-attenuated inversion recovery magnetic resonance imaging findings and prognosis in sudden sensorineural hearing loss. <i>Laryngoscope</i> , 2008 , 118, 1433-7	3.6	98
30	Three-dimensional fluid-attenuated inversion recovery magnetic resonance imaging findings in a patient with cochlear otosclerosis. <i>Auris Nasus Larynx</i> , 2008 , 35, 269-72	2.2	20
29	Effect of an enlarged endolymphatic duct on bone conduction threshold. <i>Acta Oto-Laryngologica</i> , 2008 , 128, 534-8	1.6	1
28	Imaging of endolymphatic and perilymphatic fluid at 3T after intratympanic administration of gadolinium-diethylene-triamine pentaacetic acid. <i>American Journal of Neuroradiology</i> , 2008 , 29, 724-6	4.4	76
27	Imaging endolymphatic hydrops at 3 tesla using 3D-FLAIR with intratympanic Gd-DTPA administration. <i>Magnetic Resonance in Medical Sciences</i> , 2008 , 7, 85-91	2.9	46
26	Communication between cochlear perilymph and cerebrospinal fluid through the cochlear modiolus visualized after intratympanic administration of Gd-DTPA. <i>Radiation Medicine</i> , 2008 , 26, 597-602		21
25	Separate visualization of endolymphatic space, perilymphatic space and bone by a single pulse sequence; 3D-inversion recovery imaging utilizing real reconstruction after intratympanic Gd-DTPA administration at 3 Tesla. <i>European Radiology</i> , 2008 , 18, 920-4	8	118
24	Visualization of endolymphatic hydrops in patients with Meniere's disease. <i>Laryngoscope</i> , 2007 , 117, 415-20	3.6	395
23	Cochlear modiolus and lateral semicircular canal in sudden deafness. <i>Acta Oto-Laryngologica</i> , 2007 , 127, 1157-61	1.6	8
22	3D-FLAIR MRI findings in a patient with Ramsay Hunt syndrome. <i>Acta Oto-Laryngologica</i> , 2007 , 127, 547-9.6		21

21	Three-dimensional fluid-attenuated inversion recovery magnetic resonance imaging investigation of inner ear disturbances in cases of middle ear cholesteatoma with labyrinthine fistula. <i>Otology and Neurotology</i> , 2007 , 28, 1029-33	2.6	37
20	Response of cochlear blood flow to prostaglandin E1 applied topically to the round window. <i>Acta Oto-Laryngologica</i> , 2006 , 126, 232-6	1.6	11
19	3D-FLAIR magnetic resonance imaging in the evaluation of mumps deafness. <i>International Journal of Pediatric Otorhinolaryngology</i> , 2006 , 70, 2115-7	1.7	29
18	Lateral semicircular canal and vertigo in patients with large vestibular aqueduct syndrome. <i>Otology and Neurotology</i> , 2006 , 27, 788-92	2.6	11
17	Blood flow to the promontory in cochlear otosclerosis. <i>Clinical Otolaryngology</i> , 2006 , 31, 110-5		13
16	Inner ear hemorrhage in systemic lupus erythematosus. <i>Laryngoscope</i> , 2006 , 116, 826-8	3.6	37
15	Three-dimensional fluid-attenuated inversion recovery magnetic resonance imaging findings in patients with sudden sensorineural hearing loss. <i>Laryngoscope</i> , 2006 , 116, 1451-4	3.6	88
14	Prompt contrast enhancement of cerebrospinal fluid space in the fundus of the internal auditory canal: observations in patients with meningeal diseases on 3D-FLAIR images at 3 Tesla. <i>Magnetic Resonance in Medical Sciences</i> , 2006 , 5, 151-5	2.9	13
13	Sudden sensorineural hearing loss associated with inner ear anomaly. <i>Otology and Neurotology</i> , 2005 , 26, 241-6	2.6	16
12	Enlarged endolymphatic duct and sac syndrome: relationship between MR findings and genotype of mutation in Pendred syndrome gene. <i>Magnetic Resonance Imaging</i> , 2004 , 22, 25-30	3.3	10
11	Magnetic resonance imaging of endolymphatic sac in acute low-tone sensorineural hearing loss without vertigo. <i>Orl</i> , 2003 , 65, 254-60	2	6
10	Contrast enhancement of the cochlear aqueduct in MR imaging: its frequency and clinical significance. <i>Neuroradiology</i> , 2003 , 45, 626-30	3.2	3
9	Disorders of cochlear blood flow. <i>Brain Research Reviews</i> , 2003 , 43, 17-28		163
8	Imaging of a congenital perilymphatic fistula. <i>International Journal of Pediatric Otorhinolaryngology</i> , 2003 , 67, 421-5	1.7	10
7	Tympanometric findings in patients with enlarged vestibular aqueducts. <i>Laryngoscope</i> , 2002 , 112, 1642-6	3.6	34
6	Contrast-enhanced MR imaging of the endolymphatic sac in patients with sudden hearing loss. <i>European Radiology</i> , 2002 , 12, 1121-6	8	25
5	Phenotypes associated with replacement of His by Arg in the Pendred syndrome gene. <i>European Journal of Endocrinology</i> , 2001 , 145, 697-703	6.5	34
4	Expression of midkine in the cochlea. <i>Hearing Research</i> , 2001 , 160, 10-4	3.9	3

3	Effects of anterior inferior cerebellar artery occlusion on cochlear blood flow--a comparison between laser-Doppler and microsphere methods. <i>Hearing Research</i> , 2001 , 162, 85-90	3.9	19
2	MR imaging of the cochlear modiolus: area measurement in healthy subjects and in patients with a large endolymphatic duct and sac. <i>Radiology</i> , 1999 , 213, 819-23	20.5	45
1	MR imaging of the inner ear: comparison of a three-dimensional fast spin-echo sequence with use of a dedicated quadrature-surface coil with a gadolinium-enhanced spoiled gradient-recalled sequence. <i>Radiology</i> , 1998 , 208, 679-85	20.5	37