## Kaoruko Iida

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/9065151/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Isonitrogenous low-carbohydrate diet elicits specific changes in metabolic gene expression in the skeletal muscle of exercise-trained mice. PLoS ONE, 2022, 17, e0262875.	1.1	1
2	Relationship between vitamin D receptor gene polymorphisms (Bsml, Taql, Apal, and Fokl) and calcium intake on bone mass in young Japanese women. BMC Women's Health, 2021, 21, 76.	0.8	3
3	Protective effect of Bacteroides fragilis LPS on Escherichia coli LPS-induced inflammatory changes in human monocytic cells and in a rheumatoid arthritis mouse model. Immunology Letters, 2021, 233, 48-56.	1.1	9
4	Association of Hours of Paid Work with Dietary Intake and Quality in Japanese Married Women: A Cross-Sectional Study. Nutrients, 2021, 13, 3005.	1.7	2
5	Association of alcohol consumption with prevalence of fatty liver after adjustment for dietary patterns: Cross-sectional analysis of Japanese middle-aged adults. Clinical Nutrition, 2020, 39, 1580-1586.	2.3	2
6	Gallic acid regulates adipocyte hypertrophy and suppresses inflammatory gene expression induced by the paracrine interaction between adipocytes and macrophages in vitro and in vivo. Nutrition Research, 2020, 73, 58-66.	1.3	23
7	Daidzein promotes the expression of oxidative phosphorylation- and fatty acid oxidation-related genes via an estrogen-related receptor α pathway to decrease lipid accumulation in muscle cells. Journal of Nutritional Biochemistry, 2020, 77, 108315.	1.9	19
8	Associations between nutritional adequacy and insomnia symptoms in Japanese men and women aged 18–69 years: a cross-sectional study. Sleep Health, 2020, 6, 197-204.	1.3	10
9	Associations Between Health Literacy and Underweight and Overweight Among Japanese Adults Aged 20 to 39 Years: A Cross-Sectional Study. Health Education and Behavior, 2020, 47, 631-639.	1.3	4
10	Effect of Cdx2 Polymorphism on the Relationship between Dietary Calcium Intake and Peak Bone Mass in Young Japanese Women. Nutrients, 2020, 12, 191.	1.7	2
11	Gallic Acid Inhibits Lipid Accumulation via AMPK Pathway and Suppresses Apoptosis and Macrophage-Mediated Inflammation in Hepatocytes. Nutrients, 2020, 12, 1479.	1.7	38
12	No association between fruits or vegetables and non-alcoholic fatty liver disease in middle-aged men and women. Nutrition, 2019, 61, 119-124.	1.1	34
13	Lactate Promotes Myoblast Differentiation and Myotube Hypertrophy via a Pathway Involving MyoD In Vitro and Enhances Muscle Regeneration In Vivo. International Journal of Molecular Sciences, 2018, 19, 3649.	1.8	42
14	<i>Terminalia bellirica</i> (Gaertn.) Roxb. Extract and Gallic Acid Attenuate LPS-Induced Inflammation and Oxidative Stress via MAPK/NF- <i>l°</i> B and Akt/AMPK/Nrf2 Pathways. Oxidative Medicine and Cellular Longevity, 2018, 2018, 1-15.	1.9	93
15	Relation of Bone Mass to Vitamin D Receptor Gene Polymorphism and Lifestyle Factors in Japanese Female College Students. Journal of Hard Tissue Biology, 2018, 27, 281-286.	0.2	1
16	Short-term and long-term ketogenic diet therapy and the addition of exercise have differential impacts on metabolic gene expression in the mouse energy-consuming organs heart and skeletal muscle. Nutrition Research, 2018, 60, 77-86.	1.3	20
17	Carbohydrate intake during early pregnancy is inversely associated with abnormal glucose challenge test results in Japanese pregnant women. Diabetes/Metabolism Research and Reviews, 2017, 33, e2898.	1.7	6
18	Association between rice, bread, and noodle intake and the prevalence of non-alcoholic fatty liver disease in Japanese middle-aged men and women. Clinical Nutrition, 2017, 36, 1601-1608.	2.3	33

Kaoruko lida

#	Article	IF	CITATIONS
19	Impact of Exercise and Nutrition on Bone Mass. Journal of Hard Tissue Biology, 2017, 26, 381-385.	0.2	1
20	Terminalia bellirica Extract Inhibits Low-Density Lipoprotein Oxidation and Macrophage Inflammatory Response in Vitro. Antioxidants, 2016, 5, 20.	2.2	19
21	The Dietary Isoflavone Daidzein Reduces Expression of Pro-Inflammatory Genes through PPARα/γ and JNK Pathways in Adipocyte and Macrophage Co-Cultures. PLoS ONE, 2016, 11, e0149676.	1.1	74
22	Milk-derived peptide Val-Pro-Pro (VPP) inhibits obesity-induced adipose inflammation via an angiotensin-converting enzyme (ACE) dependent cascade. Molecular Nutrition and Food Research, 2015, 59, 2502-2510.	1.5	43
23	Dietary isoflavone daidzein promotes Tfam expression that increases mitochondrial biogenesis in C2C12 muscle cells. Journal of Nutritional Biochemistry, 2015, 26, 1193-1199.	1.9	26
24	Daidzein regulates proinflammatory adipokines thereby improving obesityâ€related inflammation through PPARγ. Molecular Nutrition and Food Research, 2014, 58, 718-726.	1.5	54