

# Zhimin Chen

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9068793/publications.pdf>

Version: 2024-02-01

17  
papers

236  
citations

1162889

8  
h-index

1199470

12  
g-index

19  
all docs

19  
docs citations

19  
times ranked

415  
citing authors

#	ARTICLE	IF	CITATIONS
1	Does statins promote vascular calcification in chronic kidney disease?. European Journal of Clinical Investigation, 2017, 47, 137-148.	1.7	62
2	Vertebral bone density associates with coronary artery calcification and is an independent predictor of poor outcome in end-stage renal disease patients. Bone, 2016, 92, 50-57.	1.4	42
3	Bone mineral density and mortality in end-stage renal disease patients. CKJ: Clinical Kidney Journal, 2020, 13, 307-321.	1.4	36
4	Inverse J-shaped relation between coronary arterial calcium density and mortality in advanced chronic kidney disease. Nephrology Dialysis Transplantation, 2020, 35, 1202-1211.	0.4	20
5	High serum uric acid level is a mortality risk factor in peritoneal dialysis patients: a retrospective cohort study. Nutrition and Metabolism, 2019, 16, 52.	1.3	19
6	High alkaline phosphatase and low intact parathyroid hormone associate with worse clinical outcome in peritoneal dialysis patients. Peritoneal Dialysis International, 2021, 41, 236-243.	1.1	13
7	Bone mineral density of extremities is associated with coronary calcification and biopsy-verified vascular calcification in living-donor renal transplant recipients. Journal of Bone and Mineral Metabolism, 2017, 35, 536-543.	1.3	8
8	Differences in association of lower bone mineral density with higher coronary calcification in female and male end-stage renal disease patients. BMC Nephrology, 2019, 20, 59.	0.8	8
9	Effects of CD20+ B-cell infiltration into allografts on kidney transplantation outcomes: a systematic review and meta-analysis. Oncotarget, 2017, 8, 37935-37941.	0.8	8
10	The Role of an Integrated Care Model for Kidney Disease in the Development of Peritoneal Dialysis: A Single-Center Experience in China. Peritoneal Dialysis International, 2014, 34, 55-58.	1.1	7
11	Tacrolimus dose requirement based on the CYP3A5 genotype in renal transplant patients. Oncotarget, 2017, 8, 81285-81294.	0.8	7
12	Dialysate cell-free mitochondrial DNA fragments as a marker of intraperitoneal inflammation and peritoneal solute transport rate in peritoneal dialysis. BMC Nephrology, 2019, 20, 128.	0.8	6
13	SP356PULMONARY DYSFUNCTION IS LINKED TO MALNUTRITION AND INFLAMMATION IN CHRONIC KIDNEY DISEASE PATIENTS. Nephrology Dialysis Transplantation, 2016, 31, i208-i208.	0.4	0
14	SP323HIGH FIBRINOGEN LEVELS ARE INDEPENDENTLY ASSOCIATED WITH INCREASED MORTALITY IN PATIENTS WITH CHRONIC KIDNEY DISEASE (CKD). Nephrology Dialysis Transplantation, 2018, 33, i452-i452.	0.4	0
15	MO673: Association of Mean Corpuscular Volume with Mortality in Kidney Failure Patients: Opposite Findings in Chinese and Swedish Kidney Failure Patients. Nephrology Dialysis Transplantation, 2022, 37, .	0.4	0
16	MO708: Association of Atherogenic Index of Plasma (AIP) with Mortality in Chinese and Swedish End-Stage Kidney Disease Patients: Results from Two Cohorts. Nephrology Dialysis Transplantation, 2022, 37, .	0.4	0
17	MO681: Association of Framingham Risk Score with Mortality in Kidney Failure Patients from China and Sweden. Nephrology Dialysis Transplantation, 2022, 37, .	0.4	0