Petri Bono

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/9078422/publications.pdf

Version: 2024-02-01

257450 118850 4,257 68 24 62 citations h-index g-index papers 69 69 69 6702 all docs docs citations times ranked citing authors

| # | Article | IF | Citations |
|----|--|------|-----------|
| 1 | Nivolumab alone and nivolumab plus ipilimumab in recurrent small-cell lung cancer (CheckMate 032): a multicentre, open-label, phase $1/2$ trial. Lancet Oncology, The, 2016, 17, 883-895. | 10.7 | 1,091 |
| 2 | Nivolumab monotherapy in recurrent metastatic urothelial carcinoma (CheckMate 032): a multicentre, open-label, two-stage, multi-arm, phase 1/2 trial. Lancet Oncology, The, 2016, 17, 1590-1598. | 10.7 | 594 |
| 3 | Randomized, Controlled, Double-Blind, Cross-Over Trial Assessing Treatment Preference for Pazopanib Versus Sunitinib in Patients With Metastatic Renal Cell Carcinoma: PISCES Study. Journal of Clinical Oncology, 2014, 32, 1412-1418. | 1.6 | 381 |
| 4 | Initial efficacy of anti-lymphocyte activation gene-3 (anti–LAG-3; BMS-986016) in combination with nivolumab (nivo) in pts with melanoma (MEL) previously treated with anti–PD-1/PD-L1 therapy Journal of Clinical Oncology, 2017, 35, 9520-9520. | 1.6 | 188 |
| 5 | Nivolumab Alone and With Ipilimumab in Previously Treated Metastatic Urothelial Carcinoma: CheckMate 032 Nivolumab 1 mg/kg Plus Ipilimumab 3 mg/kg Expansion Cohort Results. Journal of Clinical Oncology, 2019, 37, 1608-1616. | 1.6 | 185 |
| 6 | Adjuvant Imatinib for High-Risk GI Stromal Tumor: Analysis of a Randomized Trial. Journal of Clinical Oncology, 2016, 34, 244-250. | 1.6 | 174 |
| 7 | Activity and safety of ODM-201 in patients with progressive metastatic castration-resistant prostate cancer (ARADES): an open-label phase 1 dose-escalation and randomised phase 2 dose expansion trial. Lancet Oncology, The, 2014, 15, 975-985. | 10.7 | 172 |
| 8 | High LYVE-1–Positive Lymphatic Vessel Numbers Are Associated with Poor Outcome in Breast Cancer. Clinical Cancer Research, 2004, 10, 7144-7149. | 7.0 | 156 |
| 9 | Adjuvant Capecitabine, Docetaxel, Cyclophosphamide, and Epirubicin for Early Breast Cancer: Final Analysis of the Randomized FinXX Trial. Journal of Clinical Oncology, 2012, 30, 11-18. | 1.6 | 114 |
| 10 | Afatinib alone or afatinib plus vinorelbine versus investigator's choice of treatment for HER2-positive breast cancer with progressive brain metastases after trastuzumab, lapatinib, or both (LUX-Breast 3): a randomised, open-label, multicentre, phase 2 trial. Lancet Oncology, The, 2015, 16, 1700-1710. | 10.7 | 108 |
| 11 | Second-line targeted therapies after nivolumab-ipilimumab failure in metastatic renal cell carcinoma. European Journal of Cancer, 2019, 108, 33-40. | 2.8 | 96 |
| 12 | Adjuvant Capecitabine in Combination With Docetaxel, Epirubicin, and Cyclophosphamide for Early Breast Cancer. JAMA Oncology, 2017, 3, 793. | 7.1 | 74 |
| 13 | Adjuvant capecitabine in combination with docetaxel and cyclophosphamide plus epirubicin for breast cancer: an open-label, randomised controlled trial. Lancet Oncology, The, 2009, 10, 1145-1151. | 10.7 | 65 |
| 14 | Hypertension as predictor of sunitinib treatment outcome in metastatic renal cell carcinoma. Acta Oncol \tilde{A}^3 gica, 2011, 50, 569-573. | 1.8 | 63 |
| 15 | First-in-human Phase 1 open label study of the BET inhibitor ODM-207 in patients with selected solid tumours. British Journal of Cancer, 2020, 123, 1730-1736. | 6.4 | 63 |
| 16 | Elevated serum endostatin is associated with poor outcome in patients with non-Hodgkin lymphoma. Cancer, 2003, 97, 2767-2775. | 4.1 | 52 |
| 17 | CheckMate-032: Phase I/II, open-label study of safety and activity of nivolumab (nivo) alone or with ipilimumab (ipi) in advanced and metastatic (A/M) gastric cancer (GC) Journal of Clinical Oncology, 2016, 34, 4010-4010. | 1.6 | 50 |
| 18 | Pharmacokinetics, Antitumor Activity, and Safety of ODM-201 in Patients with Chemotherapy-naive Metastatic Castration-resistant Prostate Cancer: An Open-label Phase 1 Study. European Urology, 2016, 69, 834-840. | 1.9 | 49 |

| # | Article | IF | Citations |
|----|--|-----|-----------|
| 19 | Sunitinibâ€induced hypertension, neutropaenia and thrombocytopaenia as predictors of good prognosis in patients with metastatic renal cell carcinoma. BJU International, 2016, 117, 110-117. | 2.5 | 47 |
| 20 | Trebananib (AMG 386) plus weekly paclitaxel with or without bevacizumab as first-line therapy for HER2-negative locally recurrent or metastatic breast cancer: A phase 2 randomized study. Breast, 2015, 24, 182-190. | 2.2 | 44 |
| 21 | Efficacy and safety of nivolumab monotherapy in metastatic urothelial cancer (mUC): Results from the phase I/II CheckMate 032 study Journal of Clinical Oncology, 2016, 34, 4501-4501. | 1.6 | 36 |
| 22 | Safety and activity of nivolumab monotherapy in advanced and metastatic (A/M) gastric or gastroesophageal junction cancer (GC/GEC): Results from the CheckMate-032 study Journal of Clinical Oncology, 2016, 34, 6-6. | 1.6 | 33 |
| 23 | Safety and Antitumour Activity of ODM-201 (BAY-1841788) in Chemotherapy-naÃ-ve and CYP17 Inhibitor-naÃ-ve Patients: Follow-up from the ARADES and ARAFOR Trials. European Urology Focus, 2018, 4, 547-553. | 3.1 | 30 |
| 24 | Systemic Blockade of Clever-1 Elicits Lymphocyte Activation Alongside Checkpoint Molecule Downregulation in Patients with Solid Tumors: Results from a Phase I/II Clinical Trial. Clinical Cancer Research, 2021, 27, 4205-4220. | 7.0 | 29 |
| 25 | Patient preference between pazopanib (Paz) and sunitinib (Sun): Results of a randomized double-blind, placebo-controlled, cross-over study in patients with metastatic renal cell carcinoma (mRCC)—PISCES study, NCT 01064310 Journal of Clinical Oncology, 2012, 30, CRA4502-CRA4502. | 1.6 | 27 |
| 26 | Spatial immunoprofiling of the intratumoral and peritumoral tissue of renal cell carcinoma patients. Modern Pathology, 2021, 34, 2229-2241. | 5.5 | 25 |
| 27 | Prospective Observational Study of Pazopanib in Patients with Advanced Renal Cell Carcinoma (PRINCIPAL Study). Oncologist, 2019, 24, 491-497. | 3.7 | 22 |
| 28 | CheckMate 025 phase III trial: Outcomes by key baseline factors and prior therapy for nivolumab (NIVO) versus everolimus (EVE) in advanced renal cell carcinoma (RCC) Journal of Clinical Oncology, 2016, 34, 498-498. | 1.6 | 21 |
| 29 | Association of Angiopoietin-2 and Ki-67 Expression with Vascular Density and Sunitinib Response in Metastatic Renal Cell Carcinoma. PLoS ONE, 2016, 11, e0153745. | 2.5 | 20 |
| 30 | Safety and Antitumour Activity of ODM-201 (BAY-1841788) in Castration-resistant, CYP17 Inhibitor-naÃ-ve Prostate Cancer: Results from Extended Follow-up of the ARADES Trial. European Urology Focus, 2017, 3, 606-614. | 3.1 | 18 |
| 31 | Patient preference between pazopanib (Paz) and sunitinib (Sun): Results of a randomized double-blind, placebo-controlled, cross-over study in patients with metastatic renal cell carcinoma (mRCC)—PISCES study, NCT 01064310 Journal of Clinical Oncology, 2012, 30, CRA4502-CRA4502. | 1.6 | 18 |
| 32 | Correlation of c-Met Expression and Outcome in Patients With Renal Cell Carcinoma Treated With Sunitinib. Clinical Genitourinary Cancer, 2017, 15, 487-494. | 1.9 | 16 |
| 33 | Angiotensin Inhibitors as Treatment of Sunitinib/Pazopanib-induced Hypertension inÂMetastatic Renal Cell Carcinoma. Clinical Genitourinary Cancer, 2017, 15, 384-390.e3. | 1.9 | 14 |
| 34 | Adjuvant Capecitabine for Early Breast Cancer: 15-Year Overall Survival Results From a Randomized Trial. Journal of Clinical Oncology, 2022, , JCO2102054. | 1.6 | 14 |
| 35 | VEGFR3 and CD31 as prognostic factors in renal cell cancer. Anticancer Research, 2015, 35, 921-7. | 1.1 | 13 |
| 36 | Outcome of surgery for patients with renal cell carcinoma and tumour thrombus in the era of modern targeted therapy. Scandinavian Journal of Urology, 2016, 50, 380-386. | 1.0 | 12 |

| # | Article | IF | CITATIONS |
|----|---|-----|-----------|
| 37 | Cancer costs and outcomes in the Finnish population 2004–2014. Acta Oncológica, 2018, 57, 297-303. | 1.8 | 10 |
| 38 | Tumor PIK3CA mutations, lymphocyte infiltration, and recurrence-free survival (RFS) in early breast cancer (BC): Results from the FinHER trial Journal of Clinical Oncology, 2012, 30, 507-507. | 1.6 | 10 |
| 39 | Phase I/II, open-label study of nivolumab (anti-PD-1; BMS-936558, ONO-4538) as monotherapy or combined with ipilimumab in advanced or metastatic solid tumors Journal of Clinical Oncology, 2014, 32, TPS3114-TPS3114. | 1.6 | 10 |
| 40 | Nivolumab monotherapy in metastatic urothelial carcinoma: Longer-term efficacy and safety results from the CheckMate 032 study Journal of Clinical Oncology, 2018, 36, 414-414. | 1.6 | 10 |
| 41 | Combined Angiogenesis and Proliferation Markers' Expressions as Long-Term Prognostic Factors in Renal Cell Cancer. Clinical Genitourinary Cancer, 2016, 14, e283-e289. | 1.9 | 9 |
| 42 | Real-world Effectiveness and Safety of Pazopanib in Patients With Intermediate Prognostic Risk Advanced Renal Cell Carcinoma. Clinical Genitourinary Cancer, 2019, 17, e526-e533. | 1.9 | 9 |
| 43 | Outcomes in Patients With Metastatic Renal Cell Carcinoma Who Develop Everolimus-Related Hyperglycemia and Hypercholesterolemia: Combined Subgroup Analyses of the RECORD-1 and REACT Trials. Clinical Genitourinary Cancer, 2016, 14, 406-414. | 1.9 | 8 |
| 44 | p95HER2 Methionine 611 Carboxy-Terminal Fragment Is Predictive of Trastuzumab Adjuvant Treatment Benefit in the FinHer Trial. Clinical Cancer Research, 2018, 24, 3046-3052. | 7.0 | 8 |
| 45 | Three vs. 1 year of adjuvant imatinib (IM) for operable high-risk GIST: The second planned analysis of the randomized SSGXVIII/AIO trial Journal of Clinical Oncology, 2015, 33, 10505-10505. | 1.6 | 8 |
| 46 | ODM-204, a Novel Dual Inhibitor of CYP17A1 and Androgen Receptor: Early Results from Phase I Dose Escalation in Men with Castration-resistant Prostate Cancer. European Urology Focus, 2020, 6, 63-70. | 3.1 | 7 |
| 47 | Decrease of Pro-Angiogenic Monocytes Predicts Clinical Response to Anti-Angiogenic Treatment in Patients with Metastatic Renal Cell Carcinoma. Cells, 2022, 11, 17. | 4.1 | 7 |
| 48 | Cancer costs and outcomes for common cancer sites in the Finnish population between 2009–2014. Acta Oncológica, 2018, 57, 983-988. | 1.8 | 6 |
| 49 | Modularising outpatient care delivery: A mixed methods case study at a Finnish University Hospital. Health Services Management Research, 2018, 31, 195-204. | 1.7 | 6 |
| 50 | Phase I/IIa, open-label, multicentre study to evaluate the optimal dosing and safety of ODM-203 in patients with advanced or metastatic solid tumours. ESMO Open, 2020, 5, e001081. | 4.5 | 6 |
| 51 | Novel Angiogenesis Markers as Long-Term Prognostic Factors in Patients With Renal Cell Cancer. Clinical Genitourinary Cancer, 2017, 15, e15-e24. | 1.9 | 5 |
| 52 | ODM-201 and the CNS: A clinical perspective Journal of Clinical Oncology, 2014, 32, 275-275. | 1.6 | 4 |
| 53 | A study of two ODM-201 formulations with a safety and tolerability extension phase in patients with metastatic chemotherapy-naive castration-resistant prostate cancer (CRPC) Journal of Clinical Oncology, 2014, 32, 115-115. | 1.6 | 3 |
| 54 | Adjuvant capesitabine in combination with docetaxel (T), epirubicin (E), and cyclophosphamide (C) in the treatment of early breast cancer (BC): 10-year survival results from the randomized FinXX trial Journal of Clinical Oncology, 2016, 34, 1001-1001. | 1.6 | 3 |

| # | Article | IF | CITATIONS |
|----|--|-----|-----------|
| 55 | Dose escalation study of ODM-203, a selective dual FGFR/VEGFR inhibitor, in patients with advanced solid tumours Journal of Clinical Oncology, 2016, 34, 2576-2576. | 1.6 | 3 |
| 56 | A single-institution experience with bevacizumab in the treatment of metastatic colorectal cancer and in conjunction with liver resection. OncoTargets and Therapy, 2014, 7, 1177. | 2.0 | 2 |
| 57 | Pharmacokinetics, activity, and safety of ODM-201 in chemotherapy-naÃ-ve patients with metastatic castration-resistant prostate cancer: An open-label phase I trial with long-term extension Journal of Clinical Oncology, 2015, 33, 230-230. | 1.6 | 2 |
| 58 | Promising clinical benefit rates in advanced cancers alongside potential biomarker correlation in a phase I/II trial investigating bexmarilimab, a novel macrophage-guided immunotherapy Journal of Clinical Oncology, 2022, 40, 2645-2645. | 1.6 | 2 |
| 59 | Helicobacter pylori and gastrointestinal symptoms in diagnostics and adjuvant chemotherapy of colorectal cancer. Oncology Letters, 2014, 7, 553-559. | 1.8 | 1 |
| 60 | Long-term efficacy and safety of androgen receptor inhibitor ODM-201 in ARADES phase I/II trial Journal of Clinical Oncology, 2014, 32, 5079-5079. | 1.6 | 1 |
| 61 | A phase III comparative study of nivolumab (anti-PD-1; BMS-936558; ONO-4538) versus everolimus in patients (pts) with advanced or metastatic renal cell carcinoma (mRCC) previously treated with antiangiogenic therapy Journal of Clinical Oncology, 2013, 31, TPS4592-TPS4592. | 1.6 | 1 |
| 62 | Reply to M. Isik et al. Journal of Clinical Oncology, 2010, 28, e335-e336. | 1.6 | 0 |
| 63 | Principal: A prospective observational study of real-world treatment patterns and treatment outcomes in patients with advanced or metastatic renal cell carcinoma (mRCC) receiving pazopanib Journal of Clinical Oncology, 2014, 32, TPS4600-TPS4600. | 1.6 | 0 |
| 64 | Correlation of endothelial angiopoietin-2 expression with tumor angiogenesis and response to sunitinib in metastatic renal cell carcinoma Journal of Clinical Oncology, 2015, 33, 461-461. | 1.6 | 0 |
| 65 | ODM-204 a novel dual inhibitor of CYP17A1 and androgen receptor: Early results from phase I dose escalation in men with castration-resistant prostate cancer Journal of Clinical Oncology, 2017, 35, 246-246. | 1.6 | 0 |
| 66 | Prospective, multinational, observational study of real-world treatment outcomes with pazopanib in patients with advanced or metastatic renal cell carcinoma (PRINCIPAL study) Journal of Clinical Oncology, 2018, 36, 4574-4574. | 1.6 | 0 |
| 67 | Comparison of clinical outcomes with first-line pazopanib in clinical trial eligible and non-clinical trial eligible patients with renal cell carcinoma Journal of Clinical Oncology, 2018, 36, 4561-4561. | 1.6 | 0 |

Modelling treatment benefit for bexmarilimab (an anti-Clever-1 antibody and a novel macrophage) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50

e14500-e14500.