## Tohru Yamakuni

## List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/9081145/tohru-yamakuni-publications-by-year.pdf

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

86	2,912	27	51
papers	citations	h-index	g-index
97 ext. papers	3,123 ext. citations	3.6 avg, IF	4.23 L-index

#	Paper	IF	Citations
86	Effect of methoxyflavones contained in Kaempferia parviflora on CRE-mediated transcription in PC12D cells. <i>Bioorganic and Medicinal Chemistry Letters</i> , <b>2020</b> , 30, 127606	2.9	1
85	Extracellular Esynuclein enters dopaminergic cells by modulating flotillin-1-assisted dopamine transporter endocytosis. <i>FASEB Journal</i> , <b>2019</b> , 33, 10240-10256	0.9	10
84	A novel mechanism of depression: role for connexins. <i>European Neuropsychopharmacology</i> , <b>2018</b> , 28, 483-498	1.2	14
83	Imidacloprid, a neonicotinoid insecticide, facilitates tyrosine hydroxylase transcription and phenylethanolamine N-methyltransferase mRNA expression to enhance catecholamine synthesis and its nicotine-evoked elevation in PC12D cells. <i>Toxicology</i> , <b>2018</b> , 394, 84-92	4.4	14
82	Royal jelly coordinately enhances hippocampal neuronal expression of somatostatin and neprilysin genes conferring neuronal protection against toxic soluble amyloid-lbligomers implicated in Alzheimer disease pathogenesis. <i>Journal of Functional Foods</i> , <b>2018</b> , 51, 28-38	5.1	3
81	Ginsenoside Rg1 alleviates corticosterone-induced dysfunction of gap junctions in astrocytes. <i>Journal of Ethnopharmacology</i> , <b>2017</b> , 208, 207-213	5	23
80	Fermented Citrus reticulata (ponkan) fruit squeezed draff that contains a large amount of 4'-demethylnobiletin prevents MK801-induced memory impairment. <i>Journal of Natural Medicines</i> , <b>2017</b> , 71, 617-631	3.3	5
79	Ginsenoside Rg1-induced antidepressant effects involve the protection of astrocyte gap junctions within the prefrontal cortex. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , <b>2017</b> , 75, 183-191	5.5	22
78	Oral sinensetin, but not nobiletin alone, prevents MK801-induced impairment of memory formation in mice, like nobiletin-rich chinpi, a kampo medicine. <i>Traditional &amp; Kampo Medicine</i> , <b>2017</b> , 4, 116-120	0.7	
77	Tyrosine hydroxylase gene expression is facilitated by alcohol followed by the degradation of the protein by ubiquitin proteasome system. <i>Neuroendocrinology Letters</i> , <b>2017</b> , 38, 43-49	0.3	2
76	Dopamine or biopterin deficiency potentiates phosphorylation at (40)Ser and ubiquitination of tyrosine hydroxylase to be degraded by the ubiquitin proteasome system. <i>Biochemical and Biophysical Research Communications</i> , <b>2015</b> , 465, 53-8	3.4	22
75	Four new myrsinol diterpenes from Euphorbia prolifera. <i>Journal of Natural Medicines</i> , <b>2013</b> , 67, 333-8	3.3	7
74	Potent activity of nobiletin-rich Citrus reticulata peel extract to facilitate cAMP/PKA/ERK/CREB signaling associated with learning and memory in cultured hippocampal neurons: identification of the substances responsible for the pharmacological action. <i>Journal of Neural Transmission</i> , <b>2013</b> ,	4.3	46
73	SLC10A4 is a protease-activated transporter that transports bile acids. <i>Journal of Biochemistry</i> , <b>2013</b> , 154, 93-101	3.1	9
72	Nobiletin-rich Citrus reticulata peels, a kampo medicine for Alzheimer's disease: a case series. <i>Geriatrics and Gerontology International</i> , <b>2013</b> , 13, 236-8	2.9	34
71	Nobiletin induces inhibitions of Ras activity and mitogen-activated protein kinase kinase/extracellular signal-regulated kinase signaling to suppress cell proliferation in C6 rat glioma cells. <i>Biological and Pharmaceutical Bulletin</i> , <b>2013</b> , 36, 540-7	2.3	19
70	New myrsinol diterpenes from Euphorbia prolifera and their inhibitory activities on LPS-induced NO production. <i>Bioorganic and Medicinal Chemistry Letters</i> , <b>2012</b> , 22, 3612-8	2.9	24

69	Repeated treatment with nicotine induces phosphorylation of NMDA receptor NR2B subunit in the brain regions involved in behavioral sensitization. <i>Neuroscience Letters</i> , <b>2012</b> , 524, 133-8	3.3	14
68	Three new iridoids from the roots of Valeriana jatamansi. <i>Journal of Natural Medicines</i> , <b>2012</b> , 66, 653-7	3.3	12
67	Isolation, structural elucidation, and neuroprotective effects of iridoids from Valeriana jatamansi. <i>Bioscience, Biotechnology and Biochemistry</i> , <b>2012</b> , 76, 1401-3	2.1	16
66	Honeybee royal jelly and nobiletin stimulate CRE-mediated transcription in ERK-independent and -dependent fashions, respectively, in PC12D cells. <i>Journal of Pharmacological Sciences</i> , <b>2011</b> , 116, 384-7	3.7	9
65	High-performance liquid chromatography with photodiode array detection for determination of nobiletin content in the brain and serum of mice administrated the natural compound. <i>Analytical and Bioanalytical Chemistry</i> , <b>2011</b> , 400, 3635-41	4.4	38
64	Neuroprotective Kaurane Diterpenes from Fritillaria ebeiensis. <i>Bioscience, Biotechnology and Biochemistry</i> , <b>2011</b> , 75, 1386-8	2.1	17
63	Dual effects of nobiletin, a citrus polymethoxy flavone, on catecholamine secretion in cultured bovine adrenal medullary cells. <i>Journal of Neurochemistry</i> , <b>2010</b> , 114, 1030-8	6	8
62	Two distinct mechanisms for actin capping protein regulationsteric and allosteric inhibition. <i>PLoS Biology</i> , <b>2010</b> , 8, e1000416	9.7	58
61	Nobiletin improves brain ischemia-induced learning and memory deficits through stimulation of CaMKII and CREB phosphorylation. <i>Brain Research</i> , <b>2009</b> , 1295, 218-29	3.7	101
60	4'-Demethylnobiletin, a bioactive metabolite of nobiletin enhancing PKA/ERK/CREB signaling, rescues learning impairment associated with NMDA receptor antagonism via stimulation of the ERK cascade. <i>Biochemistry</i> , <b>2009</b> , 48, 7713-21	3.2	57
59	Neurobehavioral effects of tetrabromobisphenol A, a brominated flame retardant, in mice. <i>Toxicology Letters</i> , <b>2009</b> , 189, 78-83	4.4	93
58	Nobiletin, a citrus flavonoid with neurotrophic action, augments protein kinase A-mediated phosphorylation of the AMPA receptor subunit, GluR1, and the postsynaptic receptor response to glutamate in murine hippocampus. <i>European Journal of Pharmacology</i> , <b>2008</b> , 578, 194-200	5.3	66
57	A novel diol-derivative of chalcone produced by bioconversion, 3-(2,3-dihydroxyphenyl)-1-phenylpropan-1-one, activates PKA/MEK/ERK signaling and antagonizes Abeta-inhibitor of the cascade in cultured rat CNS neurons. <i>European Journal of Pharmacology</i> ,	5.3	15
56	Nobiletin, a citrus flavonoid, improves memory impairment and Abeta pathology in a transgenic mouse model of Alzheimer's disease. <i>Journal of Pharmacology and Experimental Therapeutics</i> , <b>2008</b> , 326, 739-44	4.7	171
55	Beneficial effects of Ajuga decumbens on osteoporosis and arthritis. <i>Biological and Pharmaceutical Bulletin</i> , <b>2008</b> , 31, 1199-204	2.3	26
54	Nobiletin, a citrus flavonoid, reverses learning impairment associated with N-methyl-D-aspartate receptor antagonism by activation of extracellular signal-regulated kinase signaling. <i>Journal of Pharmacology and Experimental Therapeutics</i> , <b>2007</b> , 321, 784-90	4.7	53
53	Nobiletin, a citrus flavonoid that improves memory impairment, rescues bulbectomy-induced cholinergic neurodegeneration in mice. <i>Journal of Pharmacological Sciences</i> , <b>2007</b> , 105, 122-6	3.7	107
52	Clerodane diterpenoids and flavonoids with NGF-potentiating activity from the aerial parts of Baccharis gaudichaudiana. <i>Chemical and Pharmaceutical Bulletin</i> , <b>2007</b> , 55, 1532-4	1.9	17

51	Ca2+ channel activating action of maitotoxin in cultured brainstem neurons. <i>European Journal of Pharmacology</i> , <b>2006</b> , 536, 223-31	5.3	12
50	Three-membered ring sesquiterpenoids with NGF-potentiating activity from the roots of Valeriana fauriei. <i>Planta Medica</i> , <b>2006</b> , 72, 373-5	3.1	15
49	Proinsulin C-peptide stimulates a PKC/IkappaB/NF-kappaB signaling pathway to activate COX-2 gene transcription in Swiss 3T3 fibroblasts. <i>Journal of Biochemistry</i> , <b>2006</b> , 139, 1083-8	3.1	17
48	Bioactive ent-clerodane diterpenoids from the aerial parts of Baccharis gaudichaudiana. <i>Journal of Natural Products</i> , <b>2006</b> , 69, 274-6	4.9	22
47	Garcinone B reduces prostaglandin E2 release and NF-kappaB-mediated transcription in C6 rat glioma cells. <i>Neuroscience Letters</i> , <b>2006</b> , 394, 206-10	3.3	25
46	Nobiletin restoring beta-amyloid-impaired CREB phosphorylation rescues memory deterioration in Alzheimer's disease model rats. <i>Neuroscience Letters</i> , <b>2006</b> , 400, 230-4	3.3	106
45	Iridoids and sesquiterpenoids with NGF-potentiating activity from the rhizomes and roots of Valeriana fauriei. <i>Chemical and Pharmaceutical Bulletin</i> , <b>2006</b> , 54, 123-5	1.9	18
44	Metronomic scheduling of a cyclic hexapeptide Ra-VII for anti-angiogenesis, tumor vessel maturation and anti-tumor activity. <i>Cancer Science</i> , <b>2006</b> , 97, 665-74	6.9	10
43	Mechanism of neurotrophic action of nobiletin in PC12D cells. <i>Biochemistry</i> , <b>2005</b> , 44, 13683-91	3.2	93
42	Intracellular cAMP controls a physical association of V-1 with CapZ in cultured mammalian endocrine cells. <i>Biochemical and Biophysical Research Communications</i> , <b>2005</b> , 331, 181-6	3.4	10
41	Nobiletin and its related flavonoids with CRE-dependent transcription-stimulating and neuritegenic activities. <i>Biochemical and Biophysical Research Communications</i> , <b>2005</b> , 337, 1330-6	3.4	123
40	Ephedorae herba decreases lipopolysaccharide-induced cyclooxgenase-2 protein expression and NF-kappaB-dependent transcription in C6 rat glioma cells. <i>Journal of Pharmacological Sciences</i> , <b>2005</b> , 98, 327-30	3.7	18
39	IC101 induces apoptosis by Akt dephosphorylation via an inhibition of heat shock protein 90-ATP binding activity accompanied by preventing the interaction with Akt in L1210 cells. <i>Journal of Pharmacology and Experimental Therapeutics</i> , <b>2004</b> , 310, 1288-95	4.7	22
38	Evidence for the involvement of protein kinase C in acidic pH-induced contraction in spontaneously hypertensive rat aorta. <i>Pharmacology</i> , <b>2004</b> , 71, 10-6	2.3	8
37	gamma-Mangostin inhibits inhibitor-kappaB kinase activity and decreases lipopolysaccharide-induced cyclooxygenase-2 gene expression in C6 rat glioma cells. <i>Molecular Pharmacology</i> , <b>2004</b> , 66, 667-74	4.3	90
36	Acidosis-induced protein tyrosine phosphorylation depends on Ca2+ influx via voltage-dependent Ca2+ channels in SHR aorta. <i>European Journal of Pharmacology</i> , <b>2004</b> , 504, 105-11	5.3	1
35	Nardosinone enhances nerve growth factor-induced neurite outgrowth in a mitogen-activated protein kinase- and protein kinase C-dependent manner in PC12D cells. <i>Journal of Pharmacological Sciences</i> , <b>2003</b> , 93, 122-5	3.7	27
34	Nardosinone, the first enhancer of neurite outgrowth-promoting activity of staurosporine and dibutyryl cyclic AMP in PC12D cells. <i>Developmental Brain Research</i> , <b>2003</b> , 145, 177-83		25

33	Overexpression of V-1 prevents nitric oxide-induced cell death: involvement of enhanced tetrahydrobiopterin biosynthesis. <i>Journal of Neuroscience Research</i> , <b>2003</b> , 72, 716-25	4.4	
32	Down-regulation of an ankyrin repeat-containing protein, V-1, during skeletal muscle differentiation and its re-expression in the regenerative process of muscular dystrophy.  Neuromuscular Disorders, 2003, 13, 32-41	2.9	13
31	Identification of ATF-2 as a transcriptional regulator for the tyrosine hydroxylase gene. <i>Journal of Biological Chemistry</i> , <b>2002</b> , 277, 40768-74	5.4	28
30	Stimulated tyrosine phosphorylation of phosphatidylinositol 3-kinase causes acidic pH-induced contraction in spontaneously hypertensive rat aorta. <i>Journal of Pharmacology and Experimental Therapeutics</i> , <b>2002</b> , 303, 1255-64	4.7	9
29	Enhanced expression of GTP cyclohydrolase I in V-1-overexpressing PC12D cells. <i>Biochemical and Biophysical Research Communications</i> , <b>2002</b> , 293, 962-8	3.4	11
28	Expression of V-1, a novel catecholamine biosynthesis regulatory protein, is enhanced by hypertension in atrial myocytes of Dahl salt-sensitive rats. <i>Biochemical and Biophysical Research Communications</i> , <b>2002</b> , 298, 793-7	3.4	5
27	Glucocorticoid inhibits expression of V-1, a catecholamine biosynthesis regulatory protein, in cultured adrenal medullary cells. <i>FEBS Letters</i> , <b>2002</b> , 528, 166-70	3.8	5
26	V-1, a catecholamine biosynthesis regulatory protein, positively controls catecholamine secretion in PC12D cells. <i>FEBS Letters</i> , <b>2002</b> , 530, 94-8	3.8	9
25	Picrosides I and II, selective enhancers of the mitogen-activated protein kinase-dependent signaling pathway in the action of neuritogenic substances on PC12D cells. <i>Life Sciences</i> , <b>2002</b> , 71, 1821-35	6.8	35
24	Enhancement of Noradrenergic Phenotype Expression in Transgenic Mice Overexpressing V-1, A Cytoplasmic Ankyrin Repeat Protein. <i>Advances in Behavioral Biology</i> , <b>2002</b> , 53-56		
23	Two steroidal saponins from Camassia cusickii induce L1210 cell death through the apoptotic mechanism. <i>Canadian Journal of Physiology and Pharmacology</i> , <b>2001</b> , 79, 953-958	2.4	6
22	Potentiation of nerve growth factor-action by picrosides I and II, natural iridoids, in PC12D cells. <i>European Journal of Pharmacology</i> , <b>2000</b> , 406, 203-8	5.3	52
21	Molecular and functional characterization of a novel mouse transient receptor potential protein homologue TRP7. Ca(2+)-permeable cation channel that is constitutively activated and enhanced by stimulation of G protein-coupled receptor. <i>Journal of Biological Chemistry</i> , <b>1999</b> , 274, 27359-70	5.4	373
20	A new regulatory protein of catecholamine synthesizing-enzyme expression. <i>Advances in Pharmacology</i> , <b>1998</b> , 42, 30-2	5.7	
19	A novel protein containing Cdc10/SWI6 motifs regulates expression of mRNA encoding catecholamine biosynthesizing enzymes. <i>Journal of Biological Chemistry</i> , <b>1998</b> , 273, 27051-4	5.4	33
18	Murine central neurons express a novel member of the cdc10/SWI6 motif-containing protein superfamily. <i>Molecular Brain Research</i> , <b>1996</b> , 40, 203-13		11
17	Differential gene expression of fibroblast growth factor receptor isoforms in rat ovary. <i>Molecular and Cellular Endocrinology</i> , <b>1994</b> , 104, 75-80	4.4	22
16	Production and secretion of nerve growth factor by clonal striated muscle cell line, G8-1.  Neurochemistry International, 1992, 21, 251-8	4.4	4

15	Localization of gene expression of calbindin in the brain of adult rats. <i>Neuroscience Letters</i> , <b>1992</b> , 138, 211-5	3.3	27
14	Regulation of nerve growth factor and nerve growth factor receptor production by NMDA in C6 glioma cells. <i>Molecular Brain Research</i> , <b>1992</b> , 14, 35-42		21
13	A rat cerebellar protein containing the cdc10/SWI6 motif. FEBS Journal, 1992, 207, 615-20		31
12	Development and migration of Purkinje cells in the mouse cerebellar primordium. <i>Anatomy and Embryology</i> , <b>1991</b> , 184, 195-212		100
11	Production of nerve growth factor in rat skeletal muscle. <i>Neuroscience Letters</i> , <b>1991</b> , 132, 5-7	3.3	52
10	Transient appearance of immunoreactivity for Ca-binding protein (spot 35-calbindin) in small principal neurons in the superior cervical ganglion of pre-weanling rats. <i>Journal of the Autonomic Nervous System</i> , <b>1991</b> , 35, 25-31		8
9	Localization of spot 35-calbindin (rat cerebellar calbindin) in the anterior pituitary of the rat: developmental and sexual differences. <i>Archives of Histology and Cytology</i> , <b>1990</b> , 53, 585-91		11
8	Expression of immunoreactivity for Ca-binding protein, spot 35 in the interstitial cell of the rat pineal organ. <i>The Histochemical Journal</i> , <b>1990</b> , 22, 4-10		17
7	An immunohistochemical study of the ontogeny of the horizontal cell in the rat retina using an antiserum against spot 35 protein, a novel Purkinje cell-specific protein, as a marker. <i>The Anatomical Record</i> , <b>1988</b> , 222, 103-9		14
6	Expression of beta-nerve growth factor mRNA in rat glioma cells and astrocytes from rat brain. <i>FEBS Letters</i> , <b>1987</b> , 223, 117-21	3.8	55
5	An immunohistochemical study on the ontogeny of cells immunoreactive for spot 35 protein, a novel Purkinje cell-specific protein, in the rat cerebellum. <i>Developmental Brain Research</i> , <b>1986</b> , 394, 225-	-31	28
4	A purkinje cell-specific protein (spot 35 protein) showing wide distribution in the endocrine system of some mammals. An immunohistochemical study <i>Acta Histochemica Et Cytochemica</i> , <b>1986</b> , 19, 545-55	<del>1</del> .9	10
3	Involvement of spot 35 protein, a cerebellar protein, in modulation of Purkinje cell activity of the rat cerebellum. <i>European Journal of Pharmacology</i> , <b>1985</b> , 108, 309-13	5.3	12
2	IMMUNOHISTOCHEMICAL DEMONSTRATION OF A CEREBELLAR PROTEIN (SPOT 35 PROTEIN) IN SOME SENSORY CELLS OF GUINEA PIGS. <i>Biomedical Research</i> , <b>1985</b> , 6, 329-334	1.5	21
1	Isolation and immunohistochemical localization of a cerebellar protein. <i>Neuroscience Letters</i> , <b>1984</b> , 45, 235-40	3.3	88