

Yuya Watanabe

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9081803/publications.pdf>

Version: 2024-02-01

27
papers

953
citations

471509

17
h-index

552781

26
g-index

27
all docs

27
docs citations

27
times ranked

1180
citing authors

#	ARTICLE	IF	CITATIONS
1	Adherence to the food-based Japanese dietary guidelines and prevalence of poor oral health-related quality of life among older Japanese adults in the Kyoto-Kameoka study. <i>British Journal of Nutrition</i> , 2022, 128, 467-476.	2.3	3
2	Diet quality and physical or comprehensive frailty among older adults. <i>European Journal of Nutrition</i> , 2022, 61, 2451-2462.	3.9	11
3	Relationship between Standing Trunk Extension Angle and Medial Elbow Injuries in Young Baseball Pitchers. <i>International Journal of Environmental Research and Public Health</i> , 2022, 19, 3895.	2.6	0
4	Dose-Response Relationship Between Life-Space Mobility and Mortality in Older Japanese Adults: A Prospective Cohort Study. <i>Journal of the American Medical Directors Association</i> , 2022, 23, 1869.e7-1869.e18.	2.5	9
5	Validation of the Kihon Checklist and the frailty screening index for frailty defined by the phenotype model in older Japanese adults. <i>BMC Geriatrics</i> , 2022, 22, .	2.7	21
6	Association Between the Prevalence of Frailty and Doubly Labeled Water-Calibrated Energy Intake Among Community-Dwelling Older Adults. <i>Journals of Gerontology - Series A Biological Sciences and Medical Sciences</i> , 2021, 76, 876-884.	3.6	23
7	Motion Analysis Focusing on Rotational Movements of Professional Female Baseball Pitchers: Comparison with Male University Baseball Pitchers. <i>International Journal of Environmental Research and Public Health</i> , 2021, 18, 13342.	2.6	1
8	Association between Mixing Ability of Masticatory Functions Measured Using Color-Changing Chewing Gum and Frailty among Japanese Older Adults: The Kyoto-Kameoka Study. <i>International Journal of Environmental Research and Public Health</i> , 2020, 17, 4555.	2.6	16
9	Objectively Measured Daily Step Counts and Prevalence of Frailty in 3,616 Older Adults. <i>Journal of the American Geriatrics Society</i> , 2020, 68, 2310-2318.	2.6	36
10	A U-Shaped Relationship between the Prevalence of Frailty and Body Mass Index in Community-Dwelling Japanese Older Adults: The Kyoto-Kameoka Study. <i>Journal of Clinical Medicine</i> , 2020, 9, 1367.	2.4	57
11	Comprehensive geriatric intervention in community-dwelling older adults: a cluster-randomized controlled trial. <i>Journal of Cachexia, Sarcopenia and Muscle</i> , 2020, 11, 26-37.	7.3	24
12	Consumption of green tea but not coffee is associated with the oral health-related quality of life among an older Japanese population: Kyoto-Kameoka cross-sectional study. <i>European Journal of Clinical Nutrition</i> , 2019, 73, 577-584.	2.9	8
13	Estimation of Energy Intake by a Food Frequency Questionnaire: Calibration and Validation with the Doubly Labeled Water Method in Japanese Older People. <i>Nutrients</i> , 2019, 11, 1546.	4.1	22
14	Relationship Between Physical Fitness at the End of Preseason and the Inseason Game Performance in Japanese Female Professional Baseball Players. <i>Journal of Strength and Conditioning Research</i> , 2019, 33, 1580-1588.	2.1	9
15	Association between echo intensity and attenuation of skeletal muscle in young and older adults: a comparison between ultrasonography and computed tomography. <i>Clinical Interventions in Aging</i> , 2018, Volume 13, 1871-1878.	2.9	39
16	Association between the Frequency of Protein-Rich Food Intakes and Kihon-Checklist Frailty Indices in Older Japanese Adults: The Kyoto-Kameoka Study. <i>Nutrients</i> , 2018, 10, 84.	4.1	17
17	Sociodemographic and physical predictors of non-participation in community based physical checkup among older neighbors: a case-control study from the Kyoto-Kameoka longitudinal study, Japan. <i>BMC Public Health</i> , 2018, 18, 568.	2.9	19
18	Chapter 3 Prevention of sarcopenia. <i>Geriatrics and Gerontology International</i> , 2018, 18, 23-27.	1.5	20

#	ARTICLE	IF	CITATIONS
19	Comprehensive geriatric intervention program with and without weekly class-style exercise: research protocol of a cluster randomized controlled trial in Kyoto-Kameoka Study. <i>Clinical Interventions in Aging</i> , 2018, Volume 13, 1019-1033.	2.9	13
20	Sex Difference in the Association Between Protein Intake and Frailty: Assessed Using the Kihon Checklist Indexes Among Older Adults. <i>Journal of the American Medical Directors Association</i> , 2018, 19, 801-805.	2.5	26
21	Prevalence of Frailty Assessed by Fried and Kihon Checklist Indexes in a Prospective Cohort Study: Design and Demographics of the Kyoto-Kameoka Longitudinal Study. <i>Journal of the American Medical Directors Association</i> , 2017, 18, 733.e7-733.e15.	2.5	68
22	Frequency of Fruit and Vegetable Consumption and the Oral Health-Related Quality of Life among Japanese Elderly: A Cross-Sectional Study from the Kyoto-Kameoka Study. <i>Nutrients</i> , 2017, 9, 1362.	4.1	23
23	Effect of resistance training using bodyweight in the elderly: Comparison of resistance exercise movement between slow and normal speed movement. <i>Geriatrics and Gerontology International</i> , 2015, 15, 1270-1277.	1.5	46
24	Effect of very low-intensity resistance training with slow movement on muscle size and strength in healthy older adults. <i>Clinical Physiology and Functional Imaging</i> , 2014, 34, 463-470.	1.2	82
25	Increased Muscle Size and Strength From Slow-Movement, Low-Intensity Resistance Exercise and Tonic Force Generation. <i>Journal of Aging and Physical Activity</i> , 2013, 21, 71-84.	1.0	59
26	Comparison of single- or multifrequency bioelectrical impedance analysis and spectroscopy for assessment of appendicular skeletal muscle in the elderly. <i>Journal of Applied Physiology</i> , 2013, 115, 812-818.	2.5	82
27	Echo intensity obtained from ultrasonography images reflecting muscle strength in elderly men. <i>Clinical Interventions in Aging</i> , 2013, 8, 993.	2.9	219