Elizabeth Tutton

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/9083381/publications.pdf

Version: 2024-02-01

567281 526287 38 777 15 27 citations h-index g-index papers 41 41 41 905 citing authors docs citations times ranked all docs

#	Article	IF	CITATIONS
1	Qualitative interview study exploring the patient experience of living with axial spondyloarthritis and fatigue: difficult, demanding and draining. BMJ Open, 2022, 12, e053958.	1.9	7
2	Flexibility and resistance exercises versus usual care for improving pain and function after distal radius fracture in adults aged 50 years or over: protocol for the WISE randomised multicentre feasibility trial. Pilot and Feasibility Studies, 2022, 8, 55.	1,2	2
3	Development of the Warwick Axial Spondyloarthritis faTigue and Energy questionnaire (WASTEd) – a new patient-reported outcome measure. Rheumatology Advances in Practice, 2022, 6, rkac027.	0.7	2
4	Patient and informal carer experience of hip fracture: a qualitative study using interviews and observation in acute orthopaedic trauma. BMJ Open, 2021, 11, e042040.	1.9	5
5	Reamed intramedullary nailing versus circular frame external fixation for segmental tibial fractures (STIFF-F): a mixed methods feasibility study. Pilot and Feasibility Studies, 2021, 7, 93.	1.2	1
6	A qualitative study of parents' and their child's experience of a medial epicondyle fracture. Bone & Joint Open, 2021, 2, 359-364.	2.6	3
7	Chronic pain with neuropathic characteristics after surgery for major trauma to the lower limb: prevalence, predictors, and association with pain severity, disability, and quality of life in the UK WHiST trial. Bone and Joint Journal, 2021, 103-B, 1047-1054.	4.4	16
8	Unattainable equipoise in randomized controlled trials. Bone & Joint Open, 2021, 2, 486-492.	2.6	2
9	Co-producing a multi-stakeholder Core Outcome Set for distal Tibia and Ankle fractures (COSTA): a study protocol. Trials, 2021, 22, 443.	1.6	3
10	P70â€∫Evaluating the WASTEd: a new co-produced patient-reported measure of axial spondyloarthritis (axSpA) fatigue and energy. Rheumatology, 2020, 59, .	1.9	0
11	P71â€fWorking collaboratively with patients as research partners in the co-production of the WASTEd: a measure of fatigue and energy in axial spondyloarthritis (axSpA). Rheumatology, 2020, 59, .	1.9	O
12	A mixed-methods systematic review of patients' experience of being invited to participate in surgical randomised controlled trials. Social Science and Medicine, 2020, 253, 112961.	3.8	18
13	Facilitating trial recruitment: A qualitative study of patient and staff experiences of an orthopaedic trauma trial. Trials, 2019, 20, 492.	1.6	15
14	A qualitative study of patients' experience of recovery after a distal femoral fracture. Injury, 2019, 50, 1750-1755.	1.7	9
15	I becomes we, but where is me? The unity–division paradox when caring for a relative with dementia: A qualitative study. International Journal of Older People Nursing, 2019, 14, e12263.	1.3	5
16	Intramedullary nails versus distal locking plates for fracture of the distal femur: results from the Trial of Acute Femoral Fracture Fixation (TrAFFix) randomised feasibility study and process evaluation. BMJ Open, 2019, 9, e026810.	1.9	16
17	Patient experience of long-term recovery after open fracture of the lower limb: a qualitative study using interviews in a community setting. BMJ Open, 2019, 9, e031261.	1.9	34
18	Progressive functional exercise versus best practice advice for adults aged 50 years or over after ankle fracture: protocol for a pilot randomised controlled trial in the UK - the Ankle Fracture Treatment: Enhancing Rehabilitation (AFTER) study. BMJ Open, 2019, 9, e030877.	1.9	8

#	Article	IF	CITATIONS
19	Understanding assessment on a hospital ward for older people: A qualitative study. Journal of Advanced Nursing, 2019, 75, 850-861.	3.3	4
20	Retrograde intramedullary nail fixation compared with fixed-angle plate fixation for fracture of the distal femur: the TrAFFix feasibility RCT. Health Technology Assessment, 2019, 23, 1-132.	2.8	8
21	Three-Year Follow-up of a Trial of Close Contact Casting vs Surgery for Initial Treatment of Unstable Ankle Fractures in Older Adults. JAMA - Journal of the American Medical Association, 2018, 319, 1274.	7.4	19
22	Understanding informal carers' experiences of caring for older people with a hip fracture: a systematic review of qualitative studies. Disability and Rehabilitation, 2018, 40, 740-750.	1.8	26
23	Participation in a trial in the emergency situation: a qualitative study of patient experience in the UK WOLLF trial. Trials, 2018, 19, 328.	1.6	10
24	Assessing fatigue in adults with axial spondyloarthritis: a systematic review of the quality and acceptability of patient-reported outcome measures. Rheumatology Advances in Practice, 2018, 2, rky017.	0.7	8
25	Effect of Negative Pressure Wound Therapy vs Standard Wound Management on 12-Month Disability Among Adults With Severe Open Fracture of the Lower Limb. JAMA - Journal of the American Medical Association, 2018, 319, 2280.	7.4	137
26	Negative-pressure wound therapy versus standard dressings for adults with an open lower limb fracture: the WOLLF RCT. Health Technology Assessment, 2018, 22, 1-162.	2.8	42
27	Enabling the flow of compassionate care: a grounded theory study. BMC Health Services Research, 2017, 17, 174.	2.2	32
28	Understanding patient and relative/carer experience of hip fracture in acute care: A qualitative study protocol. International Journal of Orthopaedic and Trauma Nursing, 2017, 25, 36-41.	0.9	3
29	Close Contact Casting vs Surgery for Initial Treatment of Unstable Ankle Fractures in Older Adults. JAMA - Journal of the American Medical Association, 2016, 316, 1455.	7.4	94
30	The Ankle Injury Management (AIM) trial: a pragmatic, multicentre, equivalence randomised controlled trial and economic evaluation comparing close contact casting with open surgical reduction and internal fixation in the treatment of unstable ankle fractures in patients aged over 60 years. Health Technology Assessment, 2016, 20, 1-158.	2.8	73
31	Protocol for a randomised controlled trial of standard wound management versus negative pressure wound therapy in the treatment of adult patients with an open fracture of the lower limb: UK Wound management of Open Lower Limb Fractures (UK WOLFF). BMJ Open, 2015, 5, e009087.	1.9	23
32	Patients' experiences of technology and care in adult intensive care. Journal of Advanced Nursing, 2015, 71, 2051-2061.	3.3	38
33	Ankle Injury Management (AIM): design of a pragmatic multi-centre equivalence randomised controlled trial comparing Close Contact Casting (CCC) to Open surgical Reduction and Internal Fixation (ORIF) in the treatment of unstable ankle fractures in patients over 60Âyears. BMC Musculoskeletal Disorders, 2014, 15, 79.	1.9	14
34	Staff and patient views of the concept of hope on a stroke unit: a qualitative study. Journal of Advanced Nursing, 2012, 68, 2061-2069.	3.3	20
35	Hope in orthopaedic trauma: A qualitative study. International Journal of Nursing Studies, 2012, 49, 872-879.	5.6	18
36	Fluid optimisation using a peripherally inserted central catheter (PICC) following proximal femoral fracture: Lessons learnt from a feasibility study. Journal of Orthopaedic Nursing, 2009, 13, 11-18.	0.2	4

ELIZABETH TUTTON

#	Article	IF	CITATION
37	An exploration of hope as a concept for nursing. Journal of Orthopaedic Nursing, 2009, 13, 119-127.	0.2	32
38	Professional nursing culture on a trauma unit: experiences of patients and staff. Journal of Advanced Nursing, 2008, 61, 145-153.	3.3	24