Giuseppe Gargiulo

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/9083810/publications.pdf

Version: 2024-02-01

393982 377514 35 1,459 19 34 citations g-index h-index papers 35 35 35 1565 docs citations times ranked citing authors all docs

#	Article	lF	CITATIONS
1	Neonicotinoid clothianidin adversely affects insect immunity and promotes replication of a viral pathogen in honey bees. Proceedings of the National Academy of Sciences of the United States of America, 2013, 110, 18466-18471.	3.3	531
2	Assembly of transcriptionally active chromatin in Xenopus oocytes requires specific DNA binding factors. Cell, 1984, 38, 511-521.	13.5	81
3	A simple procedure for parallel sequence analysis of both strands of 5′-labeled DNA. Gene, 1983, 23, 175-183.	1.0	76
4	Building up the <i>Drosophila</i> eggshell: First of all the eggshell genes must be transcribed. Developmental Dynamics, 2008, 237, 2061-2072.	0.8	75
5	Escherichia coli single-strand binding protein stabilizes specific denatured sites in superhelical DNA. Nature, 1983, 303, 770-774.	13.7	72
6	Chromatin fine structure of the histone gene complex of Drosophila melanogaster. Nucleic Acids Research, 1983, 11, 421-440.	6.5	65
7	Analogous cleavage of DNA by micrococcal nuclease and a 1,10-phenanthroline-cuprous complex. Nucleic Acids Research, 1982, 10, 5823-5834.	6.5	64
8	Analysis of the chromatin assembled in germinal vesicles of Xenopus oocytes. Journal of Molecular Biology, 1983, 170, 699-722.	2.0	60
9	EcR-B1 and Usp nuclear hormone receptors regulate expression of the VM32E eggshell gene during Drosophila oogenesis. Developmental Biology, 2009, 328, 541-551.	0.9	39
10	Cell Survival and Polarity of Drosophila Follicle Cells Require the Activity of Ecdysone Receptor B1 Isoform. Genetics, 2009, 181, 165-175.	1.2	36
11	Specific domains drive VM32E protein distribution and integration in <i>Drosophila</i> eggshell layers. Journal of Cell Science, 2001, 114, 2819-2829.	1.2	32
12	Drosophila vitelline membrane cross-linking requires the fs(1)Nasrat, fs(1)polehole and chorion genes activities. Development Genes and Evolution, 2001, 211, 573-580.	0.4	30
13	Extracellular NME proteins: a player or a bystander?. Laboratory Investigation, 2018, 98, 248-257.	1.7	29
14	dAkt kinase controls follicle cell size duringDrosophila oogenesis. Developmental Dynamics, 2005, 232, 845-854.	0.8	28
15	Notch signaling during development requires the function of awd, the Drosophila homolog of human metastasis suppressor gene Nm23. BMC Biology, 2014, 12, 12.	1.7	28
16	The impact on microtubule network of a bracovirus lîºB-like protein. Cellular and Molecular Life Sciences, 2010, 67, 1699-1712.	2.4	21
17	A polydnavirus-encoded ANK protein has a negative impact on steroidogenesis and development. Insect Biochemistry and Molecular Biology, 2018, 95, 26-32.	1.2	21
18	Drosophila VHL tumor-suppressor gene regulates epithelial morphogenesis by promoting microtubule and aPKC stability. Development (Cambridge), 2010, 137, 1493-1503.	1.2	20

#	Article	IF	Citations
19	Genetic, functional and evolutionary characterization of scox, the Drosophila melanogaster ortholog of the human SCO1 gene. Mitochondrion, 2010, 10, 433-448.	1.6	20
20	A Polydnavirus ANK Protein Acts as Virulence Factor by Disrupting the Function of Prothoracic Gland Steroidogenic Cells. PLoS ONE, 2014, 9, e95104.	1.1	19
21	Notch signaling links interactions between the C/EBP homolog slow border cells and the GILZ homolog bunched during cell migration. Developmental Biology, 2007, 305, 217-231.	0.9	17
22	Drosophila 4EHP is essential for the larval–pupal transition and required in the prothoracic gland for ecdysone biosynthesis. Developmental Biology, 2016, 410, 14-23.	0.9	16
23	Vasa protein is localized in the germ cells and in the oocyte-associated pyriform follicle cells during early oogenesis in the lizard Podarcis sicula. Development Genes and Evolution, 2009, 219, 361-367.	0.4	13
24	Dynamin controls extracellular level of Awd/Nme1 metastasis suppressor protein. Naunyn-Schmiedeberg's Archives of Pharmacology, 2016, 389, 1171-1182.	1.4	13
25	Spatial activation and repression of the Drosophila vitelline membrane gene VM32E are switched by a complex cis-regulatory system. Developmental Dynamics, 2000, 218, 499-506.	0.8	9
26	The ecdysone receptor signalling regulates microvilli formation in follicular epithelial cells. Cellular and Molecular Life Sciences, 2016, 73, 409-425.	2.4	7
27	Evidence for a novel function of Awd in maintenance of genomic stability. Scientific Reports, 2017, 7, 16820.	1.6	7
28	Expression of Drosophila melanogaster F elements in vivo. Molecular Genetics and Genomics, 1994, 245, 152-159.	2.4	6
29	Dpp signaling down-regulates the expression of VM32E eggshell gene during Drosophila oogenesis. Developmental Dynamics, 2006, 235, 768-775.	0.8	6
30	Egfr signaling modulates VM32E gene expression during Drosophila oogenesis. Development Genes and Evolution, 2007, 217, 529-540.	0.4	6
31	Changes in abo phenotypic expression without increase in rDNA in Drosophila melanogaster. Molecular Genetics and Genomics, 1986, 205, 366-371.	2.4	5
32	Vps28 Is Involved in the Intracellular Trafficking of Awd, the Drosophila Homolog of NME1/2. Frontiers in Physiology, 2019, 10, 983.	1.3	4
33	Comparative Expression Profiling of Wild Type Drosophila Malpighian Tubules and von Hippel-Lindau Haploinsufficient Mutant. Frontiers in Physiology, 2019, 10, 619.	1.3	2
34	The Impact of Drosophila Awd/NME1/2 Levels on Notch and Wg Signaling Pathways. International Journal of Molecular Sciences, 2020, 21, 7257.	1.8	1
35	<i>Drosophila VHL</i> tumor-suppressor gene regulates epithelial morphogenesis by promoting microtubule and aPKC stability. Journal of Cell Science, 2010, 123, e1-e1.	1.2	0

3