

Naoki Mori

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/9095435/naoki-mori-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

83
papers

2,077
citations

27
h-index

44
g-index

93
ext. papers

2,235
ext. citations

4.2
avg, IF

4.35
L-index

#	Paper	IF	Citations
83	Exportin-1 is critical for cell proliferation and survival in adult T cell leukemia.. <i>Investigational New Drugs</i> , 2022 , 1	4.3	0
82	FX1, a BCL6 inhibitor, reactivates BCL6 target genes and suppresses HTLV-1-infected T cells. <i>Investigational New Drugs</i> , 2021 , 1	4.3	0
81	Importin β regulates cell growth and survival during adult T cell leukemia/lymphoma therapy. <i>Investigational New Drugs</i> , 2021 , 39, 317-329	4.3	1
80	The anti-malaria agent artesunate exhibits cytotoxic effects in primary effusion lymphoma. <i>Investigational New Drugs</i> , 2021 , 39, 111-121	4.3	0
79	The antipsychotic drug pimozide is effective against human T-cell leukemia virus type 1-infected T cells. <i>European Journal of Pharmacology</i> , 2021 , 908, 174373	5.3	1
78	MALT-1 as a novel therapeutic target for adult T-cell leukemia. <i>European Journal of Haematology</i> , 2020 , 105, 460-467	3.8	5
77	Evaluation of artesunate for the treatment of adult T-cell leukemia/lymphoma. <i>European Journal of Pharmacology</i> , 2020 , 872, 172953	5.3	17
76	The role of CUDC-907, a dual phosphoinositide-3 kinase and histone deacetylase inhibitor, in inhibiting proliferation of adult T-cell leukemia. <i>European Journal of Haematology</i> , 2020 , 105, 763-772	3.8	2
75	Mitotic kinase PBK/TOPK as a therapeutic target for adult T-cell leukemia/lymphoma. <i>International Journal of Oncology</i> , 2018 , 53, 801-814	4.4	15
74	Anti-adult T-cell leukemia/lymphoma activity of cerdulatinib, a dual SYK/JAK kinase inhibitor. <i>International Journal of Oncology</i> , 2018 , 53, 1681-1690	4.4	10
73	Effects of NVP-BEZ235, a dual phosphatidylinositol 3-kinase/mammalian target of rapamycin inhibitor, on HTLV-1-infected T-cell lines. <i>Oncology Letters</i> , 2018 , 15, 5311-5317	2.6	6
72	Expression and significance of Pim-3 kinase in adult T-cell leukemia. <i>European Journal of Haematology</i> , 2017 , 99, 495-504	3.8	4
71	In vitro and in vivo anti-primary effusion lymphoma activities of fucoidan extracted from <i>Cladosiphon okamuranus</i> Tokida. <i>Oncology Reports</i> , 2017 , 38, 3197-3204	3.5	6
70	Butein inhibits NF- κ B, AP-1 and Akt activation in adult T-cell leukemia/lymphoma. <i>International Journal of Oncology</i> , 2017 , 51, 633-643	4.4	15
69	Peridinin, a carotenoid, inhibits proliferation and survival of HTLV-1-infected T-cell lines. <i>International Journal of Oncology</i> , 2016 , 49, 1713-21	4.4	12
68	Efficiency of AUY922 in mice with adult T-cell leukemia/lymphoma. <i>Oncology Letters</i> , 2016 , 12, 387-392	2.6	4
67	Activation of PKC- β in HTLV-1-infected T cells. <i>International Journal of Oncology</i> , 2015 , 46, 1609-18	4.4	5

66	Constitutive expression of IRF-5 in HTLV-1-infected T cells. <i>International Journal of Oncology</i> , 2015 , 47, 361-9	4.4	3
65	Induction of β - γ by Epstein-Barr virus latent membrane protein-1 and CD30. <i>International Journal of Oncology</i> , 2015 , 47, 2197-207	4.4	8
64	Abundant expression of HMGB1 in human T-cell lymphotropic virus type I-infected T-cell lines and high plasma levels of HMGB1 in patients with adult T-cell leukemia. <i>Oncology Letters</i> , 2014 , 7, 1239-1242	2.6	11
63	Human T cell leukemia virus type I tax-induced β - γ modulates tax-dependent and tax-independent gene expression in T cells. <i>Neoplasia</i> , 2013 , 15, 1110-24	6.4	9
62	CD69 overexpression by human T-cell leukemia virus type 1 Tax transactivation. <i>Biochimica Et Biophysica Acta - Molecular Cell Research</i> , 2013 , 1833, 1542-52	4.9	11
61	Cytotoxic effects of fucoidan nanoparticles against osteosarcoma. <i>Marine Drugs</i> , 2013 , 11, 4267-78	6	49
60	Protein phosphatase 2A as a potential target for treatment of adult T cell leukemia. <i>Current Cancer Drug Targets</i> , 2013 , 13, 829-42	2.8	3
59	Heat Shock Protein 90 Inhibitor NVP-AUY922 Has Potent Anti-Tumor Activity With Adult T-Cell Leukemia-Lymphoma Cells. <i>Blood</i> , 2013 , 122, 1829-1829	2.2	0
58	Honokiol induces cell cycle arrest and apoptosis via inhibition of survival signals in adult T-cell leukemia. <i>Biochimica Et Biophysica Acta - General Subjects</i> , 2012 , 1820, 879-87	4	30
57	Mechanisms of virus immune evasion lead to development from chronic inflammation to cancer formation associated with human papillomavirus infection. <i>Oncology Reviews</i> , 2012 , 6, e17	4.3	20
56	Pathogenesis of Metastatic Calcification and Acute Pancreatitis in Adult T-Cell Leukemia under Hypercalcemic State. <i>Leukemia Research and Treatment</i> , 2012 , 2012, 128617		6
55	Acquired hemophilia A in a patient with essential thrombocythemia. <i>Acta Haematologica</i> , 2012 , 127, 170-2	2.7	4
54	Beneficial effects of fucoidan in patients with chronic hepatitis C virus infection. <i>World Journal of Gastroenterology</i> , 2012 , 18, 2225-30	5.6	29
53	Fucoidan therapy decreases the proviral load in patients with human T-lymphotropic virus type-1-associated neurological disease. <i>Antiviral Therapy</i> , 2011 , 16, 89-98	1.6	38
52	Retraction: Curcumin targets Akt cell survival signaling pathway in HTLV-I-infected T-cell lines. <i>Cancer Science</i> , 2011 , 102, 498	6.9	2
51	Retraction: Anti-adult T-cell leukemia effects of a novel synthetic retinoid, Am80 (Tamibarotene). <i>Cancer Science</i> , 2011 , 102, 499	6.9	1
50	Retraction: Epstein-Barr virus-encoded latent membrane protein 1 activates β -catenin signaling in B lymphocytes. <i>Cancer Science</i> , 2011 , 102, 500	6.9	
49	Effects of AZD1152, a selective Aurora B kinase inhibitor, on Burkitt's and Hodgkin's lymphomas. <i>Biochemical Pharmacology</i> , 2011 , 81, 1106-15	6	17

48	Activation of AID by human T-cell leukemia virus Tax oncoprotein and the possible role of its constitutive expression in ATL genesis. <i>Carcinogenesis</i> , 2011 , 32, 110-9	4.6	28
47	Interleukin-1 alpha produced by human T-cell leukaemia virus type I-infected T cells induces intercellular adhesion molecule-1 expression on lung epithelial cells. <i>Journal of Medical Microbiology</i> , 2011 , 60, 1750-1761	3.2	9
46	Induction of CD69 expression by cagPAI-positive Helicobacter pylori infection. <i>World Journal of Gastroenterology</i> , 2011 , 17, 3691-9	5.6	2
45	Cell signaling modifiers for molecular targeted therapy in ATLL. <i>Frontiers in Bioscience - Landmark</i> , 2009 , 14, 1479-89	2.8	11
44	Targeting Bcl-2 Family Proteins in Adult T-Cell Leukemia/Lymphoma: In Vitro and In Vivo Effects of a Novel Bcl-2 Family Inhibitor ABT-737.. <i>Blood</i> , 2009 , 114, 1685-1685	2.2	
43	Rearranged NF-kappa B2 gene in an adult T-cell leukemia cell line. <i>Cancer Science</i> , 2008 , 99, 792-8	6.9	9
42	Human T-Cell Leukemia Virus Type 1 Tax Activates Caveolin-1 Gene Expression. <i>Blood</i> , 2008 , 112, 2816-2816		
41	Activation of hypoxia-inducible factor 1 in human T-cell leukaemia virus type 1-infected cell lines and primary adult T-cell leukaemia cells. <i>Biochemical Journal</i> , 2007 , 406, 317-23	3.8	36
40	Curcumin (diferuloylmethane) inhibits constitutive active NF-kappaB, leading to suppression of cell growth of human T-cell leukemia virus type I-infected T-cell lines and primary adult T-cell leukemia cells. <i>International Journal of Cancer</i> , 2006 , 118, 765-72	7.5	69
39	NIK-333 inhibits growth of human T-cell leukemia virus type I-infected T-cell lines and adult T-cell leukemia cells in association with blockade of nuclear factor-kappaB signal pathway. <i>Molecular Cancer Therapeutics</i> , 2006 , 5, 704-12	6.1	14
38	Human T-cell leukemia virus type 1 tax dysregulates beta-catenin signaling. <i>Journal of Virology</i> , 2006 , 80, 10497-505	6.6	29
37	Efficient intervention of growth and infiltration of primary adult T-cell leukemia cells by an HIV protease inhibitor, ritonavir. <i>Blood</i> , 2006 , 107, 716-24	2.2	64
36	Curcumin suppresses constitutive activation of AP-1 by downregulation of JunD protein in HTLV-1-infected T-cell lines. <i>Leukemia Research</i> , 2006 , 30, 313-21	2.7	44
35	Transactivation of CCL20 Gene by CD30 in Hodgkin's Lymphoma.. <i>Blood</i> , 2006 , 108, 2258-2258	2.2	
34	Elevated expression of CD30 in adult T-cell leukemia cell lines: possible role in constitutive NF-kappaB activation. <i>Retrovirology</i> , 2005 , 2, 29	3.6	35
33	Fucoidan extracted from Cladosiphon okamuranus Tokida induces apoptosis of human T-cell leukemia virus type 1-infected T-cell lines and primary adult T-cell leukemia cells. <i>Nutrition and Cancer</i> , 2005 , 52, 189-201	2.8	102
32	Transcriptional activation of survivin through the NF-kappaB pathway by human T-cell leukemia virus type I tax. <i>International Journal of Cancer</i> , 2005 , 115, 967-74	7.5	132
31	Dengue virus-induced apoptosis in hepatic cells is partly mediated by Apo2 ligand/tumour necrosis factor-related apoptosis-inducing ligand. <i>Journal of General Virology</i> , 2005 , 86, 1055-1065	4.9	45

30	Suppression of Cell Growth and Induction of Apoptosis of HTLV-I-Infected T-Cell Lines and Primary ATL Cells by Galectin-9.. <i>Blood</i> , 2005 , 106, 4810-4810	2.2	
29	Apoptosis induced by the histone deacetylase inhibitor FR901228 in human T-cell leukemia virus type 1-infected T-cell lines and primary adult T-cell leukemia cells. <i>Journal of Virology</i> , 2004 , 78, 4582-90	6.6	55
28	Elevated expression of CCL5/RANTES in adult T-cell leukemia cells: possible transactivation of the CCL5 gene by human T-cell leukemia virus type I tax. <i>International Journal of Cancer</i> , 2004 , 111, 548-57	7.5	32
27	Tax-Independent Constitutive IB Kinase Activation in Adult T-Cell Leukemia Cells. <i>Neoplasia</i> , 2004 , 6, 266-278	6.4	66
26	Rapid tumor formation of human T-cell leukemia virus type 1-infected cell lines in novel NOD-SCID/gammac(null) mice: suppression by an inhibitor against NF-kappaB. <i>Journal of Virology</i> , 2003 , 77, 5286-94	6.6	94
25	Repression of Smad-dependent transforming growth factor-beta signaling by Epstein-Barr virus latent membrane protein 1 through nuclear factor-kappaB. <i>International Journal of Cancer</i> , 2003 , 105, 661-8	7.5	27
24	Helicobacter pylori induces matrix metalloproteinase-9 through activation of nuclear factor kappaB. <i>Gastroenterology</i> , 2003 , 124, 983-92	13.3	97
23	Helicobacter pylori induces RANTES through activation of NF-kappa B. <i>Infection and Immunity</i> , 2003 , 71, 3748-56	3.7	19
22	Activation of cyclin D1 and D2 promoters by human T-cell leukemia virus type I tax protein is associated with IL-2-independent growth of T cells. <i>International Journal of Cancer</i> , 2002 , 99, 378-85	7.5	43
21	Bay 11-7082 inhibits transcription factor NF-kappaB and induces apoptosis of HTLV-I-infected T-cell lines and primary adult T-cell leukemia cells. <i>Blood</i> , 2002 , 100, 1828-34	2.2	250
20	Human T-cell leukemia virus type I Tax transactivates the matrix metalloproteinase-9 gene: potential role in mediating adult T-cell leukemia invasiveness. <i>Blood</i> , 2002 , 99, 1341-9	2.2	55
19	Helicobacter pylori-mediated transcriptional regulation of the human beta-defensin 2 gene requires NF-kappaB. <i>Cellular Microbiology</i> , 2001 , 3, 115-23	3.9	78
18	Constitutive expression of the cyclooxygenase-2 gene in T-cell lines infected with human T cell leukemia virus type I. <i>International Journal of Cancer</i> , 2001 , 94, 813-9	7.5	17
17	Human T-cell leukemia virus type 1 tax protein activates transcription through AP-1 site by inducing DNA binding activity in T cells. <i>Virology</i> , 2001 , 279, 38-46	3.6	60
16	Human T-cell leukemia virus type I tax protein induces the expression of anti-apoptotic gene Bcl-xL in human T-cells through nuclear factor-kappaB and c-AMP responsive element binding protein pathways. <i>Virus Genes</i> , 2001 , 22, 279-87	2.3	71
15	Interleukin-2-dependent but not independent T-cell lines infected with human T-cell leukemia virus type 1 selectively express CD45RO, a marker for persistent infection in vivo. <i>Virus Genes</i> , 2001 , 23, 263-71	2.3	5
14	Lactacystin activates FLICE (caspase 8) protease and induces apoptosis in Fas-resistant adult T-cell leukemia cell lines. <i>European Journal of Haematology</i> , 2000 , 64, 315-22	3.8	15
13	Activation of oncogenic transcription factor AP-1 in T cells infected with human T cell leukemia virus type 1. <i>AIDS Research and Human Retroviruses</i> , 2000 , 16, 1603-6	1.6	29

12	Interleukin-15 stimulates the proliferation of peripheral blood leukemic cells from adult T-cell leukemia patients. <i>European Journal of Haematology</i> , 1998 , 61, 278-9	3.8	3
11	High levels of the DNA-binding activity of E2F in adult T-cell leukemia and human T-cell leukemia virus type I-infected cells: possible enhancement of DNA-binding of E2F by the human T-cell leukemia virus I transactivating protein, <i>Tax</i> . <i>European Journal of Haematology</i> , 1997 , 58, 114-20	3.8	7
10	Characterization and regulation of interleukin-4 receptor in adult T-cell leukemia cells. <i>European Journal of Haematology</i> , 1996 , 56, 241-7	3.8	4
9	Interleukin-4 inhibits the production of interleukin-1 by adult T-cell leukemia cells. <i>European Journal of Haematology</i> , 1995 , 55, 121-5	3.8	6
8	Interleukin-1 alpha as an autocrine growth factor for acute lymphoblastic leukaemia cells. <i>British Journal of Haematology</i> , 1994 , 86, 386-8	4.5	12
7	Interleukin-4 induces proliferation of adult T-cell leukemia cells. <i>European Journal of Haematology</i> , 1993 , 50, 133-40	3.8	9
6	Angiotropic lymphoma of paranasal sinuses with initial symptoms of oculomotor nerve palsy. <i>Internal Medicine</i> , 1993 , 32, 237-42	1.1	4
5	Enhancing effect of interleukin-2 on production of parathyroid hormone-related protein by adult T-cell leukemia cells. <i>Japanese Journal of Cancer Research</i> , 1993 , 84, 425-30		8
4	An HTLV-I carrier who developed CD4+ T-CLL expressed the IL-2 receptor beta chain alone without expressing the alpha chain. <i>British Journal of Haematology</i> , 1993 , 85, 185-7	4.5	1
3	Studies on the hepatotoxicity induced by bis (tributyltin) oxide. <i>Archives of Toxicology</i> , 1992 , 66, 182-7	5.8	19
2	Effects of dietary zinc deficiency on protein secretory functions of the mouse testis. <i>Archives of Histology and Cytology</i> , 1991 , 54, 401-10		8
1	Corneal edema induced by bis (tributyltin) oxide. <i>Archives of Toxicology</i> , 1991 , 65, 651-5	5.8	7