

L Del Peso

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9103087/publications.pdf>

Version: 2024-02-01

63
papers

6,813
citations

126858

33
h-index

128225

60
g-index

69
all docs

69
docs citations

69
times ranked

8933
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|---|------|-----------|
| 1 | Interleukin-3-Induced Phosphorylation of BAD Through the Protein Kinase Akt. <i>Science</i> , 1997, 278, 687-689. | 6.0 | 2,085 |
| 2 | Nod1, an Apaf-1-like Activator of Caspase-9 and Nuclear Factor- κ B. <i>Journal of Biological Chemistry</i> , 1999, 274, 14560-14567. | 1.6 | 639 |
| 3 | An Induced Proximity Model for NF- κ B Activation in the Nod1/RICK and RIP Signaling Pathways. <i>Journal of Biological Chemistry</i> , 2000, 275, 27823-27831. | 1.6 | 478 |
| 4 | Hypoxia Induces the Activation of the Phosphatidylinositol 3-Kinase/Akt Cell Survival Pathway in PC12 Cells. <i>Journal of Biological Chemistry</i> , 2001, 276, 22368-22374. | 1.6 | 217 |
| 5 | RICK, a Novel Protein Kinase Containing a Caspase Recruitment Domain, Interacts with CLARP and Regulates CD95-mediated Apoptosis. <i>Journal of Biological Chemistry</i> , 1998, 273, 12296-12300. | 1.6 | 215 |
| 6 | Hypoxia Promotes Glycogen Accumulation through Hypoxia Inducible Factor (HIF)-Mediated Induction of Glycogen Synthase 1. <i>PLoS ONE</i> , 2010, 5, e9644. | 1.1 | 209 |
| 7 | Identification of a functional hypoxia-responsive element that regulates the expression of the egl nine homologue 3 (egl3/phd3) gene. <i>Biochemical Journal</i> , 2005, 390, 189-197. | 1.7 | 194 |
| 8 | Genome-wide identification of hypoxia-inducible factor binding sites and target genes by a probabilistic model integrating transcription-profiling data and in silico binding site prediction. <i>Nucleic Acids Research</i> , 2010, 38, 2332-2345. | 6.5 | 179 |
| 9 | The von Hippel Lindau/Hypoxia-inducible Factor (HIF) Pathway Regulates the Transcription of the HIF-Proline Hydroxylase Genes in Response to Low Oxygen. <i>Journal of Biological Chemistry</i> , 2003, 278, 48690-48695. | 1.6 | 155 |
| 10 | Rho proteins induce metastatic properties in vivo. <i>Oncogene</i> , 1997, 15, 3047-3057. | 2.6 | 153 |
| 11 | Hypoxia Inducible Factor 1-Alpha (HIF-1 Alpha) Is Induced during Reperfusion after Renal Ischemia and Is Critical for Proximal Tubule Cell Survival. <i>PLoS ONE</i> , 2012, 7, e33258. | 1.1 | 133 |
| 12 | Regulation of the forkhead transcription factor FKHR, but not the PAX3-FKHR fusion protein, by the serine/threonine kinase Akt. <i>Oncogene</i> , 1999, 18, 7328-7333. | 2.6 | 125 |
| 13 | Lack of Evidence for the Involvement of the Phosphoinositide 3-Kinase/Akt Pathway in the Activation of Hypoxia-inducible Factors by Low Oxygen Tension. <i>Journal of Biological Chemistry</i> , 2002, 277, 13508-13517. | 1.6 | 103 |
| 14 | The Transcription Factor Encyclopedia. <i>Genome Biology</i> , 2012, 13, R24. | 13.9 | 103 |
| 15 | The SIN3A histone deacetylase complex is required for a complete transcriptional response to hypoxia. <i>Nucleic Acids Research</i> , 2018, 46, 120-133. | 6.5 | 96 |
| 16 | Caenorhabditis elegans EGL-1 Disrupts the Interaction of CED-9 with CED-4 and Promotes CED-3 Activation. <i>Journal of Biological Chemistry</i> , 1998, 273, 33495-33500. | 1.6 | 93 |
| 17 | Targeting tumour hypoxia to prevent cancer metastasis. From biology, biosensing and technology to drug development: the METOXIA consortium. <i>Journal of Enzyme Inhibition and Medicinal Chemistry</i> , 2015, 30, 689-721. | 2.5 | 93 |
| 18 | Rho-regulated signals induce apoptosis in vitro and in vivo by a p53-independent, but Bcl2 dependent pathway. <i>Oncogene</i> , 1998, 17, 1855-1869. | 2.6 | 92 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 19 | EFNA3 long noncoding RNAs induced by hypoxia promote metastatic dissemination. <i>Oncogene</i> , 2015, 34, 2609-2620. | 2.6 | 91 |
| 20 | Generation of phosphorylcholine as an essential event in the activation of Raf-1 and MAP-kinases in growth factors-induced mitogenic stimulation. <i>Journal of Cellular Biochemistry</i> , 1995, 57, 141-149. | 1.2 | 89 |
| 21 | Linking extracellular survival signals and the apoptotic machinery. <i>Current Opinion in Neurobiology</i> , 1998, 8, 613-618. | 2.0 | 83 |
| 22 | Activation of type D phospholipase by serum stimulation and ras-induced transformation in NIH3T3 cells. <i>Oncogene</i> , 1994, 9, 1387-95. | 2.6 | 74 |
| 23 | Vitamin D differentially regulates colon stem cells in patient-derived normal and tumor organoids. <i>FEBS Journal</i> , 2020, 287, 53-72. | 2.2 | 67 |
| 24 | miR-127 Protects Proximal Tubule Cells against Ischemia/Reperfusion: Identification of Kinesin Family Member 3B as miR-127 Target. <i>PLoS ONE</i> , 2012, 7, e44305. | 1.1 | 59 |
| 25 | The Ras family of GTPases in cancer cell invasion. <i>Cellular and Molecular Life Sciences</i> , 2000, 57, 65-76. | 2.4 | 56 |
| 26 | Disruption of the CED-9-Â-CED-4 Complex by EGL-1 Is a Critical Step for Programmed Cell Death in <i>Caenorhabditis elegans</i> . <i>Journal of Biological Chemistry</i> , 2000, 275, 27205-27211. | 1.6 | 56 |
| 27 | ERK5/BMK1 Is a Novel Target of the Tumor Suppressor VHL: Implication in Clear Cell Renal Carcinoma. <i>Neoplasia</i> , 2013, 15, 649-IN17. | 2.3 | 53 |
| 28 | Interaction between PARP-1 and HIF-2Î± in the hypoxic response. <i>Oncogene</i> , 2014, 33, 891-898. | 2.6 | 47 |
| 29 | Improving analysis of transcription factor binding sites within ChIP-Seq data based on topological motif enrichment. <i>BMC Genomics</i> , 2014, 15, 472. | 1.2 | 47 |
| 30 | Cooperativity of Stress-Responsive Transcription Factors in Core Hypoxia-Inducible Factor Binding Regions. <i>PLoS ONE</i> , 2012, 7, e45708. | 1.1 | 46 |
| 31 | ERK2, but Not ERK1, Mediates Acquired and "De novo" Resistance to Imatinib Mesylate: Implication for CML Therapy. <i>PLoS ONE</i> , 2009, 4, e6124. | 1.1 | 41 |
| 32 | TFEA.ChIP: a tool kit for transcription factor binding site enrichment analysis capitalizing on ChIP-seq datasets. <i>Bioinformatics</i> , 2019, 35, 5339-5340. | 1.8 | 41 |
| 33 | Hypoxia-inducible factors and cancer. <i>Clinical and Translational Oncology</i> , 2007, 9, 278-289. | 1.2 | 37 |
| 34 | Comparative Study of Organoids from Patient-Derived Normal and Tumor Colon and Rectal Tissue. <i>Cancers</i> , 2020, 12, 2302. | 1.7 | 37 |
| 35 | Identification of a region on hypoxia-inducible-factor prolyl 4-hydroxylases that determines their specificity for the oxygen degradation domains. <i>Biochemical Journal</i> , 2007, 408, 231-240. | 1.7 | 36 |
| 36 | Hypoxia and Chromatin: A Focus on Transcriptional Repression Mechanisms. <i>Biomedicines</i> , 2018, 6, 47. | 1.4 | 35 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 37 | Specific oncolytic effect of a new hypoxia-inducible factor-dependent replicative adenovirus on von Hippel-Lindau-defective renal cell carcinomas. <i>Cancer Research</i> , 2003, 63, 6877-84. | 0.4 | 33 |
| 38 | Analysis of HIF-prolyl hydroxylases binding to substrates. <i>Biochemical and Biophysical Research Communications</i> , 2006, 351, 313-320. | 1.0 | 32 |
| 39 | Regulatory and Functional Connection of Microphthalmia-Associated Transcription Factor and Anti-Metastatic Pigment Epithelium Derived Factor in Melanoma. <i>Neoplasia</i> , 2014, 16, 529-542. | 2.3 | 30 |
| 40 | Down-regulation of Hypoxia-inducible Factor-2 in PC12 Cells by Nerve Growth Factor Stimulation. <i>Journal of Biological Chemistry</i> , 2003, 278, 31895-31901. | 1.6 | 28 |
| 41 | Induction of apoptosis by rho in NIH 3T3 cells requires two complementary signals. Ceramides function as a progression factor for apoptosis. <i>Oncogene</i> , 1995, 11, 2657-65. | 2.6 | 28 |
| 42 | Hypoxia Negatively Regulates Antimetastatic PEDF in Melanoma Cells by a Hypoxia Inducible Factor-Independent, Autophagy Dependent Mechanism. <i>PLoS ONE</i> , 2012, 7, e32989. | 1.1 | 27 |
| 43 | Activation of phospholipase D by growth factors and oncogenes in murine fibroblasts follow alternative but cross-talking pathways. <i>Biochemical Journal</i> , 1997, 322, 519-528. | 1.7 | 26 |
| 44 | The human <i>PKP2</i> /plakophilin-2 gene is induced by Wnt/ β -catenin in normal and colon cancer-associated fibroblasts. <i>International Journal of Cancer</i> , 2018, 142, 792-804. | 2.3 | 26 |
| 45 | Classification of Airflow Limitation Based on <i>z</i> -Score Underestimates Mortality in Patients with Chronic Obstructive Pulmonary Disease. <i>American Journal of Respiratory and Critical Care Medicine</i> , 2017, 196, 298-305. | 2.5 | 24 |
| 46 | Vitamin D and Wnt3A have additive and partially overlapping modulatory effects on gene expression and phenotype in human colon fibroblasts. <i>Scientific Reports</i> , 2019, 9, 8085. | 1.6 | 23 |
| 47 | Ras protein is involved in the physiological regulation of phospholipase D by platelet derived growth factor. <i>Oncogene</i> , 2000, 19, 431-437. | 2.6 | 21 |
| 48 | Disruption of the CED-9/CED-4 Complex by EGL-1 is a Critical Step for Programmed Cell Death in <i>C. elegans</i> . <i>Journal of Biological Chemistry</i> , 2000, 275, 27205-11. | 1.6 | 21 |
| 49 | Intussusceptive Vascular Remodeling Precedes Pathological Neovascularization. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2019, 39, 1402-1418. | 1.1 | 20 |
| 50 | Activation of phospholipase D by ras proteins is independent of protein kinase C. , 1996, 61, 599-608. | | 17 |
| 51 | The use of an active learning approach to teach metabolism to students of nutrition and dietetics. <i>Biochemistry and Molecular Biology Education</i> , 2013, 41, 131-138. | 0.5 | 15 |
| 52 | Accumulation of hypoxia-inducible factor-1 α through a novel electrophilic, thiol antioxidant-sensitive mechanism. <i>Cellular Signalling</i> , 2007, 19, 2098-2105. | 1.7 | 14 |
| 53 | Metabolic labeling of RNA uncovers the contribution of transcription and decay rates on hypoxia-induced changes in RNA levels. <i>Rna</i> , 2020, 26, 1006-1022. | 1.6 | 13 |
| 54 | A role for insulator elements in the regulation of gene expression response to hypoxia. <i>Nucleic Acids Research</i> , 2012, 40, 1916-1927. | 6.5 | 11 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 55 | Identification of non-coding genetic variants in samples from hypoxemic respiratory disease patients that affect the transcriptional response to hypoxia. <i>Nucleic Acids Research</i> , 2016, 44, gkw811. | 6.5 | 8 |
| 56 | A yeast three-hybrid system that reconstitutes mammalian hypoxia inducible factor regulatory machinery. <i>BMC Cell Biology</i> , 2008, 9, 18. | 3.0 | 7 |
| 57 | lnc RNAs, hypoxia and metastasis. <i>Oncoscience</i> , 2015, 2, 795-796. | 0.9 | 6 |
| 58 | Hypoxia compensates cell cycle arrest with progenitor differentiation during angiogenesis. <i>FASEB Journal</i> , 2020, 34, 6654-6674. | 0.2 | 6 |
| 59 | Modulation of phospholipase D by Ras proteins mediated by its effectors Ral-GDS, PI3K and Raf-1. <i>International Journal of Oncology</i> , 2002, 21, 477. | 1.4 | 5 |
| 60 | Non-invasive monitoring of hypoxia-inducible factor activation by optical imaging during antiangiogenic treatment in a xenograft model of ovarian carcinoma. <i>International Journal of Oncology</i> , 2011, 39, 543-52. | 1.4 | 3 |
| 61 | Apoptosis and cancer. , 2000, 2, 180-190. | | 2 |
| 62 | Hypoxia classifier for transcriptome datasets. <i>BMC Bioinformatics</i> , 2022, 23, . | 1.2 | 1 |
| 63 | Hypoxia-inducible factor and cancer. , 2004, 6, 3-11. | | 0 |