

Ken-ichiro Kubo

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9107366/publications.pdf>

Version: 2024-02-01

45
papers

3,278
citations

201385

27
h-index

253896

43
g-index

46
all docs

46
docs citations

46
times ranked

4057
citing authors

#	ARTICLE	IF	CITATIONS
1	Rhythmic activation of excitatory neurons in the mouse frontal cortex improves the prefrontal cortex-mediated cognitive function. <i>Cerebral Cortex</i> , 2022, 32, 5243-5258.	1.6	1
2	Dab1-deficient deep layer neurons prevent Dab1-deficient superficial layer neurons from entering the cortical plate. <i>Neuroscience Research</i> , 2022, 180, 23-35.	1.0	2
3	Comprehensive characterization of migration profiles of murine cerebral cortical neurons during development using FlashTag labeling. <i>IScience</i> , 2021, 24, 102277.	1.9	15
4	Increased densities of white matter neurons as a cross-disease feature of neuropsychiatric disorders. <i>Psychiatry and Clinical Neurosciences</i> , 2020, 74, 166-175.	1.0	8
5	Human neocortical development as a basis to understand mechanisms underlying neurodevelopmental disabilities in extremely preterm infants. <i>Journal of Obstetrics and Gynaecology Research</i> , 2020, 46, 2242-2250.	0.6	1
6	Both excitatory and inhibitory neurons transiently form clusters at the outermost region of the developing mammalian cerebral neocortex. <i>Journal of Comparative Neurology</i> , 2019, 527, 1577-1597.	0.9	11
7	Drebrin-like (Dbnl) Controls Neuronal Migration via Regulating N-Cadherin Expression in the Developing Cerebral Cortex. <i>Journal of Neuroscience</i> , 2019, 39, 678-691.	1.7	22
8	ApoER2 Controls Not Only Neuronal Migration in the Intermediate Zone But Also Termination of Migration in the Developing Cerebral Cortex. <i>Cerebral Cortex</i> , 2018, 28, 223-235.	1.6	34
9	Reelin transiently promotes N-cadherin-dependent neuronal adhesion during mouse cortical development. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2017, 114, 2048-2053.	3.3	46
10	Excessive activation of AhR signaling disrupts neuronal migration in the hippocampal CA1 region in the developing mouse. <i>Journal of Toxicological Sciences</i> , 2017, 42, 25-30.	0.7	20
11	Association of impaired neuronal migration with cognitive deficits in extremely preterm infants. <i>JCI Insight</i> , 2017, 2, .	2.3	21
12	Impaired dendritic growth and positioning of cortical pyramidal neurons by activation of aryl hydrocarbon receptor signaling in the developing mouse. <i>PLoS ONE</i> , 2017, 12, e0183497.	1.1	11
13	In Utero Bisphenol A Exposure Induces Abnormal Neuronal Migration in the Cerebral Cortex of Mice. <i>Frontiers in Endocrinology</i> , 2016, 7, 7.	1.5	8
14	Reelin and Neuropsychiatric Disorders. <i>Frontiers in Cellular Neuroscience</i> , 2016, 10, 229.	1.8	143
15	SUMOylation of DISC1: A Potential Role in Neural Progenitor Proliferation in the Developing Cortex. <i>Molecular Neuropsychiatry</i> , 2016, 2, 20-27.	3.0	4
16	DISC1 a key molecular lead in psychiatry and neurodevelopment: No-More Disrupted-in-Schizophrenia 1. <i>Molecular Psychiatry</i> , 2016, 21, 1488-1489.	4.1	61
17	Resilience in schizophrenia: A comparative study between a remote island and an urban area in Japan. <i>Schizophrenia Research</i> , 2016, 171, 92-96.	1.1	9
18	The COUP-TFII/Neuropilin-2 is a molecular switch steering diencephalon-derived GABAergic neurons in the developing mouse brain. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2015, 112, E4985-94.	3.3	37

#	ARTICLE	IF	CITATIONS
19	Cellular dynamics of neuronal migration in the hippocampus. <i>Frontiers in Neuroscience</i> , 2015, 9, 135.	1.4	51
20	Developmental origin of abnormal dendritic growth in the mouse brain induced by in utero disruption of aryl hydrocarbon receptor signaling. <i>Neurotoxicology and Teratology</i> , 2015, 52, 42-50.	1.2	35
21	Reelin has a preventive effect on phencyclidine-induced cognitive and sensory-motor gating deficits. <i>Neuroscience Research</i> , 2015, 96, 30-36.	1.0	30
22	Importance of Reelin C-Terminal Region in the Development and Maintenance of the Postnatal Cerebral Cortex and Its Regulation by Specific Proteolysis. <i>Journal of Neuroscience</i> , 2015, 35, 4776-4787.	1.7	64
23	Neuronal Heterotopias Affect the Activities of Distant Brain Areas and Lead to Behavioral Deficits. <i>Journal of Neuroscience</i> , 2015, 35, 12432-12445.	1.7	36
24	Reelin receptors ApoER2 and VLDLR are expressed in distinct spatiotemporal patterns in developing mouse cerebral cortex. <i>Journal of Comparative Neurology</i> , 2015, 523, 463-478.	0.9	53
25	Characterization of the dipeptide repeat protein in the molecular pathogenesis of c9FTD/ALS. <i>Human Molecular Genetics</i> , 2015, 24, 1630-1645.	1.4	136
26	Hippocampal Pyramidal Neurons Switch from a Multipolar Migration Mode to a Novel "Climbing" Migration Mode during Development. <i>Journal of Neuroscience</i> , 2014, 34, 1115-1126.	1.7	61
27	How does Reelin control neuronal migration and layer formation in the developing mammalian neocortex?. <i>Neuroscience Research</i> , 2014, 86, 50-58.	1.0	111
28	Dab1-mediated colocalization of multi-adaptor protein CIN85 with Reelin receptors, ApoER2 and VLDLR, in neurons. <i>Genes To Cells</i> , 2013, 18, 410-424.	0.5	10
29	Reelin Controls Neuronal Positioning by Promoting Cell-Matrix Adhesion via Inside-Out Activation of Integrin $\alpha 5 \beta 1$. <i>Neuron</i> , 2012, 76, 353-369.	3.8	156
30	DISC1-dependent switch from progenitor proliferation to migration in the developing cortex. <i>Nature</i> , 2011, 473, 92-96.	13.7	181
31	Reelin inhibits migration of sympathetic preganglionic neurons in the spinal cord of the chick. <i>Journal of Comparative Neurology</i> , 2011, 519, 1970-1978.	0.9	4
32	The Outermost Region of the Developing Cortical Plate Is Crucial for Both the Switch of the Radial Migration Mode and the Dab1-Dependent "Inside-Out" Lamination in the Neocortex. <i>Journal of Neuroscience</i> , 2011, 31, 9426-9439.	1.7	104
33	GABAergic Precursor Transplantation into the Prefrontal Cortex Prevents Phencyclidine-Induced Cognitive Deficits. <i>Journal of Neuroscience</i> , 2011, 31, 14116-14125.	1.7	49
34	Disrupted-in-Schizophrenia-1 (Disc1) is necessary for migration of the pyramidal neurons during mouse hippocampal development. <i>Human Molecular Genetics</i> , 2011, 20, 2834-2845.	1.4	55
35	Ectopic Reelin Induces Neuronal Aggregation with a Normal Birthdate-Dependent "Inside-Out" Alignment in the Developing Neocortex. <i>Journal of Neuroscience</i> , 2010, 30, 10953-10966.	1.7	68
36	Migration defects by DISC1 knockdown in C57BL/6, 129X1/SvJ, and ICR strains via in utero gene transfer and virus-mediated RNAi. <i>Biochemical and Biophysical Research Communications</i> , 2010, 400, 631-637.	1.0	38

#	ARTICLE	IF	CITATIONS
37	Knockdown of DISC1 by In Utero Gene Transfer Disturbs Postnatal Dopaminergic Maturation in the Frontal Cortex and Leads to Adult Behavioral Deficits. <i>Neuron</i> , 2010, 65, 480-489.	3.8	275
38	Rab GTPases-Dependent Endocytic Pathways Regulate Neuronal Migration and Maturation through N-Cadherin Trafficking. <i>Neuron</i> , 2010, 67, 588-602.	3.8	291
39	Recruitment of PCM1 to the Centrosome by the Cooperative Action of DISC1 and BBS4. <i>Archives of General Psychiatry</i> , 2008, 65, 996.	13.8	124
40	Regulation of the interaction of Disabled-1 with CIN85 by phosphorylation with Cyclin-dependent kinase 5. <i>Genes To Cells</i> , 2007, 12, 1315-1327.	0.5	17
41	A schizophrenia-associated mutation of DISC1 perturbs cerebral cortex development. <i>Nature Cell Biology</i> , 2005, 7, 1167-1178.	4.6	532
42	Cell and molecular mechanisms that control cortical layer formation in the brain. <i>Keio Journal of Medicine</i> , 2003, 52, 8-20.	0.5	22
43	Secreted Reelin molecules form homodimers. <i>Neuroscience Research</i> , 2002, 43, 381-388.	1.0	98
44	Reelin molecules assemble together to form a large protein complex, which is inhibited by the function-blocking CR-50 antibody. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2000, 97, 9729-9734.	3.3	126
45	Pax-6 is required for thalamocortical pathway formation in fetal rats. , 1999, 408, 147-160.		97