

Che-Yi Chou

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/9110559/che-yi-chou-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

40
papers

380
citations

10
h-index

18
g-index

42
ext. papers

484
ext. citations

3.6
avg, IF

3.16
L-index

#	Paper	IF	Citations
40	Incidence of different types of subsequent fractures and related mortality in Taiwan.. <i>Archives of Osteoporosis</i> , 2022 , 17, 55	2.9	0
39	Hip fracture in patients with non-dialysis chronic kidney disease stage 5. <i>Scientific Reports</i> , 2021 , 11, 20591	4.1	0
38	Geriatric Functional Impairment Using the Integrated Care for Older People (ICOPE) Approach in Community-Dwelling Elderly and Its Association with Dyslipidemia. <i>Vascular Health and Risk Management</i> , 2021 , 17, 389-394	4.4	2
37	Urine phthalate metabolites are associated with urothelial cancer in chronic kidney disease patients. <i>Chemosphere</i> , 2021 , 273, 127834	8.4	0
36	Association of diabetes, education level, and care dependency with use of temporary vascular access in patients with chronic kidney disease. <i>Seminars in Dialysis</i> , 2021 , 34, 130-136	2.5	
35	A Matrix Metalloproteinase-2-Based Nomogram to Assess the Risk of Encapsulating Peritoneal Sclerosis in Peritoneal Dialysis Patients. <i>BioMed Research International</i> , 2021 , 2021, 6666441	3	1
34	The MicroRNA Prediction Models as Ancillary Diagnosis Biomarkers for Urothelial Carcinoma in Patients With Chronic Kidney Disease. <i>Frontiers in Medicine</i> , 2021 , 8, 726214	4.9	1
33	Text mining in a literature review of urothelial cancer using topic model. <i>BMC Cancer</i> , 2020 , 20, 462	4.8	6
32	Identification of GSN and LAMC2 as Key Prognostic Genes of Bladder Cancer by Integrated Bioinformatics Analysis. <i>Cancers</i> , 2020 , 12,	6.6	10
31	Early prediction of acquiring acute kidney injury for older inpatients using most effective laboratory test results. <i>BMC Medical Informatics and Decision Making</i> , 2020 , 20, 36	3.6	3
30	Wilforine resensitizes multidrug resistant cancer cells via competitive inhibition of P-glycoprotein. <i>Phytomedicine</i> , 2020 , 71, 153239	6.5	7
29	Association between asthma and risk of benign prostatic hyperplasia: a retrospective population-based study. <i>Aging Male</i> , 2020 , 23, 599-606	2.1	2
28	Urinary tract infection is associated with hypokalemia: a case control study. <i>BMC Urology</i> , 2020 , 20, 108	2.2	0
27	Maximum Glomerular Filtration Decline Rate is Associated with Mortality and Poor Renal Outcome in Chronic Kidney Disease Patients. <i>Blood Purification</i> , 2019 , 48, 131-137	3.1	1
26	Development and validation of a nomogram for urothelial cancer in patients with chronic kidney disease. <i>Scientific Reports</i> , 2019 , 9, 3473	4.9	4
25	Abdominal aortic calcification score predicts the occurrence of coronary artery disease in middle-aged peritoneal dialysis patients. <i>Nephrology</i> , 2019 , 24, 336-340	2.2	6
24	Hypothyroidism is associated with all-cause mortality in a national cohort of chronic haemodialysis patients. <i>Nephrology</i> , 2018 , 23, 559-564	2.2	4

23	Abdominal aortic calcification score can predict future coronary artery disease in hemodialysis patients: a 5-year prospective cohort study. <i>BMC Nephrology</i> , 2018 , 19, 313	2.7	9
22	Indirect comparison of efficacy and safety between immune checkpoint inhibitors and antiangiogenic therapy in advanced non-small-cell lung cancer. <i>Scientific Reports</i> , 2018 , 8, 9686	4.9	0
21	Prolonged QT interval is linked to all-cause and cardiac mortality in chronic peritoneal dialysis patients. <i>Nephrology</i> , 2017 , 22, 436-440	2.2	5
20	Increased Risk of Bullous Pemphigoid after First-Ever Stroke: A Population-Based Study. <i>Neurodegenerative Diseases</i> , 2017 , 17, 166-170	2.3	5
19	Qi deficiency is associated with depression in chronic hemodialysis patients. <i>Complementary Therapies in Medicine</i> , 2017 , 30, 102-106	3.5	5
18	Effect of Long-term Incretin-Based Therapies on Ischemic Heart Diseases in Patients with Type 2 Diabetes Mellitus: A Network Meta-analysis. <i>Scientific Reports</i> , 2017 , 7, 15795	4.9	11
17	Transformation of 5-D itch scale and numerical rating scale in chronic hemodialysis patients. <i>BMC Nephrology</i> , 2017 , 18, 56	2.7	18
16	The Shortest QRS Duration of an Electrocardiogram Might Be an Optimal Electrocardiographic Predictor for Response to Cardiac Resynchronization Therapy. <i>International Heart Journal</i> , 2017 , 58, 530-535	1.8	3
15	Chronic Osteomyelitis Is Associated With Increased Risk of New-Onset Atrial Fibrillation: Evidence From a Nationwide Cohort of 23 Million People. <i>Canadian Journal of Cardiology</i> , 2016 , 32, 1388-1395	3.8	2
14	Upper gastrointestinal bleeding as a risk factor for dialysis and all-cause mortality: a cohort study of chronic kidney disease patients in Taiwan. <i>BMJ Open</i> , 2016 , 6, e010439	3	8
13	Loss of Residual Renal Function is Associated With Vascular Calcification in Hemodialysis Patients. <i>Therapeutic Apheresis and Dialysis</i> , 2016 , 20, 27-30	1.9	6
12	Multidisciplinary care in patients with chronic kidney disease: A systematic review and meta-analysis. <i>European Journal of Internal Medicine</i> , 2015 , 26, 640-5	3.9	44
11	Topiramate may not increase risk of urolithiasis: A nationwide population-based cohort study. <i>Seizure: the Journal of the British Epilepsy Association</i> , 2015 , 29, 86-9	3.2	6
10	Hyperphosphate-Induced Myocardial Hypertrophy through the GATA-4/NFAT-3 Signaling Pathway Is Attenuated by ERK Inhibitor Treatment. <i>CardioRenal Medicine</i> , 2015 , 5, 79-88	2.8	21
9	Longer dialysis times are associated with a lower chance of dementia among chronic hemodialysis patients. <i>Therapeutic Apheresis and Dialysis</i> , 2015 , 19, 50-5	1.9	
8	Patients with diabetes as the primary kidney disease have a worse survival than patients with comorbid diabetes in chronic haemodialysis patients. <i>Nephrology</i> , 2015 , 20, 155-60	2.2	4
7	Upper gastrointestinal bleeding in patients with CKD. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2014 , 9, 1354-9	6.9	27
6	Peptic ulcer disease risk in chronic kidney disease: ten-year incidence, ulcer location, and ulcerogenic effect of medications. <i>PLoS ONE</i> , 2014 , 9, e87952	3.7	33

5	Association of walking with survival and RRT among patients with CKD stages 3-5. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2014 , 9, 1183-9	6.9	62
4	Comparing risk of new onset diabetes mellitus in chronic kidney disease patients receiving peritoneal dialysis and hemodialysis using propensity score matching. <i>PLoS ONE</i> , 2014 , 9, e87891	3.7	19
3	Waist-to-Height Ratio and Obesity in Chinese 2012 , 2007-2015		
2	Norepinephrine and hospital mortality in critically ill patients undergoing continuous renal replacement therapy. <i>Artificial Organs</i> , 2011 , 35, E11-7	2.6	13
1	Waist-to-height ratio is the best index of obesity in association with chronic kidney disease. <i>Nutrition</i> , 2007 , 23, 788-93	4.8	32