David Centurion

List of Publications by Citations

Source: https://exaly.com/author-pdf/9111014/david-centurion-publications-by-citations.pdf

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

79 papers 1,399 citations h-index g-index

81 1,513 5 4.06 ext. papers ext. citations avg, IF L-index

#	Paper	IF	Citations
79	Cardiovascular responses produced by 5-hydroxytriptamine:a pharmacological update on the receptors/mechanisms involved and therapeutic implications. <i>Naunyn-Schmiedebergts Archives of Pharmacology</i> , 2007 , 376, 45-63	3.4	154
78	Migraine: pathophysiology, pharmacology, treatment and future trends. <i>Current Vascular Pharmacology</i> , 2003 , 1, 71-84	3.3	98
77	Pharmacological profile of the clonidine-induced inhibition of vasodepressor sensory outflow in pithed rats: correlation with a2A/2C-adrenoceptors. <i>British Journal of Pharmacology</i> , 2008 , 154, 1160-1	186	78
76	Cardiovascular alterations after spinal cord injury: an overview. <i>Current Medicinal Chemistry Cardiovascular and Hematological Agents</i> , 2004 , 2, 133-48		69
75	Canine external carotid vasoconstriction to methysergide, ergotamine and dihydroergotamine: role of 5-HT1B/1D receptors and alpha2-adrenoceptors. <i>British Journal of Pharmacology</i> , 1999 , 126, 585-94	8.6	51
74	Mediation of 5-HT-induced external carotid vasodilatation in GR 127935-pretreated vagosympathectomized dogs by the putative 5-HT7 receptor. <i>British Journal of Pharmacology</i> , 1997 , 120, 1319-27	8.6	50
73	The canine external carotid vasoconstrictor 5-HT1 receptor: blockade by 5-HT1B (SB224289), but not by 5-HT1D (BRL15572) receptor antagonists. <i>European Journal of Pharmacology</i> , 1998 , 362, 69-72	5.3	40
72	Experimental migraine models and their relevance in migraine therapy. <i>Cephalalgia</i> , 2006 , 26, 642-59	6.1	36
71	The suprachiasmatic nucleus is part of a neural feedback circuit adapting blood pressure response. <i>Neuroscience</i> , 2014 , 266, 197-207	3.9	35
70	The 5-HT1-like receptors mediating inhibition of sympathetic vasopressor outflow in the pithed rat: operational correlation with the 5-HT1A, 5-HT1B and 5-HT1D subtypes. <i>British Journal of Pharmacology</i> , 1998 , 124, 1001-11	8.6	35
69	Characterization of putative 5-HT7 receptors mediating tachycardia in the cat. <i>British Journal of Pharmacology</i> , 1997 , 121, 1187-95	8.6	32
68	5-HT7, but not 5-HT2B, receptors mediate hypotension in vagosympathectomized rats. <i>European Journal of Pharmacology</i> , 2004 , 502, 239-42	5.3	32
67	Pharmacological profile of the 5-HT-induced inhibition of cardioaccelerator sympathetic outflow in pithed rats: correlation with 5-HT1 and putative 5-ht5A/5B receptors. <i>British Journal of Pharmacology</i> , 2003 , 140, 725-35	8.6	29
66	Further characterization of the 5-HT1 receptors mediating cardiac sympatho-inhibition in pithed rats: pharmacological correlation with the 5-HT1B and 5-HT1D subtypes. <i>Naunyn-Schmiedebergts Archives of Pharmacology</i> , 2004 , 369, 220-7	3.4	28
65	The 5-HT(1) receptors inhibiting the rat vasodepressor sensory CGRPergic outflow: further involvement of 5-HT(1F), but not 5-HT(1A) or 5-HT(1D), subtypes. <i>European Journal of Pharmacology</i> , 2011 , 659, 233-43	5.3	27
64	Current and prospective pharmacological targets in relation to antimigraine action. <i>Naunyn-Schmiedebergts Archives of Pharmacology</i> , 2008 , 378, 371-94	3.4	26
63	Potential vascular alpha1-adrenoceptor blocking properties of an array of 5-HT receptor ligands in the rat. <i>European Journal of Pharmacology</i> , 2006 , 535, 234-42	5.3	26

(2001-2008)

62	Pharmacological profile of the clonidine-induced inhibition of vasodepressor sensory outflow in pithed rats: correlation with alpha(2A/2C)-adrenoceptors. <i>British Journal of Pharmacology</i> , 2008 , 154, 51-9	8.6	24	
61	The atypical 5-HT2 receptor mediating tachycardia in pithed rats: pharmacological correlation with the 5-HT2A receptor subtype. <i>British Journal of Pharmacology</i> , 2002 , 135, 1531-9	8.6	23	
60	5-Hydroxytryptamine inhibits the tachycardia induced by selective preganglionic sympathetic stimulation in pithed rats. <i>Life Sciences</i> , 1999 , 64, 1839-47	6.8	22	
59	Operational characteristics of the 5-HT1-like receptors mediating external carotid vasoconstriction in vagosympathectomized dogs. Close resemblance to the 5-HT1D receptor subtype. <i>Naunyn-Schmiedebergts Archives of Pharmacology</i> , 1996 , 354, 550-6	3.4	22	
58	Characterization of the postjunctional alpha 2C-adrenoceptor mediating vasoconstriction to UK14304 in porcine pulmonary veins. <i>British Journal of Pharmacology</i> , 2007 , 151, 186-94	8.6	21	
57	5-HT1B receptors, alpha2A/2C- and, to a lesser extent, alpha1-adrenoceptors mediate the external carotid vasoconstriction to ergotamine in vagosympathectomised dogs. <i>Naunyn-Schmiedebergts Archives of Pharmacology</i> , 2004 , 370, 46-53	3.4	21	
56	Mediation of 5-HT-induced internal carotid vasodilatation in GR127935- and ritanserin-pretreated dogs by 5-HT7 receptors. <i>Naunyn-Schmiedebergts Archives of Pharmacology</i> , 2000 , 362, 169-76	3.4	21	
55	Donitriptan, but not sumatriptan, inhibits capsaicin-induced canine external carotid vasodilatation via 5-HT1B rather than 5-HT1D receptors. <i>British Journal of Pharmacology</i> , 2006 , 149, 82-91	8.6	19	
54	5-HT1B receptors and alpha 2A/2C-adrenoceptors mediate external carotid vasoconstriction to dihydroergotamine. <i>European Journal of Pharmacology</i> , 2004 , 484, 287-90	5.3	19	
53	Pharmacological evidence that alpha2A- and alpha2C-adrenoceptors mediate the inhibition of cardioaccelerator sympathetic outflow in pithed rats. <i>European Journal of Pharmacology</i> , 2007 , 554, 20	5- ⁵ t-3	17	
52	Further pharmacological analysis of the orphan 5-HT receptors mediating feline vasodepressor responses: close resemblance to the 5-HT7, receptor. <i>Naunyn-Schmiedebergts Archives of Pharmacology</i> , 2000 , 361, 665-71	3.4	17	
51	Crosstalk of vascular 5-HT1 receptors with other receptors: clinical implications. Neuropharmacology, 2008, 55, 986-93	5.5	16	
50	Unravelling the pharmacological profile of the canine external carotid vasodilator U -HT1-likeU receptors: coexistence of sympatho-inhibitory 5-HT1B and postjunctional 5-HT7 receptors. <i>Naunyn-Schmiedebergts Archives of Pharmacology</i> , 2001 , 363, 73-80	3.4	16	
49	Spinal sumatriptan inhibits capsaicin-induced canine external carotid vasodilatation via 5-HT1B rather than 5-HT1D receptors. <i>European Journal of Pharmacology</i> , 2009 , 615, 133-8	5.3	14	
48	Activation of 5-HT1B receptors inhibits the vasodepressor sensory CGRPergic outflow in pithed rats. <i>European Journal of Pharmacology</i> , 2010 , 637, 131-7	5.3	14	
47	Pharmacological characterization of the inhibition by moxonidine and agmatine on the cardioaccelerator sympathetic outflow in pithed rats. <i>European Journal of Pharmacology</i> , 2009 , 616, 17	5-82	13	
46	Evidence that some imidazoline derivatives inhibit peripherally the vasopressor sympathetic outflow in pithed rats. <i>Autonomic Neuroscience: Basic and Clinical</i> , 2008 , 143, 40-5	2.4	13	
45	The GR127935-sensitive 5-HT(1) receptors mediating canine internal carotid vasoconstriction: resemblance to the 5-HT(1B), but not to the 5-HT(1D) or 5-ht(1F), receptor subtype. <i>British Journal of Pharmacology</i> 2001 , 132, 991-8	8.6	13	

44	The 🛘 -adrenoceptors mediating inhibition of the vasopressor sympathetic outflow in pithed rats: pharmacological correlation with 🖺 A, 🖺 B and 🗸 C subtypes. <i>European Journal of Pharmacology</i> , 2013 , 718, 245-52	5.3	12
43	Evidence for 5-HT(1B/1D) and 5-HT(2A) receptors mediating constriction of the canine internal carotid circulation. <i>British Journal of Pharmacology</i> , 2001 , 132, 983-90	8.6	11
42	Fructose-Induced Insulin Resistance as a Model of Neuropathic Pain in Rats. <i>Neuroscience</i> , 2019 , 404, 233-245	3.9	10
41	Postjunctional IC-adrenoceptors mediate vasoconstriction in rat tail artery: influence of precontraction and temperature on vasoreactivity. <i>Naunyn-Schmiedebergts Archives of Pharmacology</i> , 2010 , 382, 487-97	3.4	10
40	Pharmacological evidence that NaHS inhibits the vasopressor responses induced by stimulation of the preganglionic sympathetic outflow in pithed rats. <i>European Journal of Pharmacology</i> , 2016 , 770, 40-	5 5.3	9
39	Pharmacological characterization of the mechanisms involved in the vasorelaxation induced by progesterone and 17Destradiol on isolated canine basilar and internal carotid arteries. <i>Steroids</i> , 2014 , 89, 33-40	2.8	9
38	Pharmacological evidence that Call+ channels and, to a lesser extent, K+ channels mediate the relaxation of testosterone in the canine basilar artery. <i>Steroids</i> , 2011 , 76, 409-15	2.8	9
37	The dopamine receptors mediating inhibition of the sympathetic vasopressor outflow in pithed rats: pharmacological correlation with the D(2) -like type. <i>Basic and Clinical Pharmacology and Toxicology</i> , 2011 , 109, 506-12	3.1	9
36	Phenylephrine contracts porcine pulmonary veins via alpha(1B)-, alpha(1D)-, and alpha(2)-adrenoceptors. <i>European Journal of Pharmacology</i> , 2009 , 613, 86-92	5.3	9
35	Pharmacological characterization of ergotamine-induced inhibition of the cardioaccelerator sympathetic outflow in pithed rats. <i>Naunyn-Schmiedebergts Archives of Pharmacology</i> , 2009 , 379, 137-48	3.4	9
34	Effect of some acute and prophylactic antimigraine drugs on the vasodepressor sensory CGRPergic outflow in pithed rats. <i>Life Sciences</i> , 2009 , 84, 125-31	6.8	9
33	Clonidine inhibits the canine external carotid vasodilatation to capsaicin by alpha2A/2C-adrenoceptors. <i>European Journal of Pharmacology</i> , 2006 , 543, 68-76	5.3	9
32	Pharmacological profile of the vascular responses to dopamine in the canine external carotid circulation. <i>Basic and Clinical Pharmacology and Toxicology</i> , 2003 , 92, 165-72		9
31	Pharmacological analysis of the mechanisms involved in the tachycardic and vasopressor responses to the antimigraine agent, isometheptene, in pithed rats. <i>Life Sciences</i> , 2004 , 74, 3223-34	6.8	9
30	A-adrenoceptors, but not nitric oxide, mediate the peripheral cardiac sympatho-inhibition of moxonidine. <i>European Journal of Pharmacology</i> , 2016 , 782, 35-43	5.3	8
29	Pharmacological evidence that spinal (PC)- and, to a lesser extent, (PA)-adrenoceptors inhibit capsaicin-induced vasodilatation in the canine external carotid circulation. <i>European Journal of Pharmacology</i> , 2012 , 683, 204-10	5.3	8
28	Pharmacological identification of the Endrenoceptor subtypes mediating the vasopressor responses to B-HT 933 in pithed rats. <i>European Journal of Pharmacology</i> , 2012 , 691, 118-24	5.3	8
27	Pharmacological evaluation of metformin and N-benzylbiguanide, a novel analogue of metformin, on the vasopressor responses to adrenergic system stimulation in pithed rats with fructose-induced insulin resistance. <i>European Journal of Pharmacology</i> , 2017 , 814, 313-323	5.3	7

(2013-2019)

26	Histopathological and biochemical changes in the development of nonalcoholic fatty liver disease induced by high-sucrose diet at different times. <i>Canadian Journal of Physiology and Pharmacology</i> , 2019 , 97, 23-36	2.4	7	
25	Lack of effect of the adenosine A1 receptor agonist, GR79236, on capsaicin-induced CGRP release in anaesthetized pigs. <i>Cephalalgia</i> , 2005 , 25, 1082-90	6.1	6	
24	Chronic administration of NaHS and L-Cysteine restores cardiovascular changes induced by high-fat diet in rats. <i>European Journal of Pharmacology</i> , 2019 , 863, 172707	5.3	5	
23	Synthesis and In Vitro AMPK Activation of Cycloalkyl/Alkarylbiguanides with Robust In Vivo Antihyperglycemic Action. <i>Journal of Chemistry</i> , 2017 , 2017, 1-8	2.3	5	
22	Therapeutic effect of treatment with metformin and/or 4-hydroxychalcone in male Wistar rats with nonalcoholic fatty liver disease. <i>European Journal of Pharmacology</i> , 2019 , 863, 172699	5.3	4	
21	Pharmacological identification of 1 - and 2 -adrenoceptor subtypes involved in the vasopressor responses induced by ergotamine in pithed rats. <i>European Journal of Pharmacology</i> , 2013 , 715, 262-9	5.3	4	
20	A61603-induced contractions of the porcine meningeal artery are mediated by alpha1- and alpha2-adrenoceptors. <i>Basic and Clinical Pharmacology and Toxicology</i> , 2007 , 100, 279-85	3.1	4	
19	Pharmacological analysis of the cardiac sympatho-inhibitory actions of moxonidine and agmatine in pithed spontaneously hypertensive rats. <i>European Journal of Pharmacology</i> , 2016 , 791, 25-36	5.3	4	
18	NaHS prejunctionally inhibits the cardioaccelerator sympathetic outflow in pithed rats. <i>European Journal of Pharmacology</i> , 2018 , 823, 35-40	5.3	3	
17	Pharmacological evidence that metformin blocks the vasopressor responses mediated by stimulation of Eand Endrenoceptors in pithed rats. <i>European Journal of Pharmacology</i> , 2018 , 820, 130-13	7 5.3	3	
16	Pharmacological profile of the inhibition by dihydroergotamine and methysergide on the cardioaccelerator sympathetic outflow in pithed rats. <i>European Journal of Pharmacology</i> , 2009 , 612, 80-	6 ^{5.3}	3	
15	GR127935 antagonizes the 5-HT1-like receptor-mediated external carotid vasoconstriction in vagosympathectomized dogs. <i>Annals of the New York Academy of Sciences</i> , 1997 , 812, 207-8	6.5	3	
14	Activation of 5-HT and 5-HT receptors enhanced a positively reinforced long-term memory. Behavioural Brain Research, 2021 , 397, 112932	3.4	3	
13	Inhibitory effect of chronic oral treatment with fluoxetine on capsaicin-induced external carotid vasodilatation in anaesthetised dogs. <i>Cephalalgia</i> , 2015 , 35, 1041-53	6.1	2	
12	Pharmacological evidence that 5-HT1A/1B/1D, 2-adrenoceptors and D2-like receptors mediate ergotamine-induced inhibition of the vasopressor sympathetic outflow in pithed rats. <i>European Journal of Pharmacology</i> , 2014 , 740, 512-21	5.3	2	
11	Hydrogen Sulfide Subchronic Treatment Improves Hypertension Induced by Traumatic Brain Injury in Rats through Vasopressor Sympathetic Outflow Inhibition. <i>Journal of Neurotrauma</i> , 2021 ,	5.4	2	
10	Evidence that chronic administration of 17D-oestradiol decreases the vasopressor responses to adrenergic system stimulation in streptozotocin-diabetic female rats. <i>Steroids</i> , 2014 , 83, 1-9	2.8	1	
9	Pharmacological evidence that dopamine inhibits the cardioaccelerator sympathetic outflow via D2-like receptors in pithed rats. <i>Journal of Pharmacological Sciences</i> , 2013 , 123, 380-91	3.7	1	

8	Cardiovascular Responses to 5-hydroxytryptamine in Methimazole-induced Hypothyroid Pithed Rats. <i>Archives of Medical Research</i> , 2020 , 51, 310-316	6.6	1
7	Fenofibrate Protects Cardiomyocytes from Hypoxia/Reperfusion- and High Glucose-Induced Detrimental Effects. <i>PPAR Research</i> , 2021 , 2021, 8895376	4.3	1
6	Potential vascular Endrenoceptor blocking properties of metformin in rat aorta and tail artery. <i>European Journal of Pharmacology</i> , 2019 , 858, 172498	5.3	О
5	Sex-dependent antiallodynic effect of ladrenergic receptor agonist tizanidine in rats with experimental neuropathic pain <i>European Journal of Pharmacology</i> , 2022 , 174855	5.3	O
4	Blocking properties of terguride at the 5-HT receptor subtypes mediating cardiovascular responses in the rat. <i>Canadian Journal of Physiology and Pharmacology</i> , 2020 , 98, 511-521	2.4	
3	Pharmacological profile of the prejunctional 5-HT1 receptors mediating inhibition of sympathetic vasopressor outflow in the pithed rat. <i>Annals of the New York Academy of Sciences</i> , 1998 , 861, 281-2	6.5	
2	Bladder Enlargement Correlates with Plasma Insulin, Not Glucose Levels in Fructose-Fed Rats. <i>FASEB Journal</i> , 2020 , 34, 1-1	0.9	
1	Pharmacological characterization of 2 -adrenoceptor subtypes mediating inhibition of sympathetic vasopressor responses to B-HT 933 in pithed rats. <i>FASEB Journal</i> , 2013 , 27, lb605	0.9	