

# Ingebjörg Seljeflot

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9130466/publications.pdf>

Version: 2024-02-01

146  
papers

3,111  
citations

147801  
31  
h-index

206112  
48  
g-index

146  
all docs

146  
docs citations

146  
times ranked

4890  
citing authors

#	ARTICLE	IF	CITATIONS
1	Probiotics to HIV-Infected Immunological Nonresponders: Altered Mucosal Immunity and Microbial Diversity Restricted to Ileum. <i>Journal of Acquired Immune Deficiency Syndromes</i> (1999), 2022, 89, 77-86.	2.1	3
2	Human Immunodeficiency Virus-Infected Immunological Nonresponders Have Colon-Restricted Gut Mucosal Immune Dysfunction. <i>Journal of Infectious Diseases</i> , 2022, 225, 661-674.	4.0	16
3	Changes in eicosapentaenoic acid and docosahexaenoic acid and risk of cardiovascular events and atrial fibrillation: A secondary analysis of the OMEMI trial. <i>Journal of Internal Medicine</i> , 2022, 291, 637-647.	6.0	22
4	Complement ratios C3bc/C3 and sC5b-9/C5 do not increase the sensitivity of detecting acute complement activation systemically. <i>Molecular Immunology</i> , 2022, 141, 273-279.	2.2	2
5	Vascular Function in Norwegian Female Elite Runners: A Cross-Sectional, Controlled Study. <i>Sports</i> , 2022, 10, 37.	1.7	5
6	The NLRP3 inflammasome activation in subcutaneous, epicardial and pericardial adipose tissue in patients with coronary heart disease undergoing coronary by-pass surgery. <i>Atherosclerosis Plus</i> , 2022, 48, 47-54.	0.7	2
7	Gene expression of fibrinolytic markers in coronary thrombi. <i>Thrombosis Journal</i> , 2022, 20, 23.	2.1	0
8	Coagulation factors XI and XII as possible targets for anticoagulant therapy. <i>Thrombosis Research</i> , 2022, 214, 53-62.	1.7	12
9	Interleukin-6 inhibition in ST-elevation myocardial infarction: Immune cell profile in the randomised ASSAIL-MI trial. <i>EBioMedicine</i> , 2022, 80, 104013.	6.1	22
10	Exercise-induced change in circulating NT-proBNP could not distinguish between patients with and without coronary artery disease: the CADENCE study. <i>Scandinavian Cardiovascular Journal</i> , 2022, 56, 107-113.	1.2	0
11	Reduced leukocyte telomere lengths and sirtuin-1 gene expression in long-term survivors of type-1 diabetes: A Dialong substudy. <i>Journal of Diabetes Investigation</i> , 2021, 12, 1183-1192.	2.4	6
12	Effects of n-3 Fatty Acid Supplements in Elderly Patients After Myocardial Infarction. <i>Circulation</i> , 2021, 143, 528-539.	1.6	180
13	Differential associations of cardiac troponin T and cardiac troponin I with coronary artery pathology and dynamics in response to short-duration exercise. <i>Clinical Biochemistry</i> , 2021, 88, 23-29.	1.9	8
14	Biomarkers of ageing and cardiac remodeling are associated with atrial fibrillation. <i>Scandinavian Cardiovascular Journal</i> , 2021, 55, 213-219.	1.2	14
15	Randomized Trial of Interleukin-6 Receptor Inhibition in Patients With Acute ST-Segment Elevation Myocardial Infarction. <i>Journal of the American College of Cardiology</i> , 2021, 77, 1845-1855.	2.8	169
16	Gut related inflammation and cardiorespiratory fitness in patients with CAD and type 2 diabetes: a sub-study of a randomized controlled trial on exercise training. <i>Diabetology and Metabolic Syndrome</i> , 2021, 13, 36.	2.7	4
17	Effects of intermittent negative pressure treatment on circulating vascular biomarkers in patients with intermittent claudication. <i>Vascular Medicine</i> , 2021, 26, 489-496.	1.5	2
18	The Inflammasome Signaling Pathway Is Actively Regulated and Related to Myocardial Damage in Coronary Thrombi from Patients with STEMI. <i>Mediators of Inflammation</i> , 2021, 2021, 1-12.	3.0	7

#	ARTICLE	IF	CITATIONS
19	Tocilizumab increases citrullinated histone 3 in non-ST segment elevation myocardial infarction. <i>Open Heart</i> , 2021, 8, e001492.	2.3	4
20	Mortality and microbial diversity after allogeneic hematopoietic stem cell transplantation: secondary analysis of a randomized nutritional intervention trial. <i>Scientific Reports</i> , 2021, 11, 11593.	3.3	9
21	Transient Reduction of FMD-Response and L-Arginine Accompanied by Increased Levels of E-Selectin, VCAM, and ICAM after Prolonged Strenuous Exercise. <i>Sports</i> , 2021, 9, 86.	1.7	2
22	Effect of intermittent and continuous caloric restriction on Sirtuin1 concentration depends on sex and body mass index. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2021, 31, 1871-1878.	2.6	10
23	Reduced L-Arginine and L-Arginine-ADMA-Ratio, and Increased SDMA after Norseman Xtreme Triathlon. <i>Sports</i> , 2021, 9, 120.	1.7	7
24	Rifaximin or <i>Saccharomyces boulardii</i> in heart failure with reduced ejection fraction: Results from the randomized GutHeart trial. <i>EBioMedicine</i> , 2021, 70, 103511.	6.1	34
25	Gut Leakage Markers in Response to Strenuous Exercise in Patients with Suspected Coronary Artery Disease. <i>Cells</i> , 2021, 10, 2193.	4.1	6
26	Immune complexes, innate immunity, and NETosis in ChAdOx1 vaccine-induced thrombocytopenia. <i>European Heart Journal</i> , 2021, 42, 4064-4072.	2.2	49
27	Complement activation is associated with poor outcome after out-of-hospital cardiac arrest. <i>Resuscitation</i> , 2021, 166, 129-136.	3.0	12
28	Serum Galectin-3 and Subsequent Risk of Coronary Heart Disease in Subjects With Childhood-Onset Type 1 Diabetes: A Cohort Study. <i>Diabetes Care</i> , 2021, 44, 810-816.	8.6	9
29	Serum Levels of Dihomo-Gamma ( $\Gamma^3$ )-Linolenic Acid (DGLA) Are Inversely Associated with Linoleic Acid and Total Death in Elderly Patients with a Recent Myocardial Infarction. <i>Nutrients</i> , 2021, 13, 3475.	4.1	9
30	One year of omega 3 polyunsaturated fatty acid supplementation does not reduce circulating prothrombotic microvesicles in elderly subjects after suffering a myocardial infarction. <i>Clinical Nutrition</i> , 2021, 40, 5674-5677.	5.0	5
31	Adiponectin in relation to exercise and physical performance in patients with type 2 diabetes and coronary artery disease. <i>Adipocyte</i> , 2021, 10, 612-620.	2.8	2
32	NETosis in Long-Term Type 1 Diabetes Mellitus and Its Link to Coronary Artery Disease. <i>Frontiers in Immunology</i> , 2021, 12, 799539.	4.8	4
33	High levels of interleukin-6 are associated with final infarct size and adverse clinical events in patients with STEMI. <i>Open Heart</i> , 2021, 8, e001869.	2.3	10
34	Soluble ST2 concentrations associate with in-hospital mortality and need for mechanical ventilation in unselected patients with COVID-19. <i>Open Heart</i> , 2021, 8, e001884.	2.3	9
35	Late awakening, prognostic factors and long-term outcome in out-of-hospital cardiac arrest – results of the prospective Norwegian Cardio-Respiratory Arrest Study (NORCAST). <i>Resuscitation</i> , 2020, 149, 170-179.	3.0	47
36	Cardiovascular rEmodelling in living kidney donorS with reduced glomerular filtration rate: rationale and design of the CENS study. <i>Blood Pressure</i> , 2020, 29, 123-134.	1.5	2

#	ARTICLE	IF	CITATIONS
37	Legumain is upregulated in acute cardiovascular events and associated with improved outcome - potentially related to anti-inflammatory effects on macrophages. <i>Atherosclerosis</i> , 2020, 296, 74-82.	0.8	14
38	Shorter Leukocyte Telomere Lengths in Healthy Relatives of Patients with Coronary Heart Disease. <i>Rejuvenation Research</i> , 2020, 23, 324-332.	1.8	9
39	Superiority of high sensitivity cardiac troponin T vs. I for long-term prognostic value in patients with chest pain; data from the Akershus cardiac Examination (ACE) 3 study. <i>Clinical Biochemistry</i> , 2020, 78, 10-17.	1.9	15
40	Very Long Chain Marine n-3 Polyunsaturated Fatty Acids in Atherothrombotic Heart Disease. A Brief Review, with a Focus on Metabolic Effects. <i>Nutrients</i> , 2020, 12, 3014.	4.1	4
41	Left ventricular dysfunction in COPD without pulmonary hypertension. <i>PLoS ONE</i> , 2020, 15, e0235075.	2.5	3
42	Circulating levels of the terminal complement complex are associated with hypercoagulability in patients with stable coronary artery disease. <i>Thrombosis Research</i> , 2020, 196, 106-108.	1.7	1
43	Markers of remodeling in subcutaneous adipose tissue are strongly associated with overweight and insulin sensitivity in healthy non-obese men. <i>Scientific Reports</i> , 2020, 10, 14055.	3.3	8
44	Collagen methionine sulfoxide and glucuronidine/LW-1 are markers of coronary artery disease in long-term survivors with type 1 diabetes. The Dialong study. <i>PLoS ONE</i> , 2020, 15, e0233174.	2.5	8
45	Double-Stranded DNA and NETs Components in Relation to Clinical Outcome After ST-Elevation Myocardial Infarction. <i>Scientific Reports</i> , 2020, 10, 5007.	3.3	22
46	Low fibre intake is associated with gut microbiota alterations in chronic heart failure. <i>ESC Heart Failure</i> , 2020, 7, 456-466.	3.1	56
47	Procoagulant activity in children and adolescents on intensive insulin therapy. <i>Pediatric Diabetes</i> , 2020, 21, 496-504.	2.9	2
48	Rosuvastatin alters the genetic composition of the human gut microbiome. <i>Scientific Reports</i> , 2020, 10, 5397.	3.3	20
49	Complement Activation in Association with Markers of Neutrophil Extracellular Traps and Acute Myocardial Infarction in Stable Coronary Artery Disease. <i>Mediators of Inflammation</i> , 2020, 2020, 1-9.	3.0	8
50	Annexin V+ Microvesicles in Children and Adolescents with Type 1 Diabetes: A Prospective Cohort Study. <i>Journal of Diabetes Research</i> , 2020, 2020, 1-8.	2.3	2
51	Neutrophil extracellular trap components and myocardial recovery in post-ischemic acute heart failure. <i>PLoS ONE</i> , 2020, 15, e0241333.	2.5	6
52	Left ventricular dysfunction in COPD without pulmonary hypertension. , 2020, 15, e0235075.		0
53	Left ventricular dysfunction in COPD without pulmonary hypertension. , 2020, 15, e0235075.		0
54	Left ventricular dysfunction in COPD without pulmonary hypertension. , 2020, 15, e0235075.		0

#	ARTICLE	IF	CITATIONS
55	Left ventricular dysfunction in COPD without pulmonary hypertension. , 2020, 15, e0235075.		0
56	Title is missing!. , 2020, 15, e0233174.		0
57	Title is missing!. , 2020, 15, e0233174.		0
58	Title is missing!. , 2020, 15, e0233174.		0
59	Title is missing!. , 2020, 15, e0233174.		0
60	Title is missing!. , 2020, 15, e0233174.		0
61	Title is missing!. , 2020, 15, e0233174.		0
62	Glucose associated NETosis in patients with ST-elevation myocardial infarction: an observational study. BMC Cardiovascular Disorders, 2019, 19, 221.	1.7	9
63	Novel biomolecules of ageing, sex differences and potential underlying mechanisms of telomere shortening in coronary artery disease. Experimental Gerontology, 2019, 119, 53-60.	2.8	23
64	A study of atherothrombotic biomarkers in welders. International Archives of Occupational and Environmental Health, 2019, 92, 1023-1031.	2.3	8
65	High Adherence to the Nordic Diet Is Associated with Lower Levels of Total and Platelet-Derived Circulating Microvesicles in a Norwegian Population. Nutrients, 2019, 11, 1114.	4.1	7
66	A brief review on resistance to P2Y12 receptor antagonism in coronary artery disease. Thrombosis Journal, 2019, 17, 11.	2.1	46
67	Effects of dietary intervention and n-3 PUFA supplementation on markers of gut-related inflammation and their association with cardiovascular events in a high-risk population. Atherosclerosis, 2019, 286, 53-59.	0.8	16
68	IgM antibodies against phosphorylcholine measured early after acute ST-elevation myocardial infarction in relation to atherosclerotic disease burden and long-term clinical outcome. PLoS ONE, 2019, 14, e0215640.	2.5	1
69	High-sensitive cardiac Troponin T and exercise stress test for evaluation of angiographically significant coronary disease. International Journal of Cardiology, 2019, 287, 1-6.	1.7	8
70	Elevated levels of circulating microvesicles in coronary artery disease patients with type 2 diabetes and albuminuria: Effects of exercise training. Diabetes and Vascular Disease Research, 2019, 16, 431-439.	2.0	10
71	Effects of exercise training on inflammasome-related mediators and their associations to glucometabolic variables in patients with combined coronary artery disease and type 2 diabetes mellitus: Sub-study of a randomized control trial. Diabetes and Vascular Disease Research, 2019, 16, 360-368.	2.0	14
72	Index of microvascular resistance to assess the effect of rosuvastatin on microvascular function in women with chest pain and no obstructive coronary artery disease: A double-blind randomized study. Catheterization and Cardiovascular Interventions, 2019, 94, 660-668.	1.7	10

#	ARTICLE	IF	CITATIONS
73	Rationale for the ASSAIL-MI-trial: a randomised controlled trial designed to assess the effect of tocilizumab on myocardial salvage in patients with acute ST-elevation myocardial infarction (STEMI). <i>Open Heart</i> , 2019, 6, e001108.	2.3	34
74	Effects of exercise training on markers of adipose tissue remodeling in patients with coronary artery disease and type 2 diabetes mellitus: sub study of the randomized controlled EXCADI trial. <i>Diabetology and Metabolic Syndrome</i> , 2019, 11, 109.	2.7	3
75	Neutrophil Extracellular Trap Components Associate with Infarct Size, Ventricular Function, and Clinical Outcome in STEMI. <i>Mediators of Inflammation</i> , 2019, 2019, 1-10.	3.0	33
76	Leukocyte telomere length and serum polyunsaturated fatty acids, dietary habits, cardiovascular risk factors and features of myocardial infarction in elderly patients. <i>BMC Geriatrics</i> , 2019, 19, 376.	2.7	10
77	Circulating markers of gut barrier function associated with disease severity in primary sclerosing cholangitis. <i>Liver International</i> , 2019, 39, 371-381.	3.9	51
78	Markers of neutrophil extracellular traps are associated with adverse clinical outcome in stable coronary artery disease. <i>European Journal of Preventive Cardiology</i> , 2018, 25, 762-769.	1.8	34
79	Antithrombotic therapy and body mass: an expert position paper of the ESC Working Group on Thrombosis. <i>European Heart Journal</i> , 2018, 39, 1672-1686f.	2.2	106
80	Effect of strenuous exercise on mediators of inflammation in patients with coronary artery disease. <i>Cytokine</i> , 2018, 105, 17-22.	3.2	13
81	Acute heart failure following myocardial infarction: complement activation correlates with the severity of heart failure in patients developing cardiogenic shock. <i>ESC Heart Failure</i> , 2018, 5, 292-301.	3.1	27
82	Impact of HIV and Type 2 diabetes on Gut Microbiota Diversity, Tryptophan Catabolism and Endothelial Dysfunction. <i>Scientific Reports</i> , 2018, 8, 6725.	3.3	35
83	Markers of Thrombin Generation Are Associated With Long-Term Clinical Outcome in Patients With ST-Segment Elevation Myocardial Infarction. <i>Clinical and Applied Thrombosis/Hemostasis</i> , 2018, 24, 1088-1094.	1.7	11
84	Preserved endothelial function in young adults with type 1 diabetes. <i>PLoS ONE</i> , 2018, 13, e0206523.	2.5	13
85	Soluble IL-1 receptor 2 is associated with left ventricular remodelling in patients with ST-elevation myocardial infarction. <i>International Journal of Cardiology</i> , 2018, 268, 187-192.	1.7	15
86	Association of IL-8 With Infarct Size and Clinical Outcomes in Patients With STEMI. <i>Journal of the American College of Cardiology</i> , 2018, 72, 187-198.	2.8	40
87	High-sensitivity troponin I in persistent atrial fibrillation â€“ relation to NT-proBNP and markers of inflammation and haemostasis. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2018, 78, 386-392.	1.2	2
88	Markers of metabolic endotoxemia as related to metabolic syndrome in an elderly male population at high cardiovascular risk: a cross-sectional study. <i>Diabetology and Metabolic Syndrome</i> , 2018, 10, 59.	2.7	35
89	Interleukin-18 and the NLR family pyrin domain containing-3 inflammasome in adipose tissue are strongly associated with glucometabolic variables in a cohort of middle-aged men. <i>Diabetes and Vascular Disease Research</i> , 2018, 15, 458-464.	2.0	4
90	Design of the GutHeartâ€”targeting gut microbiota to treat heart failureâ€”trial: a Phase II, randomized clinical trial. <i>ESC Heart Failure</i> , 2018, 5, 977-984.	3.1	39

#	ARTICLE	IF	CITATIONS
91	Increased arterial stiffness in childhood onset diabetes: a cardiovascular magnetic resonance study. European Heart Journal Cardiovascular Imaging, 2018, 19, 694-700.	1.2	12
92	Biomarkers of endothelial activation and thrombosis in tunnel construction workers exposed to airborne contaminants. International Archives of Occupational and Environmental Health, 2017, 90, 309-317.	2.3	5
93	Reduced endothelial activation after exercise is associated with improved HbA <sub>1c</sub> in patients with type 2 diabetes and coronary artery disease. Diabetes and Vascular Disease Research, 2017, 14, 94-103.	2.0	11
94	Procoagulant activity in patients with combined type 2 diabetes and coronary artery disease: No effects of long-term exercise training. Diabetes and Vascular Disease Research, 2017, 14, 144-151.	2.0	6
95	Changes in dietary pattern when including 700g of salmon per week to patients with atherosclerotic heart disease. Clinical Nutrition ESPEN, 2017, 19, 38-44.	1.2	2
96	Thrombin Generation in Patients With Suspected Venous Thromboembolism. Clinical and Applied Thrombosis/Hemostasis, 2017, 23, 416-421.	1.7	10
97	HIV-infected persons with type 2 diabetes show evidence of endothelial dysfunction and increased inflammation. BMC Infectious Diseases, 2017, 17, 234.	2.9	19
98	Monocyte-derived circulating microparticles (CD14+, CD14+/CD11b+ and CD14+/CD142+) are related to long-term prognosis for cardiovascular mortality in STEMI patients. International Journal of Cardiology, 2017, 227, 876-881.	1.7	47
99	Evaluation of circulating levels of Ccn2/connective tissue growth factor in patients with ST-elevation myocardial infarction. Scientific Reports, 2017, 7, 11945.	3.3	4
100	Pro-coagulant activity during exercise testing in patients with coronary artery disease. Thrombosis Journal, 2017, 15, 3.	2.1	9
101	A Double-Blinded Randomized Study Investigating a Possible Anti-Inflammatory Effect of Saxagliptin versus Placebo as Add-On Therapy in Patients with Both Type 2 Diabetes And Stable Coronary Artery Disease. Mediators of Inflammation, 2017, 2017, 1-9.	3.0	2
102	Osteoprotegerin levels in ST-elevation myocardial infarction: Temporal profile and association with myocardial injury and left ventricular function. PLoS ONE, 2017, 12, e0173034.	2.5	8
103	Reduced HDL function in children and young adults with type 1 diabetes. Cardiovascular Diabetology, 2017, 16, 85.	6.8	30
104	vWF/ADAMTS13 is associated with on-aspirin residual platelet reactivity and clinical outcome in patients with stable coronary artery disease. Thrombosis Journal, 2017, 15, 28.	2.1	12
105	Effect of sinus rhythm restoration on markers of thrombin generation in atrial fibrillation. Thrombosis Journal, 2017, 15, 30.	2.1	2
106	Platelet-, monocyte-derived and tissue factor-carrying circulating microparticles are related to acute myocardial infarction severity. PLoS ONE, 2017, 12, e0172558.	2.5	74
107	Effects of long-term exercise training on adipose tissue expression of fractalkine and MCP-1 in patients with type 2 diabetes and stable coronary artery disease: a substudy of a randomized controlled trial. Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy, 2016, 9, 55.	2.4	8
108	The Time Course of Markers of Neutrophil Extracellular Traps in Patients Undergoing Revascularisation for Acute Myocardial Infarction or Stable Angina Pectoris. Mediators of Inflammation, 2016, 2016, 1-8.	3.0	30



#	ARTICLE	IF	CITATIONS
109	Serum Fatty Acids, Traditional Risk Factors, and Comorbidity as Related to Myocardial Injury in an Elderly Population with Acute Myocardial Infarction. <i>Journal of Lipids</i> , 2016, 2016, 1-7.	4.8	11
110	Combined Elevated Levels of the Proinflammatory Cytokines IL-18 and IL-12 Are Associated with Clinical Events in Patients with Coronary Artery Disease: An Observational Study. <i>Metabolic Syndrome and Related Disorders</i> , 2016, 14, 242-248.	1.3	21
111	Circulating Levels of IL-6 Receptor and gp130 and Long-Term Clinical Outcomes in ST-Elevation Myocardial Infarction. <i>Journal of the American Heart Association</i> , 2016, 5, .	3.7	50
112	Associations Between Delirium and Preoperative Cerebrospinal Fluid C-Reactive Protein, Interleukin-6, and Interleukin-6 Receptor in Individuals with Acute Hip Fracture. <i>Journal of the American Geriatrics Society</i> , 2016, 64, 1456-1463.	2.6	70
113	Troponin I levels in permanent atrial fibrillation—impact of rate control and exercise testing. <i>BMC Cardiovascular Disorders</i> , 2016, 16, 79.	1.7	5
114	Effects of exercise training on carotid intima-media thickness in patients with type 2 diabetes and coronary artery disease. Influence of carotid plaques. <i>Cardiovascular Diabetology</i> , 2016, 15, 13.	6.8	50
115	Soluble RAGE and atherosclerosis in youth with type 1 diabetes: a 5-year follow-up study. <i>Cardiovascular Diabetology</i> , 2015, 14, 126.	6.8	35
116	Effects on Serum Fractalkine by Diet and Omega-3 Fatty Acid Intervention: Relation to Clinical Outcome. <i>Mediators of Inflammation</i> , 2015, 2015, 1-6.	3.0	2
117	Disease Activity in Ankylosing Spondylitis and Associations to Markers of Vascular Pathology and Traditional Cardiovascular Disease Risk Factors: A Cross-sectional Study. <i>Journal of Rheumatology</i> , 2015, 42, 645-653.	2.0	33
118	Glucosepane and oxidative markers in skin collagen correlate with intima media thickness and arterial stiffness in long-term type 1 diabetes. <i>Journal of Diabetes and Its Complications</i> , 2015, 29, 407-412.	2.3	16
119	Serum pneumoproteins in tunnel construction workers. <i>International Archives of Occupational and Environmental Health</i> , 2015, 88, 943-951.	2.3	5
120	The advanced glycation end product methylglyoxal-derived hydroimidazolone-1 and early signs of atherosclerosis in childhood diabetes. <i>Diabetes and Vascular Disease Research</i> , 2015, 12, 139-145.	2.0	37
121	Is there a role for monocyte chemoattractant protein-1 in delirium? Novel observations in elderly hip fracture patients. <i>BMC Research Notes</i> , 2015, 8, 186.	1.4	12
122	Effects of exercise training on HbA <sub>1c</sub> and VO <sub>2peak</sub> in patients with type 2 diabetes and coronary artery disease: A randomised clinical trial. <i>Diabetes and Vascular Disease Research</i> , 2015, 12, 325-333.	2.0	38
123	Inflammation in childhood type 1 diabetes; influence of glycemic control. <i>Atherosclerosis</i> , 2015, 238, 33-37.	0.8	38
124	The Effect of Intracoronary Stem Cell Injection on Markers of Leukocyte Activation in Acute Myocardial Infarction. <i>Cardiology Research</i> , 2015, 6, 209-215.	1.1	3
125	Effects of diet and/or n-3 fatty acid supplementation on components of the interleukin-6 trans-signalling system in elderly men. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2015, 75, 646-51.	1.2	0
126	Association of Interleukin 8 and Myocardial Recovery in Patients with ST-Elevation Myocardial Infarction Complicated by Acute Heart Failure. <i>PLoS ONE</i> , 2014, 9, e112359.	2.5	31



#	ARTICLE	IF	CITATIONS
127	The Time Profile of Pentraxin 3 in Patients with Acute ST-Elevation Myocardial Infarction and Stable Angina Pectoris Undergoing Percutaneous Coronary Intervention. <i>Mediators of Inflammation</i> , 2014, 2014, 1-5.	3.0	19
128	The Influence of Autologous Bone Marrow Stem Cell Transplantation on Matrix Metalloproteinases in Patients Treated for Acute ST-Elevation Myocardial Infarction. <i>Mediators of Inflammation</i> , 2014, 2014, 1-9.	3.0	4
129	Frailty indicators and functional status in older patients after colorectal cancer surgery. <i>Journal of Geriatric Oncology</i> , 2014, 5, 26-32.	1.0	84
130	Prothrombin fragment 1+2 in urine as a marker on coagulation activity in patients with suspected pulmonary embolism. <i>Thrombosis Research</i> , 2014, 134, 68-71.	1.7	3
131	Effects of omega 3 supplementation in elderly patients with acute myocardial infarction: design of a prospective randomized placebo controlled study. <i>BMC Geriatrics</i> , 2014, 14, 74.	2.7	29
132	IL-6 signalling in patients with acute ST-elevation myocardial infarction. <i>Results in Immunology</i> , 2014, 4, 8-13.	2.2	54
133	Glycoprotein 130 polymorphism predicts soluble glycoprotein 130 levels. <i>Metabolism: Clinical and Experimental</i> , 2014, 63, 647-653.	3.4	11
134	Fractalkine levels are elevated early after PCI-treated ST-elevation myocardial infarction; no influence of autologous bone marrow derived stem cell injection. <i>Cytokine</i> , 2014, 69, 131-135.	3.2	15
135	The MMP-9 -1562 C/T Polymorphism in the Presence of Metabolic Syndrome Increases the Risk of Clinical Events in Patients with Coronary Artery Disease. <i>PLoS ONE</i> , 2014, 9, e106816.	2.5	30
136	Impact of atrial fibrillation on inflammatory and fibrinolytic variables in the elderly. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2013, 73, 326-333.	1.2	11
137	Levosimendan in acute heart failure following primary percutaneous coronary interventionâ€treated acute STâ€elevation myocardial infarction. Results from the LEAF trial: a randomized, placeboâ€controlled study. <i>European Journal of Heart Failure</i> , 2013, 15, 565-572.	7.1	71
138	High Onâ€Aspirin Platelet Reactivity and Clinical Outcome in Patients With Stable Coronary Artery Disease: Results From ASCET (Aspirin Nonresponsiveness and Clopidogrel Endpoint Trial). <i>Journal of the American Heart Association</i> , 2012, 1, e000703.	3.7	61
139	Effects of similar intakes of marine<i>n</i>-3 fatty acids from enriched food products and fish oil on cardiovascular risk markers in healthy human subjects. <i>British Journal of Nutrition</i> , 2012, 107, 1339-1349.	2.3	23
140	Asymmetric Dimethylarginine Levels are Highly Associated With Atrial Fibrillation in an Elderly Population. <i>Cardiology Research</i> , 2012, 3, 109-115.	1.1	2
141	The L-Arginineâ€Asymmetric Dimethylarginine Ratio Is Strongly Related to the Severity of Chronic Heart Failure. No Effects of Exercise Training. <i>Journal of Cardiac Failure</i> , 2011, 17, 135-142.	1.7	30
142	Inflammatory responses after percutaneous coronary intervention in patients with acute myocardial infarction or stable angina pectoris. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2008, 68, 555-562.	1.2	19
143	Effect of diet or very long chain Â-3 fatty acids on progression of atherosclerosis, evaluated by carotid plaques, intima-media thickness and by pulse wave propagation in elderly men with hypercholesterolaemia. <i>European Journal of Cardiovascular Prevention and Rehabilitation</i> , 2006, 13, 325-333.	2.8	47
144	The Effect of Supplementation With Omega-3 Fatty Acids on Soluble Markers of Endothelial Function in Patients With Coronary Heart Disease. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 1999, 19, 1681-1686.	2.4	160

#	ARTICLE	IF	CITATIONS
145	n-3 fatty acids do not prevent restenosis after coronary angioplasty: results from the CART study. Journal of the American College of Cardiology, 1999, 33, 1619-1626.	2.8	106
146	Effects of Partially Hydrogenated Fish Oil, Partially Hydrogenated Soybean Oil, and Butter on Hemostatic Variables in Men. Arteriosclerosis, Thrombosis, and Vascular Biology, 1996, 16, 375-380.	2.4	45