

# Ingebjörg Seljeflot

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9130466/publications.pdf>

Version: 2024-02-01

146  
papers

3,111  
citations

147786

31  
h-index

206102

48  
g-index

146  
all docs

146  
docs citations

146  
times ranked

4890  
citing authors

#	ARTICLE	IF	CITATIONS
1	Probiotics to HIV-Infected Immunological Nonresponders: Altered Mucosal Immunity and Microbial Diversity Restricted to Ileum. <i>Journal of Acquired Immune Deficiency Syndromes</i> (1999), 2022, 89, 77-86.	2.1	3
2	Human Immunodeficiency Virus-Infected Immunological Nonresponders Have Colon-Restricted Gut Mucosal Immune Dysfunction. <i>Journal of Infectious Diseases</i> , 2022, 225, 661-674.	4.0	16
3	Changes in eicosapentaenoic acid and docosahexaenoic acid and risk of cardiovascular events and atrial fibrillation: A secondary analysis of the OMEMI trial. <i>Journal of Internal Medicine</i> , 2022, 291, 637-647.	6.0	22
4	Complement ratios C3bc/C3 and sC5b-9/C5 do not increase the sensitivity of detecting acute complement activation systemically. <i>Molecular Immunology</i> , 2022, 141, 273-279.	2.2	2
5	Vascular Function in Norwegian Female Elite Runners: A Cross-Sectional, Controlled Study. <i>Sports</i> , 2022, 10, 37.	1.7	5
6	The NLRP3 inflammasome activation in subcutaneous, epicardial and pericardial adipose tissue in patients with coronary heart disease undergoing coronary by-pass surgery. <i>Atherosclerosis Plus</i> , 2022, 48, 47-54.	0.7	2
7	Gene expression of fibrinolytic markers in coronary thrombi. <i>Thrombosis Journal</i> , 2022, 20, 23.	2.1	0
8	Coagulation factors XI and XII as possible targets for anticoagulant therapy. <i>Thrombosis Research</i> , 2022, 214, 53-62.	1.7	12
9	Interleukin-6 inhibition in ST-elevation myocardial infarction: Immune cell profile in the randomised ASSAIL-MI trial. <i>EBioMedicine</i> , 2022, 80, 104013.	6.1	22
10	Exercise-induced change in circulating NT-proBNP could not distinguish between patients with and without coronary artery disease: the CADENCE study. <i>Scandinavian Cardiovascular Journal</i> , 2022, 56, 107-113.	1.2	0
11	Reduced leukocyte telomere lengths and sirtuin-1 gene expression in long-term survivors of type-1 diabetes: A Dialong substudy. <i>Journal of Diabetes Investigation</i> , 2021, 12, 1183-1192.	2.4	6
12	Effects of n-3 Fatty Acid Supplements in Elderly Patients After Myocardial Infarction. <i>Circulation</i> , 2021, 143, 528-539.	1.6	180
13	Differential associations of cardiac troponin T and cardiac troponin I with coronary artery pathology and dynamics in response to short-duration exercise. <i>Clinical Biochemistry</i> , 2021, 88, 23-29.	1.9	8
14	Biomarkers of ageing and cardiac remodeling are associated with atrial fibrillation. <i>Scandinavian Cardiovascular Journal</i> , 2021, 55, 213-219.	1.2	14
15	Randomized Trial of Interleukin-6 Receptor Inhibition in Patients With Acute ST-Segment Elevation Myocardial Infarction. <i>Journal of the American College of Cardiology</i> , 2021, 77, 1845-1855.	2.8	169
16	Gut related inflammation and cardiorespiratory fitness in patients with CAD and type 2 diabetes: a sub-study of a randomized controlled trial on exercise training. <i>Diabetology and Metabolic Syndrome</i> , 2021, 13, 36.	2.7	4
17	Effects of intermittent negative pressure treatment on circulating vascular biomarkers in patients with intermittent claudication. <i>Vascular Medicine</i> , 2021, 26, 489-496.	1.5	2
18	The Inflammasome Signaling Pathway Is Actively Regulated and Related to Myocardial Damage in Coronary Thrombi from Patients with STEMI. <i>Mediators of Inflammation</i> , 2021, 2021, 1-12.	3.0	7

#	ARTICLE	IF	CITATIONS
19	Tocilizumab increases citrullinated histone 3 in non-ST segment elevation myocardial infarction. <i>Open Heart</i> , 2021, 8, e001492.	2.3	4
20	Mortality and microbial diversity after allogeneic hematopoietic stem cell transplantation: secondary analysis of a randomized nutritional intervention trial. <i>Scientific Reports</i> , 2021, 11, 11593.	3.3	9
21	Transient Reduction of FMD-Response and L-Arginine Accompanied by Increased Levels of E-Selectin, VCAM, and ICAM after Prolonged Strenuous Exercise. <i>Sports</i> , 2021, 9, 86.	1.7	2
22	Effect of intermittent and continuous caloric restriction on Sirtuin1 concentration depends on sex and body mass index. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2021, 31, 1871-1878.	2.6	10
23	Reduced L-Arginine and L-Arginine-ADMA-Ratio, and Increased SDMA after Norseman Xtreme Triathlon. <i>Sports</i> , 2021, 9, 120.	1.7	7
24	Rifaximin or <i>Saccharomyces boulardii</i> in heart failure with reduced ejection fraction: Results from the randomized GutHeart trial. <i>EBioMedicine</i> , 2021, 70, 103511.	6.1	34
25	Gut Leakage Markers in Response to Strenuous Exercise in Patients with Suspected Coronary Artery Disease. <i>Cells</i> , 2021, 10, 2193.	4.1	6
26	Immune complexes, innate immunity, and NETosis in ChAdOx1 vaccine-induced thrombocytopenia. <i>European Heart Journal</i> , 2021, 42, 4064-4072.	2.2	49
27	Complement activation is associated with poor outcome after out-of-hospital cardiac arrest. <i>Resuscitation</i> , 2021, 166, 129-136.	3.0	12
28	Serum Galectin-3 and Subsequent Risk of Coronary Heart Disease in Subjects With Childhood-Onset Type 1 Diabetes: A Cohort Study. <i>Diabetes Care</i> , 2021, 44, 810-816.	8.6	9
29	Serum Levels of Dihomo-Gamma ( $\Gamma^3$ )-Linolenic Acid (DGLA) Are Inversely Associated with Linoleic Acid and Total Death in Elderly Patients with a Recent Myocardial Infarction. <i>Nutrients</i> , 2021, 13, 3475.	4.1	9
30	One year of omega 3 polyunsaturated fatty acid supplementation does not reduce circulating prothrombotic microvesicles in elderly subjects after suffering a myocardial infarction. <i>Clinical Nutrition</i> , 2021, 40, 5674-5677.	5.0	5
31	Adiponectin in relation to exercise and physical performance in patients with type 2 diabetes and coronary artery disease. <i>Adipocyte</i> , 2021, 10, 612-620.	2.8	2
32	NETosis in Long-Term Type 1 Diabetes Mellitus and Its Link to Coronary Artery Disease. <i>Frontiers in Immunology</i> , 2021, 12, 799539.	4.8	4
33	High levels of interleukin-6 are associated with final infarct size and adverse clinical events in patients with STEMI. <i>Open Heart</i> , 2021, 8, e001869.	2.3	10
34	Soluble ST2 concentrations associate with in-hospital mortality and need for mechanical ventilation in unselected patients with COVID-19. <i>Open Heart</i> , 2021, 8, e001884.	2.3	9
35	Late awakening, prognostic factors and long-term outcome in out-of-hospital cardiac arrest – results of the prospective Norwegian Cardio-Respiratory Arrest Study (NORCAST). <i>Resuscitation</i> , 2020, 149, 170-179.	3.0	47
36	Cardiovascular Remodelling in living kidney donorS with reduced glomerular filtration rate: rationale and design of the CENS study. <i>Blood Pressure</i> , 2020, 29, 123-134.	1.5	2

#	ARTICLE	IF	CITATIONS
37	Legumain is upregulated in acute cardiovascular events and associated with improved outcome - potentially related to anti-inflammatory effects on macrophages. <i>Atherosclerosis</i> , 2020, 296, 74-82.	0.8	14
38	Shorter Leukocyte Telomere Lengths in Healthy Relatives of Patients with Coronary Heart Disease. <i>Rejuvenation Research</i> , 2020, 23, 324-332.	1.8	9
39	Superiority of high sensitivity cardiac troponin T vs. I for long-term prognostic value in patients with chest pain; data from the Akershus cardiac Examination (ACE) 3 study. <i>Clinical Biochemistry</i> , 2020, 78, 10-17.	1.9	15
40	Very Long Chain Marine n-3 Polyunsaturated Fatty Acids in Atherothrombotic Heart Disease. A Brief Review, with a Focus on Metabolic Effects. <i>Nutrients</i> , 2020, 12, 3014.	4.1	4
41	Left ventricular dysfunction in COPD without pulmonary hypertension. <i>PLoS ONE</i> , 2020, 15, e0235075.	2.5	3
42	Circulating levels of the terminal complement complex are associated with hypercoagulability in patients with stable coronary artery disease. <i>Thrombosis Research</i> , 2020, 196, 106-108.	1.7	1
43	Markers of remodeling in subcutaneous adipose tissue are strongly associated with overweight and insulin sensitivity in healthy non-obese men. <i>Scientific Reports</i> , 2020, 10, 14055.	3.3	8
44	Collagen methionine sulfoxide and glucuronidine/LW-1 are markers of coronary artery disease in long-term survivors with type 1 diabetes. The Dialong study. <i>PLoS ONE</i> , 2020, 15, e0233174.	2.5	8
45	Double-Stranded DNA and NETs Components in Relation to Clinical Outcome After ST-Elevation Myocardial Infarction. <i>Scientific Reports</i> , 2020, 10, 5007.	3.3	22
46	Low fibre intake is associated with gut microbiota alterations in chronic heart failure. <i>ESC Heart Failure</i> , 2020, 7, 456-466.	3.1	56
47	Procoagulant activity in children and adolescents on intensive insulin therapy. <i>Pediatric Diabetes</i> , 2020, 21, 496-504.	2.9	2
48	Rosuvastatin alters the genetic composition of the human gut microbiome. <i>Scientific Reports</i> , 2020, 10, 5397.	3.3	20
49	Complement Activation in Association with Markers of Neutrophil Extracellular Traps and Acute Myocardial Infarction in Stable Coronary Artery Disease. <i>Mediators of Inflammation</i> , 2020, 2020, 1-9.	3.0	8
50	Annexin V+ Microvesicles in Children and Adolescents with Type 1 Diabetes: A Prospective Cohort Study. <i>Journal of Diabetes Research</i> , 2020, 2020, 1-8.	2.3	2
51	Neutrophil extracellular trap components and myocardial recovery in post-ischemic acute heart failure. <i>PLoS ONE</i> , 2020, 15, e0241333.	2.5	6
52	Left ventricular dysfunction in COPD without pulmonary hypertension. , 2020, 15, e0235075.		0
53	Left ventricular dysfunction in COPD without pulmonary hypertension. , 2020, 15, e0235075.		0
54	Left ventricular dysfunction in COPD without pulmonary hypertension. , 2020, 15, e0235075.		0

#	ARTICLE	IF	CITATIONS
55	Left ventricular dysfunction in COPD without pulmonary hypertension. , 2020, 15, e0235075.		0
56	Title is missing!. , 2020, 15, e0233174.		0
57	Title is missing!. , 2020, 15, e0233174.		0
58	Title is missing!. , 2020, 15, e0233174.		0
59	Title is missing!. , 2020, 15, e0233174.		0
60	Title is missing!. , 2020, 15, e0233174.		0
61	Title is missing!. , 2020, 15, e0233174.		0
62	Glucose associated NETosis in patients with ST-elevation myocardial infarction: an observational study. BMC Cardiovascular Disorders, 2019, 19, 221.	1.7	9
63	Novel biomolecules of ageing, sex differences and potential underlying mechanisms of telomere shortening in coronary artery disease. Experimental Gerontology, 2019, 119, 53-60.	2.8	23
64	A study of atherothrombotic biomarkers in welders. International Archives of Occupational and Environmental Health, 2019, 92, 1023-1031.	2.3	8
65	High Adherence to the Nordic Diet Is Associated with Lower Levels of Total and Platelet-Derived Circulating Microvesicles in a Norwegian Population. Nutrients, 2019, 11, 1114.	4.1	7
66	A brief review on resistance to P2Y12 receptor antagonism in coronary artery disease. Thrombosis Journal, 2019, 17, 11.	2.1	46
67	Effects of dietary intervention and n-3 PUFA supplementation on markers of gut-related inflammation and their association with cardiovascular events in a high-risk population. Atherosclerosis, 2019, 286, 53-59.	0.8	16
68	IgM antibodies against phosphorylcholine measured early after acute ST-elevation myocardial infarction in relation to atherosclerotic disease burden and long-term clinical outcome. PLoS ONE, 2019, 14, e0215640.	2.5	1
69	High-sensitive cardiac Troponin T and exercise stress test for evaluation of angiographically significant coronary disease. International Journal of Cardiology, 2019, 287, 1-6.	1.7	8
70	Elevated levels of circulating microvesicles in coronary artery disease patients with type 2 diabetes and albuminuria: Effects of exercise training. Diabetes and Vascular Disease Research, 2019, 16, 431-439.	2.0	10
71	Effects of exercise training on inflammasome-related mediators and their associations to glucometabolic variables in patients with combined coronary artery disease and type 2 diabetes mellitus: Sub-study of a randomized control trial. Diabetes and Vascular Disease Research, 2019, 16, 360-368.	2.0	14
72	Index of microvascular resistance to assess the effect of rosuvastatin on microvascular function in women with chest pain and no obstructive coronary artery disease: A double-blind randomized study. Catheterization and Cardiovascular Interventions, 2019, 94, 660-668.	1.7	10

#	ARTICLE	IF	CITATIONS
73	Rationale for the ASSAIL-MI-trial: a randomised controlled trial designed to assess the effect of tocilizumab on myocardial salvage in patients with acute ST-elevation myocardial infarction (STEMI). <i>Open Heart</i> , 2019, 6, e001108.	2.3	34
74	Effects of exercise training on markers of adipose tissue remodeling in patients with coronary artery disease and type 2 diabetes mellitus: sub study of the randomized controlled EXCADI trial. <i>Diabetology and Metabolic Syndrome</i> , 2019, 11, 109.	2.7	3
75	Neutrophil Extracellular Trap Components Associate with Infarct Size, Ventricular Function, and Clinical Outcome in STEMI. <i>Mediators of Inflammation</i> , 2019, 2019, 1-10.	3.0	33
76	Leukocyte telomere length and serum polyunsaturated fatty acids, dietary habits, cardiovascular risk factors and features of myocardial infarction in elderly patients. <i>BMC Geriatrics</i> , 2019, 19, 376.	2.7	10
77	Circulating markers of gut barrier function associated with disease severity in primary sclerosing cholangitis. <i>Liver International</i> , 2019, 39, 371-381.	3.9	51
78	Markers of neutrophil extracellular traps are associated with adverse clinical outcome in stable coronary artery disease. <i>European Journal of Preventive Cardiology</i> , 2018, 25, 762-769.	1.8	34
79	Antithrombotic therapy and body mass: an expert position paper of the ESC Working Group on Thrombosis. <i>European Heart Journal</i> , 2018, 39, 1672-1686f.	2.2	106
80	Effect of strenuous exercise on mediators of inflammation in patients with coronary artery disease. <i>Cytokine</i> , 2018, 105, 17-22.	3.2	13
81	Acute heart failure following myocardial infarction: complement activation correlates with the severity of heart failure in patients developing cardiogenic shock. <i>ESC Heart Failure</i> , 2018, 5, 292-301.	3.1	27
82	Impact of HIV and Type 2 diabetes on Gut Microbiota Diversity, Tryptophan Catabolism and Endothelial Dysfunction. <i>Scientific Reports</i> , 2018, 8, 6725.	3.3	35
83	Markers of Thrombin Generation Are Associated With Long-Term Clinical Outcome in Patients With ST-Segment Elevation Myocardial Infarction. <i>Clinical and Applied Thrombosis/Hemostasis</i> , 2018, 24, 1088-1094.	1.7	11
84	Preserved endothelial function in young adults with type 1 diabetes. <i>PLoS ONE</i> , 2018, 13, e0206523.	2.5	13
85	Soluble IL-1 receptor 2 is associated with left ventricular remodelling in patients with ST-elevation myocardial infarction. <i>International Journal of Cardiology</i> , 2018, 268, 187-192.	1.7	15
86	Association of IL-8 With Infarct Size and Clinical Outcomes in Patients With STEMI. <i>Journal of the American College of Cardiology</i> , 2018, 72, 187-198.	2.8	40
87	High-sensitivity troponin I in persistent atrial fibrillation â€” relation to NT-proBNP and markers of inflammation and haemostasis. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2018, 78, 386-392.	1.2	2
88	Markers of metabolic endotoxemia as related to metabolic syndrome in an elderly male population at high cardiovascular risk: a cross-sectional study. <i>Diabetology and Metabolic Syndrome</i> , 2018, 10, 59.	2.7	35
89	Interleukin-18 and the NLR family pyrin domain containing-3 inflammasome in adipose tissue are strongly associated with glucometabolic variables in a cohort of middle-aged men. <i>Diabetes and Vascular Disease Research</i> , 2018, 15, 458-464.	2.0	4
90	Design of the GutHeartâ€”targeting gut microbiota to treat heart failureâ€”trial: a Phase II, randomized clinical trial. <i>ESC Heart Failure</i> , 2018, 5, 977-984.	3.1	39

#	ARTICLE	IF	CITATIONS
91	Increased arterial stiffness in childhood onset diabetes: a cardiovascular magnetic resonance study. <i>European Heart Journal Cardiovascular Imaging</i> , 2018, 19, 694-700.	1.2	12
92	Biomarkers of endothelial activation and thrombosis in tunnel construction workers exposed to airborne contaminants. <i>International Archives of Occupational and Environmental Health</i> , 2017, 90, 309-317.	2.3	5
93	Reduced endothelial activation after exercise is associated with improved HbA <sub>1c</sub> in patients with type 2 diabetes and coronary artery disease. <i>Diabetes and Vascular Disease Research</i> , 2017, 14, 94-103.	2.0	11
94	Procoagulant activity in patients with combined type 2 diabetes and coronary artery disease: No effects of long-term exercise training. <i>Diabetes and Vascular Disease Research</i> , 2017, 14, 144-151.	2.0	6
95	Changes in dietary pattern when including 700g of salmon per week to patients with atherosclerotic heart disease. <i>Clinical Nutrition ESPEN</i> , 2017, 19, 38-44.	1.2	2
96	Thrombin Generation in Patients With Suspected Venous Thromboembolism. <i>Clinical and Applied Thrombosis/Hemostasis</i> , 2017, 23, 416-421.	1.7	10
97	HIV-infected persons with type 2 diabetes show evidence of endothelial dysfunction and increased inflammation. <i>BMC Infectious Diseases</i> , 2017, 17, 234.	2.9	19
98	Monocyte-derived circulating microparticles (CD14+, CD14+/CD11b+ and CD14+/CD142+) are related to long-term prognosis for cardiovascular mortality in STEMI patients. <i>International Journal of Cardiology</i> , 2017, 227, 876-881.	1.7	47
99	Evaluation of circulating levels of CCN2/connective tissue growth factor in patients with ST-elevation myocardial infarction. <i>Scientific Reports</i> , 2017, 7, 11945.	3.3	4
100	Pro-coagulant activity during exercise testing in patients with coronary artery disease. <i>Thrombosis Journal</i> , 2017, 15, 3.	2.1	9
101	A Double-Blinded Randomized Study Investigating a Possible Anti-Inflammatory Effect of Saxagliptin versus Placebo as Add-On Therapy in Patients with Both Type 2 Diabetes And Stable Coronary Artery Disease. <i>Mediators of Inflammation</i> , 2017, 2017, 1-9.	3.0	2
102	Osteoprotegerin levels in ST-elevation myocardial infarction: Temporal profile and association with myocardial injury and left ventricular function. <i>PLoS ONE</i> , 2017, 12, e0173034.	2.5	8
103	Reduced HDL function in children and young adults with type 1 diabetes. <i>Cardiovascular Diabetology</i> , 2017, 16, 85.	6.8	30
104	vWF/ADAMTS13 is associated with on-aspirin residual platelet reactivity and clinical outcome in patients with stable coronary artery disease. <i>Thrombosis Journal</i> , 2017, 15, 28.	2.1	12
105	Effect of sinus rhythm restoration on markers of thrombin generation in atrial fibrillation. <i>Thrombosis Journal</i> , 2017, 15, 30.	2.1	2
106	Platelet-, monocyte-derived and tissue factor-carrying circulating microparticles are related to acute myocardial infarction severity. <i>PLoS ONE</i> , 2017, 12, e0172558.	2.5	74
107	Effects of long-term exercise training on adipose tissue expression of fractalkine and MCP-1 in patients with type 2 diabetes and stable coronary artery disease: a substudy of a randomized controlled trial. <i>Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy</i> , 2016, 9, 55.	2.4	8
108	The Time Course of Markers of Neutrophil Extracellular Traps in Patients Undergoing Revascularisation for Acute Myocardial Infarction or Stable Angina Pectoris. <i>Mediators of Inflammation</i> , 2016, 2016, 1-8.	3.0	30

#	ARTICLE	IF	CITATIONS
109	Serum Fatty Acids, Traditional Risk Factors, and Comorbidity as Related to Myocardial Injury in an Elderly Population with Acute Myocardial Infarction. <i>Journal of Lipids</i> , 2016, 2016, 1-7.	4.8	11
110	Combined Elevated Levels of the Proinflammatory Cytokines IL-18 and IL-12 Are Associated with Clinical Events in Patients with Coronary Artery Disease: An Observational Study. <i>Metabolic Syndrome and Related Disorders</i> , 2016, 14, 242-248.	1.3	21
111	Circulating Levels of IL-6 Receptor and gp130 and Long-Term Clinical Outcomes in ST-Elevation Myocardial Infarction. <i>Journal of the American Heart Association</i> , 2016, 5, .	3.7	50
112	Associations Between Delirium and Preoperative Cerebrospinal Fluid C-Reactive Protein, Interleukin-6, and Interleukin-6 Receptor in Individuals with Acute Hip Fracture. <i>Journal of the American Geriatrics Society</i> , 2016, 64, 1456-1463.	2.6	70
113	Troponin I levels in permanent atrial fibrillation—impact of rate control and exercise testing. <i>BMC Cardiovascular Disorders</i> , 2016, 16, 79.	1.7	5
114	Effects of exercise training on carotid intima-media thickness in patients with type 2 diabetes and coronary artery disease. Influence of carotid plaques. <i>Cardiovascular Diabetology</i> , 2016, 15, 13.	6.8	50
115	Soluble RAGE and atherosclerosis in youth with type 1 diabetes: a 5-year follow-up study. <i>Cardiovascular Diabetology</i> , 2015, 14, 126.	6.8	35
116	Effects on Serum Fractalkine by Diet and Omega-3 Fatty Acid Intervention: Relation to Clinical Outcome. <i>Mediators of Inflammation</i> , 2015, 2015, 1-6.	3.0	2
117	Disease Activity in Ankylosing Spondylitis and Associations to Markers of Vascular Pathology and Traditional Cardiovascular Disease Risk Factors: A Cross-sectional Study. <i>Journal of Rheumatology</i> , 2015, 42, 645-653.	2.0	33
118	Glucosepane and oxidative markers in skin collagen correlate with intima media thickness and arterial stiffness in long-term type 1 diabetes. <i>Journal of Diabetes and Its Complications</i> , 2015, 29, 407-412.	2.3	16
119	Serum pneumoproteins in tunnel construction workers. <i>International Archives of Occupational and Environmental Health</i> , 2015, 88, 943-951.	2.3	5
120	The advanced glycation end product methylglyoxal-derived hydroimidazolone-1 and early signs of atherosclerosis in childhood diabetes. <i>Diabetes and Vascular Disease Research</i> , 2015, 12, 139-145.	2.0	37
121	Is there a role for monocyte chemoattractant protein-1 in delirium? Novel observations in elderly hip fracture patients. <i>BMC Research Notes</i> , 2015, 8, 186.	1.4	12
122	Effects of exercise training on HbA <sub>1c</sub> and VO <sub>2peak</sub> in patients with type 2 diabetes and coronary artery disease: A randomised clinical trial. <i>Diabetes and Vascular Disease Research</i> , 2015, 12, 325-333.	2.0	38
123	Inflammation in childhood type 1 diabetes; influence of glycemic control. <i>Atherosclerosis</i> , 2015, 238, 33-37.	0.8	38
124	The Effect of Intracoronary Stem Cell Injection on Markers of Leukocyte Activation in Acute Myocardial Infarction. <i>Cardiology Research</i> , 2015, 6, 209-215.	1.1	3
125	Effects of diet and/or n-3 fatty acid supplementation on components of the interleukin-6 trans-signalling system in elderly men. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2015, 75, 646-51.	1.2	0
126	Association of Interleukin 8 and Myocardial Recovery in Patients with ST-Elevation Myocardial Infarction Complicated by Acute Heart Failure. <i>PLoS ONE</i> , 2014, 9, e112359.	2.5	31



#	ARTICLE	IF	CITATIONS
127	The Time Profile of Pentraxin 3 in Patients with Acute ST-Elevation Myocardial Infarction and Stable Angina Pectoris Undergoing Percutaneous Coronary Intervention. <i>Mediators of Inflammation</i> , 2014, 2014, 1-5.	3.0	19
128	The Influence of Autologous Bone Marrow Stem Cell Transplantation on Matrix Metalloproteinases in Patients Treated for Acute ST-Elevation Myocardial Infarction. <i>Mediators of Inflammation</i> , 2014, 2014, 1-9.	3.0	4
129	Frailty indicators and functional status in older patients after colorectal cancer surgery. <i>Journal of Geriatric Oncology</i> , 2014, 5, 26-32.	1.0	84
130	Prothrombin fragment 1+2 in urine as a marker on coagulation activity in patients with suspected pulmonary embolism. <i>Thrombosis Research</i> , 2014, 134, 68-71.	1.7	3
131	Effects of omega 3 supplementation in elderly patients with acute myocardial infarction: design of a prospective randomized placebo controlled study. <i>BMC Geriatrics</i> , 2014, 14, 74.	2.7	29
132	IL-6 signalling in patients with acute ST-elevation myocardial infarction. <i>Results in Immunology</i> , 2014, 4, 8-13.	2.2	54
133	Glycoprotein 130 polymorphism predicts soluble glycoprotein 130 levels. <i>Metabolism: Clinical and Experimental</i> , 2014, 63, 647-653.	3.4	11
134	Fractalkine levels are elevated early after PCI-treated ST-elevation myocardial infarction; no influence of autologous bone marrow derived stem cell injection. <i>Cytokine</i> , 2014, 69, 131-135.	3.2	15
135	The MMP-9 -1562 C/T Polymorphism in the Presence of Metabolic Syndrome Increases the Risk of Clinical Events in Patients with Coronary Artery Disease. <i>PLoS ONE</i> , 2014, 9, e106816.	2.5	30
136	Impact of atrial fibrillation on inflammatory and fibrinolytic variables in the elderly. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2013, 73, 326-333.	1.2	11
137	Levosimendan in acute heart failure following primary percutaneous coronary intervention—treated acute ST-elevation myocardial infarction. Results from the LEAF trial: a randomized, placebo-controlled study. <i>European Journal of Heart Failure</i> , 2013, 15, 565-572.	7.1	71
138	High On-Aspirin Platelet Reactivity and Clinical Outcome in Patients With Stable Coronary Artery Disease: Results From ASCET (Aspirin Nonresponsiveness and Clopidogrel Endpoint Trial). <i>Journal of the American Heart Association</i> , 2012, 1, e000703.	3.7	61
139	Effects of similar intakes of marine $\omega$ -3 fatty acids from enriched food products and fish oil on cardiovascular risk markers in healthy human subjects. <i>British Journal of Nutrition</i> , 2012, 107, 1339-1349.	2.3	23
140	Asymmetric Dimethylarginine Levels are Highly Associated With Atrial Fibrillation in an Elderly Population. <i>Cardiology Research</i> , 2012, 3, 109-115.	1.1	2
141	The L-Arginine/Asymmetric Dimethylarginine Ratio Is Strongly Related to the Severity of Chronic Heart Failure. No Effects of Exercise Training. <i>Journal of Cardiac Failure</i> , 2011, 17, 135-142.	1.7	30
142	Inflammatory responses after percutaneous coronary intervention in patients with acute myocardial infarction or stable angina pectoris. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2008, 68, 555-562.	1.2	19
143	Effect of diet or very long chain $\omega$ -3 fatty acids on progression of atherosclerosis, evaluated by carotid plaques, intima-media thickness and by pulse wave propagation in elderly men with hypercholesterolaemia. <i>European Journal of Cardiovascular Prevention and Rehabilitation</i> , 2006, 13, 325-333.	2.8	47
144	The Effect of Supplementation With Omega-3 Fatty Acids on Soluble Markers of Endothelial Function in Patients With Coronary Heart Disease. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 1999, 19, 1681-1686.	2.4	160

#	ARTICLE	IF	CITATIONS
145	n-3 fatty acids do not prevent restenosis after coronary angioplasty: results from the CART study. Journal of the American College of Cardiology, 1999, 33, 1619-1626.	2.8	106
146	Effects of Partially Hydrogenated Fish Oil, Partially Hydrogenated Soybean Oil, and Butter on Hemostatic Variables in Men. Arteriosclerosis, Thrombosis, and Vascular Biology, 1996, 16, 375-380.	2.4	45