

# Mihael Vucur

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9133712/publications.pdf>

Version: 2024-02-01

49  
papers

3,445  
citations

218592

26  
h-index

197736

49  
g-index

50  
all docs

50  
docs citations

50  
times ranked

6012  
citing authors

#	ARTICLE	IF	CITATIONS
1	Circulating levels of microRNA193a-5p predict outcome in early stage hepatocellular carcinoma. PLoS ONE, 2020, 15, e0239386.	1.1	11
2	Analysis of miR-29 Serum Levels in Patients with Neuroendocrine Tumors—Results from an Exploratory Study. Journal of Clinical Medicine, 2020, 9, 2881.	1.0	8
3	Skeletal Muscle Composition Predicts Outcome in Critically Ill Patients. , 2020, 2, e0171.		34
4	Circulating levels of soluble urokinase plasminogen activator receptor predict outcome after resection of biliary tract cancer. JHEP Reports, 2020, 2, 100080.	2.6	17
5	Life is fragile: FMRP controls cell death in liver disease. Gut, 2020, 69, 2-3.	6.1	2
6	Serum levels of bone sialoprotein correlate with portal pressure in patients with liver cirrhosis. PLoS ONE, 2020, 15, e0231701.	1.1	4
7	An NF-kappaB- and IKK-Independent Function of NEMO Prevents Hepatocarcinogenesis by Suppressing Compensatory Liver Regeneration. Cancers, 2019, 11, 999.	1.7	13
8	A Combined Score of Circulating miRNAs Allows Outcome Prediction in Critically Ill Patients. Journal of Clinical Medicine, 2019, 8, 1644.	1.0	6
9	Serum levels of miR-29, miR-122, miR-155 and miR-192 are elevated in patients with cholangiocarcinoma. PLoS ONE, 2019, 14, e0210944.	1.1	43
10	Perilipin 5 and Lipocalin 2 Expression in Hepatocellular Carcinoma. Cancers, 2019, 11, 385.	1.7	25
11	miR-155 Predicts Long-Term Mortality in Critically Ill Patients Younger than 65 Years. Mediators of Inflammation, 2019, 2019, 1-8.	1.4	12
12	Serum Levels of Kisspeptin Are Elevated in Patients with Pancreatic Cancer. Disease Markers, 2019, 2019, 1-8.	0.6	7
13	Serum Levels of miR-143 Predict Survival in Critically Ill Patients. Disease Markers, 2019, 2019, 1-10.	0.6	10
14	Serum levels of kisspeptin are elevated in critically ill patients. PLoS ONE, 2018, 13, e0206064.	1.1	8
15	Necroptosis microenvironment directs lineage commitment in liver cancer. Nature, 2018, 562, 69-75.	13.7	283
16	Circulating Biomarkers for Cholangiocarcinoma. Digestive Diseases, 2018, 36, 281-288.	0.8	18
17	Elevated Serum Levels of Mixed Lineage Kinase Domain-Like Protein Predict Survival of Patients during Intensive Care Unit Treatment. Disease Markers, 2018, 2018, 1-8.	0.6	16
18	Differential Roles of Tumor Necrosis Factor Ligand Superfamily Members as Biomarkers in Pancreatic Cancer. Journal of Clinical Medicine, 2018, 7, 175.	1.0	5

#	ARTICLE	IF	CITATIONS
19	Elevated serum levels of bone sialoprotein during ICU treatment predict long-term mortality in critically ill patients. <i>Scientific Reports</i> , 2018, 8, 9750.	1.6	3
20	The Role of miRNAs in the Pathophysiology of Liver Diseases and Toxicity. <i>International Journal of Molecular Sciences</i> , 2018, 19, 261.	1.8	96
21	IL-6 and IL-8 Serum Levels Predict Tumor Response and Overall Survival after TACE for Primary and Secondary Hepatic Malignancies. <i>International Journal of Molecular Sciences</i> , 2018, 19, 1766.	1.8	38
22	Serum levels of soluble urokinase plasminogen activator receptor (suPAR) predict outcome after resection of colorectal liver metastases. <i>Oncotarget</i> , 2018, 9, 27027-27038.	0.8	19
23	The enigma of RIPK1 in the liver: More than just a kinase. <i>Molecular and Cellular Oncology</i> , 2017, 4, e1304191.	0.3	10
24	RIPK1 Suppresses a TRAF2-Dependent Pathway to Liver Cancer. <i>Cancer Cell</i> , 2017, 31, 94-109.	7.7	115
25	A Dual Role of Caspase-8 in Triggering and Sensing Proliferation-Associated DNA Damage, a Key Determinant of Liver Cancer Development. <i>Cancer Cell</i> , 2017, 32, 342-359.e10.	7.7	122
26	CEA but not CA19-9 is an independent prognostic factor in patients undergoing resection of cholangiocarcinoma. <i>Scientific Reports</i> , 2017, 7, 16975.	1.6	65
27	miR-1224 inhibits cell proliferation in acute liver failure by targeting the antiapoptotic gene Nfib. <i>Journal of Hepatology</i> , 2017, 67, 966-978.	1.8	64
28	miR-223 represents a biomarker in acute and chronic liver injury. <i>Clinical Science</i> , 2017, 131, 1971-1987.	1.8	35
29	Elevated levels of circulating osteopontin are associated with a poor survival after resection of cholangiocarcinoma. <i>Journal of Hepatology</i> , 2017, 67, 749-757.	1.8	64
30	Elevated Omentin Serum Levels Predict Long-Term Survival in Critically Ill Patients. <i>Disease Markers</i> , 2016, 2016, 1-9.	0.6	12
31	Receptor interacting protein kinase 1 (RIPK1) in hepatocytes does not mediate murine acetaminophen toxicity. <i>Hepatology</i> , 2016, 64, 306-308.	3.6	26
32	Serum levels of S100A6 are unaltered in patients with resectable cholangiocarcinoma. <i>Clinical and Translational Medicine</i> , 2016, 5, 39.	1.7	14
33	Down-regulation of miR-192-5p protects from oxidative stress-induced acute liver injury. <i>Clinical Science</i> , 2016, 130, 1197-1207.	1.8	59
34	Î² kinase/Î² control biliary homeostasis and hepatocarcinogenesis in mice by phosphorylating the cell death mediator receptor-interacting protein kinase 1. <i>Hepatology</i> , 2016, 64, 1217-1231.	3.6	54
35	The necroptosis-inducing kinase RIPK3 dampens adipose tissue inflammation and glucose intolerance. <i>Nature Communications</i> , 2016, 7, 11869.	5.8	68
36	Histidine-rich glycoprotein promotes macrophage activation and inflammation in chronic liver disease. <i>Hepatology</i> , 2016, 63, 1310-1324.	3.6	77

#	ARTICLE	IF	CITATIONS
37	Necroptosis in Nonalcoholic Steatohepatitis. <i>Cellular and Molecular Gastroenterology and Hepatology</i> , 2015, 1, 264-265.	2.3	25
38	<scp>miR</scp>â€30c and <scp>miR</scp>â€193 are a part of the <scp>TGF</scp>â€â€dependent regulatory network controlling extracellular matrix genes in liver fibrosis. <i>Journal of Digestive Diseases</i> , 2015, 16, 513-524.	0.7	57
39	Elevated miRâ€122 serum levels are an independent marker of liver injury in inflammatory diseases. <i>Liver International</i> , 2015, 35, 1172-1184.	1.9	98
40	A positive feedback loop between <scp>RIP</scp>3 and <scp>JNK</scp> controls nonâ€alcoholic steatohepatitis. <i>EMBO Molecular Medicine</i> , 2014, 6, 1062-1074.	3.3	253
41	RIP3, a kinase promoting necroptotic cell death, mediates adverse remodelling after myocardial infarction. <i>Cardiovascular Research</i> , 2014, 103, 206-216.	1.8	257
42	Levels of Circulating miR-133a Are Elevated in Sepsis and Predict Mortality in Critically Ill Patients. <i>Critical Care Medicine</i> , 2014, 42, 1096-1104.	0.4	111
43	RIP3 Inhibits Inflammatory Hepatocarcinogenesis but Promotes Cholestasis by Controlling Caspase-8- and JNK-Dependent Compensatory Cell Proliferation. <i>Cell Reports</i> , 2013, 4, 776-790.	2.9	124
44	Serum concentrations of A Proliferation-Inducing Ligand (APRIL) are elevated in sepsis and predict mortality in critically ill patients. <i>Journal of Critical Care</i> , 2013, 28, 882.e1-882.e11.	1.0	10
45	U6 is unsuitable for normalization of serum miRNA levels in patients with sepsis or liver fibrosis. <i>Experimental and Molecular Medicine</i> , 2013, 45, e42-e42.	3.2	139
46	Circulating MicroRNA-150 Serum Levels Predict Survival in Patients with Critical Illness and Sepsis. <i>PLoS ONE</i> , 2013, 8, e54612.	1.1	138
47	Micro-RNA profiling reveals a role for miR-29 in human and murine liver fibrosis. <i>Hepatology</i> , 2011, 53, 209-218.	3.6	696
48	Mouse models of hepatocarcinogenesis: What can we learn for the prevention of human hepatocellular carcinoma?. <i>Oncotarget</i> , 2010, 1, 373-378.	0.8	43
49	Mouse models of hepatocarcinogenesis: what can we learn for the prevention of human hepatocellular carcinoma?. <i>Oncotarget</i> , 2010, 1, 373-8.	0.8	28