

# Helene L Soberg

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/913507/publications.pdf>

Version: 2024-02-01

48  
papers

1,133  
citations

394421

19  
h-index

414414

32  
g-index

51  
all docs

51  
docs citations

51  
times ranked

1389  
citing authors

#	ARTICLE	IF	CITATIONS
1	Rehabilitation interventions after traumatic brain injury: a scoping review. <i>Disability and Rehabilitation</i> , 2022, 44, 653-660.	1.8	17
2	Patient-Reported Problem Areas in Chronic Traumatic Brain Injury. <i>Journal of Head Trauma Rehabilitation</i> , 2022, 37, E336-E345.	1.7	8
3	Cost-effectiveness analysis of combined cognitive and vocational rehabilitation in patients with mild-to-moderate TBI: results from a randomized controlled trial. <i>BMC Health Services Research</i> , 2022, 22, 185.	2.2	6
4	Goal Attainment in an Individually Tailored and Home-Based Intervention in the Chronic Phase after Traumatic Brain Injury. <i>Journal of Clinical Medicine</i> , 2022, 11, 958.	2.4	5
5	Psychometric properties of the Norwegian foot function index revised short form. <i>BMC Musculoskeletal Disorders</i> , 2022, 23, 416.	1.9	1
6	What are the effects of different modes of exercise training for intermittent claudication? - A Cochrane Review summary with commentary. <i>Journal of Rehabilitation Medicine</i> , 2021, 53, jrm00181.	1.1	4
7	Rehabilitation Needs, Service Provision, and Costs in the First Year Following Traumatic Injuries: Protocol for a Prospective Cohort Study. <i>JMIR Research Protocols</i> , 2021, 10, e25980.	1.0	5
8	The effectiveness of a family-centred intervention after traumatic brain injury: A pragmatic randomised controlled trial. <i>Clinical Rehabilitation</i> , 2021, 35, 026921552110103.	2.2	9
9	Rehabilitation Goals and Effects of Goal Achievement on Outcome Following an Adapted Physical Activity-Based Rehabilitation Intervention. <i>Patient Preference and Adherence</i> , 2021, Volume 15, 1545-1555.	1.8	6
10	Patient-specific functioning related to dizziness and balance problems after traumatic brain injury – A cross sectional study using an ICF perspective. <i>Cogent Medicine</i> , 2021, 8, .	0.7	2
11	Associations between shoulder pain and functioning on the ICF checklist and the disabilities of the arm, shoulder, and hand scale – a cross-sectional study. <i>Disability and Rehabilitation</i> , 2020, 42, 3084-3091.	1.8	2
12	Family needs at one and two years after severe traumatic brain injury: a prospective study of changes and predictors. <i>Brain Injury</i> , 2020, 34, 89-97.	1.2	11
13	Effectiveness of Combining Compensatory Cognitive Training and Vocational Intervention vs. Treatment as Usual on Return to Work Following Mild-to-Moderate Traumatic Brain Injury: Interim Analysis at 3 and 6 Month Follow-Up. <i>Frontiers in Neurology</i> , 2020, 11, 561400.	2.4	12
14	The Applicability of the Patient-Specific Functional Scale (PSFS) in Rehabilitation for Patients with Acquired Brain Injury (ABI) – A Cohort Study. <i>Journal of Multidisciplinary Healthcare</i> , 2020, Volume 13, 1121-1132.	2.7	8
15	Traumatic brain injury’s needs and treatment options in the chronic phase: Study protocol for a randomized controlled community-based intervention. <i>Trials</i> , 2020, 21, 294.	1.6	15
16	Mental Health and Family Functioning in Patients and Their Family Members after Traumatic Brain Injury: A Cross-Sectional Study. <i>Brain Sciences</i> , 2020, 10, 670.	2.3	10
17	The effects of vestibular rehabilitation on dizziness and balance problems in patients after traumatic brain injury: a randomized controlled trial. <i>Clinical Rehabilitation</i> , 2019, 33, 74-84.	2.2	62
18	The family as a resource for improving patient and family functioning after traumatic brain injury: A descriptive nonrandomized feasibility study of a family-centered intervention. <i>Cogent Medicine</i> , 2019, 6, 1607433.	0.7	2

#	ARTICLE	IF	CITATIONS
19	Measurement properties of the modified and total Balance Error Scoring System – the BESS, in a healthy adult sample. <i>European Journal of Physiotherapy</i> , 2018, 20, 25-31.	1.3	7
20	One year results of a randomized controlled trial on radial Extracorporeal Shock Wave Treatment, with predictors of pain, disability and return to work in patients with subacromial pain syndrome. <i>European Journal of Physical and Rehabilitation Medicine</i> , 2018, 54, 341-350.	2.2	20
21	The Norwegian version of the QOLIBRI – a study of metric properties based on a 12-month follow-up of persons with traumatic brain injury. <i>Health and Quality of Life Outcomes</i> , 2017, 15, 14.	2.4	15
22	Longitudinal changes in brain morphology from 4 weeks to 12 months after mild traumatic brain injury: Associations with cognitive functions and clinical variables. <i>Brain Injury</i> , 2017, 31, 674-685.	1.2	21
23	Effectiveness of Radial Extracorporeal Shock Wave Therapy (rESWT) When Combined With Supervised Exercises in Patients With Subacromial Shoulder Pain: A Double-Masked, Randomized, Sham-Controlled Trial. <i>American Journal of Sports Medicine</i> , 2017, 45, 2547-2554.	4.2	18
24	White matter microstructure is associated with functional, cognitive and emotional symptoms 12 months after mild traumatic brain injury. <i>Scientific Reports</i> , 2017, 7, 13795.	3.3	39
25	Dizziness-related disability following mild-to-moderate traumatic brain injury. <i>Brain Injury</i> , 2017, 31, 1436-1444.	1.2	16
26	Transitions of patients with traumatic brain injury and multiple trauma between specialized and municipal rehabilitation services – Professionals' perspectives. <i>Cogent Medicine</i> , 2017, 4, 1320849.	0.7	7
27	Predicting Outcome 12 Months after Mild Traumatic Brain Injury in Patients Admitted to a Neurosurgery Service. <i>Frontiers in Neurology</i> , 2017, 8, 125.	2.4	25
28	The One or the Many. <i>Qualitative Health Research</i> , 2017, 27, 51-59.	2.1	6
29	Shoulder MRI features with clinical correlations in subacromial pain syndrome: a cross-sectional and prognostic study. <i>BMC Musculoskeletal Disorders</i> , 2017, 18, 469.	1.9	16
30	Combined cognitive and vocational interventions after mild to moderate traumatic brain injury: study protocol for a randomized controlled trial. <i>Trials</i> , 2017, 18, 483.	1.6	24
31	Biographical disruption, adjustment and reconstruction of everyday occupations and work participation after mild traumatic brain injury. A focus group study. <i>Disability and Rehabilitation</i> , 2016, 38, 2296-2304.	1.8	31
32	User involvement and experiential knowledge in interprofessional rehabilitation: a grounded theory study. <i>BMC Health Services Research</i> , 2016, 16, 547.	2.2	14
33	Vestibular Rehabilitation After Traumatic Brain Injury: Case Series. <i>Physical Therapy</i> , 2016, 96, 839-849.	2.4	35
34	Rehabilitation pathways and functional independence one year after severe traumatic brain injury. <i>European Journal of Physical and Rehabilitation Medicine</i> , 2016, 52, 650-661.	2.2	15
35	Physical and mental health 10 years after multiple trauma. <i>Journal of Trauma and Acute Care Surgery</i> , 2015, 78, 628-633.	2.1	40
36	Is radial Extracorporeal Shock Wave Therapy (rEWST) combined with supervised exercises (SE) more effective than sham rESWT and SE in patients with subacromial shoulder pain? Study protocol for a double-blind randomised, sham-controlled trial. <i>BMC Musculoskeletal Disorders</i> , 2015, 16, 248.	1.9	9

#	ARTICLE	IF	CITATIONS
37	Self-Perceived Health Care Needs and Delivery of Health Care Services 5 Years After Moderate-to-Severe Traumatic Brain Injury. <i>PM and R</i> , 2014, 6, 1013-1021.	1.6	47
38	Health Outcome after Major Trauma: What Are We Measuring?. <i>PLoS ONE</i> , 2014, 9, e103082.	2.5	53
39	Measurement Properties of the High-Level Mobility Assessment Tool for Mild Traumatic Brain Injury. <i>Physical Therapy</i> , 2013, 93, 900-910.	2.4	21
40	Identification of relevant International Classification of Functioning, Disability and Health categories in patients with shoulder pain: A cross-sectional study. <i>Journal of Rehabilitation Medicine</i> , 2013, 45, 662-669.	1.1	16
41	Health-related Quality of Life 12 months after severe traumatic brain injury: A prospective nationwide cohort study. <i>Journal of Rehabilitation Medicine</i> , 2013, 45, 785-791.	1.1	47
42	The Trajectory of Physical and Mental Health From Injury to 5 Years After Multiple Trauma: A Prospective, Longitudinal Cohort Study. <i>Archives of Physical Medicine and Rehabilitation</i> , 2012, 93, 765-774.	0.9	74
43	Returning to Work After Severe Multiple Injuries: Multidimensional Functioning and the Trajectory From Injury to Work at 5 Years. <i>Journal of Trauma</i> , 2011, 71, 425-434.	2.3	42
44	Mental Health and Posttraumatic Stress Symptoms 2 Years After Severe Multiple Trauma: Self-Reported Disability and Psychosocial Functioning. <i>Archives of Physical Medicine and Rehabilitation</i> , 2010, 91, 481-488.	0.9	31
45	Reliability and applicability of the ICF in coding problems, resources and goals of persons with multiple injuries. <i>Disability and Rehabilitation</i> , 2008, 30, 98-106.	1.8	31
46	Identification and comparison of rehabilitation goals after multiple injuries: An ICF analysis of the patients', physiotherapists' and other allied professionals' reported goals. <i>Journal of Rehabilitation Medicine</i> , 2008, 40, 340-346.	1.1	31
47	Long-Term Multidimensional Functional Consequences of Severe Multiple Injuries Two Years After Trauma: A Prospective Longitudinal Cohort Study. <i>Journal of Trauma</i> , 2007, 62, 461-470.	2.3	119
48	Return to Work After Severe Multiple Injuries: A Multidimensional Approach on Status 1 and 2 Years Postinjury. <i>Journal of Trauma</i> , 2007, 62, 471-481.	2.3	78